VOL. 34. Ser. A.

Title-page and Index. [Issued January 1948.]

THE REVIEW OF APPLIED ENTOMOLOGY.

SERIES A: AGRICULTURAL.

ISSUED BY THE IMPERIAL INSTITUTE OF ENTOMOLOGY.

LONDON: THE IMPERIAL INSTITUTE OF ENTOMOLOGY 41, QUEEN'S GATE, S.W. 7.

Price 6s. net.

All Rights Reserved.

IMPERIAL INSTITUTE OF ENTOMOLOGY.

Executive Council.

Lieutenant-Colonel J. G. ROBERTSON, Chairman, Canada.

Sir Patrick R. Laird, C.B., F.R.S.E., Vice-Chairman, United Kingdom.

N. E. HOLMES, Australia.

G. A. HOLMES, New Zealand.

A. P. VAN DER POST, Union of South Africa.

J. W. DULANTY, C.B., C.B.E., High Commissioner, Eire.

Colonel W. F. RENDELL, C.B.E., Newfoundland.

R. S. Mani, Officiating Deputy High Commissioner, India.

B. O. BINNS, O.B.E., Burma.

A. V. Hubbard, Southern Rhodesia.

J. G. Hibbert, M.C., Colonies, Protectorates and Mandated Territories. Sir Herbert Howard, Secretary.

Director and Editor.

W. J. HALL, M.C., D.Sc.

Assistant Director.

T. H. C. TAYLOR, D.Sc.

Assistant Editor.

H. S. BUSHELL, M.A.

Head Office—British Museum (Natural History), Cromwell Road, London, S.W.7.

Publication Office and Library.-41, Queen's Gate, London, S.W.7.

INDEX OF AUTHORS.

A reference in heavy type indicates that a paper by the author has been noticed.

Abdul Ghani, M., 287. Abdul Latif, 285. Acree jr., F., 359. Agacino, E. Morales, 265. Agenjo, R., 269. Aguilar, J. d', 363. Ahlberg, O., 37. Ahmad, Taskhir, 284. Aiyer, K. S. Padmanabha, 283, 287, 288. Alexander, C. P., 51. Alfaro, A., 263, 264. Alimdzhanov, R., 156. Allen, H. W., 142, 314. Allen, N., 356. Allen, T. C., 301, 311, 312, 313. Allmendinger, D. F., 50. Anderson, E. J., 74. Anderson, R. F., 158. André, M., 350. Andrewartha, H. G., 123, 324. Annand, P. N., 175. Anshi Lal Batra, 118. Anson, R. R., 166. Apple, J. W., 318. Aristov, M. T., 94. Arkhangel'skiĭ, P., 156. Armitage, H. M., 373. Arnaud, J., 11. Arthur, D. R., 46. Ashby, D. G., 43. Ashworth, J. T., 233. Asperen, K. van, 94. Atkins, I. M., 375. Atwood, C. E., 377.

Back, E. A., 194.
Bailey, S. F., 24.
Baker, E. W., 104, 137, 320.
Baker, F. T., 330.
Baker, H., 198.
Balduf, W. V., 292.
Balock, J. W., 343.
Baptist, B. A., 281.
Barber, G. W., 147.
Barber, H. S., 148.
Bare, O. S., 192, 322.

Barnes, H. F., 251. Barthel, W. F., 360.

Bartlett, B. R., 142. Bartlett, K. A., 219. Batra, Anshi Lal, 118. Batt, G. H., 320. Baudyš, E., 327. Beament, J. W. L., 61. Beard, R. L., 234. Bebbington, A. G., 167. Bégué, H., 66, 83, 84. Bellod, M., 266. Benlloch, M., 266, 267. Bennett, C. W., 379, 380. Benson, R. B., 14, 115. Benton, C., 250. Berezhkov, R. P., 326. Berger, G., 66. Berry, F. H., 233. Besemer, A. F. H., 87. Bianchi, F. A., 2, 275. Bierig, A., **128.** Birch, L. C., 21, 22, 122, 123, 188, 222, 223. Bishop, M. W. H., 114. Blair, K. G., 114. Blanchard, E. E., 104, 299. Boczkowska, M., 9, 32. Bodenheimer, F. S., 16, 256, 265. Bolwig, N., 8. Bond, W. E., 169. Bondar, G., 6. Bondy, F. F., 355. Bouhelier, R., 65, 66. Bovien, P., 8, 116, 160. Bowen, C. V., 388. Box, H. E., 102. Boyce, A. M., 112. Bozkaya, E., 48. Brande, J. Van den, **185.** Brandes, E. W., **379.** Brann jr., J. L., 197. Brannon, L. W., 138, 316, 317. Breakey, E. P., 75, 160. Breny, R., 213, 214. Brichet, J., 364. Bridwell, J. C., 64. Brimblecombe, A. R., 258. Broadbent, B. M., 80. Brodskii, A. L., 155. Bromley, S. W., 194. Bronson, T. E., 241, 346.

Brooks, A. R., 263.
Bruce, W. N., 337.
Bruel, W. E. van den, 161, 212.
Brues, C. T., 256.
Bruner, S. C., 172.
Brunn, L. K., 301, 313.
Brunson, M. H., 314.
Bruyn Ouboter, M. P. de, 99.
Bryden, J. W., 114.
Buckell, E. R., 227.
Bumgardner, R. J., 145.
Butt, F. H., 192.

Cairaschi, E. A., 348, 362. Caldwell, J., 182. Caldwell, N. E. H., 125, 126. Callan, E. McC., 57, 191. Callenbach, J. A., 180. Cameron, G. S., 166. Campbell, G. A., 327. Cañizo Gomez, J. del, 100, 265, 288. Carayon, J., 34. Carlson, J. W., 261. Carruth, L. A., 137. Carsner, E., 379, 380. Carter, G. A., 6. Carter, R. H., 388. Carter, Walter, 134, 307. Cartwright, O. L., 354. Cartwright, W. B., 4. Cassab, A., 19. Cassidy, T. P., 135, 346. Castillo, J. C., 5 320. Castro, A. Ruiz, 267, 268. Catanei, A., 10. Chaboussou, F., 229, 370. Chamberlin, F. S., 335. Chamberlin, J. C., 353. Chamberlin, W. J., 113. Chang (Yun-hwa), 120. Chappellier, A., 83. Charrot, 32. Chatterjee, P. N., 285. Chauvin, R., 350. Cheng (Tien-hsi), 344. Cheo (Ming-tsang), 119, 120. Chernuishev, P. K., 98. Chiarelli de Gahan, A., 298. China, W. E., 81. Chisholm, R. D., 341. Chiu (Shin-foon), 152. Clancy, D. W., 336. Clausen, R. T., 303. Clément, P., 363. Clemente, F. Gómez, 269. Coaton, W. G. H., 32.

Cockbill, G. F., 329.
Cole, E., 312.
Cole, J. R., 373.
Collett, R. W., 280.
Cook, W. C., 74.
Coons, G. H., 379.
Corbet, A. S., 252.
Cory, E. N., 245, 314.
Cottier, W., 209.
Cotton, R. T., 179.
Cox, J. A., 247.
Cressman, A. W., 80.
Cristóbal, U. López, 5, 58.
Cristol, S. J., 359.
Crombie, A. C., 97.
Crowley, D. J., 75.
Cutright, C. R., 202.

Dadin, J. M., 288. da Fonseca, J. Pinto, 334. d'Aguilar, J., 363. Dahms, R. G., 375. Dalbir Singh, 47. Das, G. M., 117. Daviault, L., **173.** Davis, A. C., 76. Davuidov, P. N., 326. Dean, G. A., 179, 200. de Bruyn Ouboter, M. P., 99. DeCoursey, J. D., 376. Défago, G., 327. de Gahan, A. Chiarelli, 298. de Lapparent, P., 258. del Cañizo Gomez, J., 100, 265, 288. DeLong, D. M., 267. Demidov, N. I., 154. Denne, R. W., 210. de Oliveira, S. J., **57.** De Saeger, H., **80, 81, 96, 281.** Deschapelles, J. B., 172. Devaux, L. A., 233. Dexheimer, F. J., 311, 312. Diakonoff, A., 128. Dick, R. D., 208. Dicke, F. F., **357.** Dicke, R. J., **311.** Dickson, R. C., 302. Dierick, G. F. E. M., 96. Dirks, C. O., **306.** Ditman, L. P., 241, 243. Doeksen, J., 95. Domínguez García-Tejero, F., 269. Douglass, J. R., **353.** Ducasse, G., 257, 329. Dumbleton, L. J., 210. Dunavan, D., 355.

Duprez, R., **229.** Durán M., L., **205, 206.** Dutt, G. R., 296. Duval, G., **334.**

Ebeling, W., 79, 134. Eckert, J. E., 310. Edelsten, H. M., 252. Eichmann, R. D., 140. Eigeman, L., 67. Elze, D. L., 17. El-Zoheiry, M. S., 18. Escherich, K., 69. Essig, E. O., 149, 318. Evelyn, S. H., 166. Everett, P., 21. Everly, R. T., 221. Ewer, R. F., 297. Ewing, K. P., 247, 345. Eyer, J. R., 308. Eyndhoven, G. L. van, 98.

Fahey, J. E., 160. Fallon, Baron F., 281. Fauré, G. Olalquiaga, 205. Ferrière, C., 46, 327. Feytaud, J., 258, 349. Finney, G. L., 142, 305. Fisher, E. H., 144, 313. Fisher, R. C., 116. Fisher, W. B., 151. Flanders, S. E., 305, 306, 314, 345. Fleck, E. E., 157, 388. Fleming, W. E., 304. Fletcher, F. W., 32. Fonseca, J. Pinto da, 334. Fonseka, W. T., 324. Forbes, R. S., 377. Forsslund, K. H., 130. Foster, W. R., 112. Fox, D. E., 353. Frankenfeld, J. C., 179. Franklin, H. J., 51. Fransen, J. J., 88, 91. Frazier, N. W., 227. Freiberg, M. A., 300. Frey, W., 89. Friend, A. H., 384. Friend, R. B., 193, 233. Frost, S. W., 57, 317. Furlong, J. R., 35. Furniss, R. L., 27.

Gadd, C. H., 46, 71, 324, 366. Gahan, A. B., 64. Gahan, A. Chiarelli de, 298.

Gahan, H., 300. Gaines, R. C., 303, 311. García-Tejero, F. Domínguez, 269. Gardner, J. C. M., 288. Gardner, M. W., 182. Garman, P., 49, 219, 234, 235, 237, Garrison, G. L., 311. Gates, D. R., 312. Gätzi, K., 388. Genest, M. E., 173. Gentner, L. G., 181. Gerritsen, J. D., **90.** Gersdorff, W. A., **217.** Ghani, M. Abdul, 287. Ghesquière, J., 81, 82, 83, 96. Ghilarov (see Gilyarov). Ghulamullah, 284. Giannotti, O., 333. Gilmore, J. U., 315. Gilyarov (Ghilarov), M. S., 35,156. Glass, E. H., 31, 128. Gnadinger, C. B., 352. Gobind Ram, 229. Golding, F. D., 14, 96. Gomez, J. del Cañizo, 100, 265, 288. Gómez Clemente, F., 269. Gómez-Menor, J., **269.** Goodhue, L. D., **239, 241, 243.** Gough, H. C., 367. Gould, C. J., 160. Gould, G. E., 148. Graham, C., 106. Graham, K., 377. Gravely, F. H., 291. Graves, H. W., 25. Greaves, T. G., 211, 231. Green, R. A., 87. Greenslade, R. M., 273. Greenwood, D. E., 235. Griot, M., 300. Guérin, H., 83. Gunn, D. L., 61. Gupta, B. D., 187. Gurcharn Singh Sohi, 296. Györfi, J., 39, 40. Gyrisko, G. G., 238.

Hadley, C. H., 51.
Haeussler, G. J., 138.
Hall, W. J., 46.
Haller, H. L., 157, 359, 360.
Hambleton, E. J., 249.
Hansberry, R., 303.
Hardy, C. H., 6.
Harman, S. W., 195, 248.

Haroon Khan, M., 282, 287. Harries, F. H., 376. Harrington, C. D., 133. Harrison, L. R., 358. Haseman, L., 49, 157. Hassan, A. S., 18, 19. Havelka, J., 212. Hawkins, J. H., 182. Hayes, R. A., 359. Hayward, K. J., 171, 291. Heinrich, C., 104. Heishman, R. C., 193. Helm, G. W. van der, 67. Hely, P. C., 207, 289, 383. Henderson, V. E., 329. Hendrickx, F. L., 153, 280. Herce, P., 20. Hering, M., 57. Hershberg, B. J., 127. Hetrick, L. A., 304. Hill, G. F., 211. Hill, R. E., 28, 223. Hille Ris Lambers, D., 367. Hincks, W. D., 252. Hinton, H. E., 58. Hodson, W. E. H., 42. Hoffmann, A., 362. Hoffmann, C. H., 158, 309, 311. Hofmaster, R. N., 376. Hoggan, I. A., 182. Holdaway, F. G., 275. Hollick, F. S. J., 329, 330. Hopping, G. R., 360. Horn, F. D., 227. Horton, J. R., 28, 157. Hough, W. S., 105, 195, 336. Houser, J. S., 78. Houten, J. G. ten, 59. Hsiang (Cheng-heng), 381. Hsiao (Tsai-yu), 128. Hubault, E., 214, 371. Hudault, E., 65, 66. Hukkinen, Y., 37, 64. Hull, R., 113. Hunt, G. M., 32, 380. Hurst, H., 85. Husås, O., 7. Hussein, M., 15. Hustache, A., 362. Huysmans, C. P., 93. Huzayyin, S. A., 15.

Icart, A., 300. Ikawa, M., 301. Il'inskiĭ, N. V., 154. Irreverre, F., 5. Isaac, W. E., **189.** Ivanov, E. N., 49. Ivy, E. E., **247.**

Jacks, H., 209. Jacobson, M., 359. Jameson, J. D., 166. Jancke, O., 90. Jannone, G., 154. Jarvis, H., 169. Jary, S. G., 69. Jenkins, C. F. H., 22, 23, 364, 365. Jenkins, M. T., 357. Jewett, H. H., 129, 183. Jodka, J. F. T., 238. Johnson, J., 182. Johnson, J. P., **105, 234, 237.** Jones, E. W., **74.** Jones, L. K., 74. Jones, M. A., 217. Jones, S., 283. Jones, S. C., 143. Jones, W. W., 190. Jourdan, M. L., 257.

Kanervo, V., 42, 128, 271. Kangas, E., 9, 38, 39, 40, 41, 64, 271. Kapur, A. P., 293. Karlemark, K. A., 7. Karpova, A. I., 253. Kassanis, B., 70. Kaufmann, O., 97. Keifer, H. H., 110. Kelsey, J. M., 58, 169, 210, 211. Kelsheimer, E. G., 192. Kennedy, J. S., 352. Kenworthy, A. L., 107. Kettlewell, H. B. D., 252. Khabirova, 49. Khan, A. H., 287. Khan, M. Haroon, 282, 287. King, H. E., 166. Kiplinger, D. C., 78. Kirkpatrick, T. W., 103. Kitchener, J. A., 35. Kjellander, E., 59. Klein, H. Z., 132. Knight, R. H., 61. Knight, R. L., 166. Knutson, H., 262. Koblitsky, L., 341. Koronéos, J., 326. Körting, A., 68. Krauss, N. L. H., 276. Kronenberg, H. G., 90.

Kuenen, D. J., **91.** Kuo (Kuang), **119.** Kuschel, G., **171.** Kvíčala, B., **71.**

Lal, K. B., 228, 297, 325. Lal Batra, Anshi, 118. Lambers, D. Hille Ris, 367. Lambert, R., 173, 377. Lamerson, P. G., 198, 199, 200. Landaluze, P. Urquijo, 288. Lane, M. C., 74. Lange jr., W. H., 30, 112. Langford, G. S., 245. Lapparent, P. de, 258. Larsson, S. G., 116, 271. Lathrop, F. H., 306. Latif, Abdul, 285. Latta, R., 316, 346. Läuger, P., 43, 160. Laurie, A., 78. Lavaur, J., 229. LeBeau, F. J., 182. Leech, H. B., 377. Lefèvre, P. C., **153**, 323. Lepage, H. S., 333. Lepesme, P., 96, 102, 328, 329, 361. Lepigre, A., 363. Leskinen, K., 64. Lesne, P., 328. Lespès, L., 257. Lever, R. J. A. W., 101, 366, 388. Levin, C., 315. Lewis, S. E., 128. Lieberman, F. V., 260. Lieu, K. O. V., 270. Lindgren, D. L., 150, 302. Ling (Yu), 119. Link, K. P., 301. Linsley, E. G., 149. Little jr., E. L., 57. Liu (Chi-ying), 118, 119. Liu (Tiao-hua), 151. Lizer y Trelles, C. A., 133. Lloyd, N. C., 382. Loosies, F. E., 97. López Cristóbal, U., 5, 58. López Mansilla, E. E., 299. Lovászy, P., 9, 38, 40, 64. Luginbill, P., 250. Lunde, R. N., 318.

Maan, W. J., **92, 93.**McClintock, J. A., **151.**MacDonald, D., **166.**

McDonald, J., 371. McGregor, E. A., 24. McGuffin, W. C., 159, 377. Machotin, A., 48. McKenzie, H. L., 112, 205. Mackie, D. B., 108, 110. McKinstry, A. H., 167. Madden, A. H., 335. Mallamaire, A., 165. Mamet, R., 165. Mann, H. D., 388. Mansilla, E. E. López, 299. Marchionatto, J. B., 299. Marcovitch, S., 315, 321. Markwood, L. N., 388. Marle, G. S. van, 95. Marlowe, R. H., 308. Márquez, V. Moreno, **266.** Marshall, Sir G. A. K., **14, 189, 206.** Marshall, J., 175. Martin, C. H., 142, 305. Martin, H., 43, 160. Martin, Hubert, 254. Martineau, R., 173. Martorell, L. F., 320. Marvin, P. H., **141.** Massee, A. M., 59, 101, 272. Mathers, W. G., 360. Mathur, R. N., 286. Matthée, J. J., **47.** Mayné, R., **213.** Mayne, W. W., **186.** Mendizábal, M., 268, 270. Menor, J. Gómez-, 269. Menzies, J. D., 74. Mesnil, L., 363. Meyers, P. G., 90. Meyrick, E., 269. Michelbacher, A. E., 111, 145, 309, 322.Miles, H. W., 256. Miller, D., 20. Mills, H. B., 180. Mitchell, B. L., 168, 332. Mitchener, A. V., 159. Mohamed, N., 18. Monte, O., 171. Morales Agacino, E., 265. Morcos, G., 19. Moreau, R. E., 288. Morel, G., 83. Moreno Márquez, V., 266. Moreton, B. D., 114. Moritz-Romanova, Z. E., 326. Morquer, R., 33. Morrison, H., 200. Morrison, H. E., 250, 318.

Mote, D. C., 79, 250, 318. Mott, L. O., 337. Moul, E. T., 145. Moutia, L. A., 101, 165. Muesebeck, C. F. W., 352. Müller, P., 43, 160. Muma, M. H., 245. Mungomery, R. W., 1. Munro, J. A., 29, 30, 146. Munson, S. C., 5.

Neal, P. A., 96. Negi, P. S., 164. Neiswander, R. B., 201, 202. Nel, R. G., 189. Nixon, G. E. J., 295. Norris, K. R., 162. Notley, F. B., 294. Novák, V., 328. Nysterakis, F., 33.

Olalquiaga Fauré, G., 205. Oliveira, S. J. de, 57. Ooststroom, S. J. van, 99. Orr, L. W., 337. Osburn, M. R., 238. Ouboter, M. P. de Bruyn, 99. Overholser, E. L., 50. Overley, F. L., 50, 74. Owens, C. E., 79.

Padmanabha Aiyer, K. S., 283, 287, 288. Paillot, A., 32, 160. Paker, M., 132. Paquet, G., 173. Parencia jr., C. R., 345. Parker, R. L., 190, 198, 199, 200. Pasfield, G., 384, 386. Pastrana, J. A., 371, 372. Patch, L. H., 221. Patel, J. S., 163. Pattabhiraman, T. V., 186. Paulian, R., 96, 361. Pearson, E. O., 168, 332. Peat, J. E., 167. Pelsma, C., 69. Peluffo, A. Trujillo, 104. Pemberton, C. E., 2, 170. Pepper, B. P., 137. Pesson, P., 349. Petersen, H. I., 7. Peterson, L. O. T., 377. Petherbridge, F. R., 216.

Petherbridge, M. A., 381. Petrova, G. I., 98. Pfleger, K., 212. Phillips, A. M., 345, 373. Pic, M., 329. Pinhey, E. C. G., 33. Pinto da Fonseca, I., 334. Planes, S., 268. Plank, H. K., 218, 219. Pollard, H. N., 336. Ponomarenko, D., 131. Poos, F. W., 244, 304, 338, 346. Porter, B. A., 107. Porter, C. E., 352. Pospelov, V., 131. Potts, S. F., 346. Pound, G. S., 182. Poutiers, R., 362. Powning, R. F., 256. Prebble, M. L., 377. Prentice, A. N., 167. Prentice, I. W., 182. Pruthi, H. S., 287. Puri, A. N., 296.

Questel, D. D., 136.

Rahman, K. A., 47, 117, 118, 285, 296, 297. Rainwater, C. F., 355. Ram, Gobind, 229. Ramirez, J. A., 320. Ramsey, R. C., **150**. Rasmussen, W. B., **250**, **318**. Raucourt, M., 66, 83, 84. Rawlins, W. A., 238. Rebel, H., 269. Redman, K., 30. Reeher, M. M., 251. Reeks, W. A., 377. Régnier, R., 10. Reingard, L. V., 381. Reingard, T. N. Zabud'ko-, 381. Reinhardt, J. F., 180. Reynolds, C. E., 160. Richards, O. W., 45. Richardson, C. H., 341, 342. Richmond, H. A., 377. Riedl, W. A., 358. Ris Lambers, D. Hille, 367. Ritcher, P. O., 292. Rivnay, E., 17, 369. Roark, R. C., 288. Robin, F., 229. Rock, P. J. G., 183.

Roebuck, A., 330.
Roepke, W., 67.
Rolfe, S. W., 69.
Romanova, Z. E. Moritz-, 326.
Romney, V. E., 135, 346.
Roonwal, M. L., 47.
Roos, K., 327.
Ross, D. M., 329.
Ruiz Castro, A., 267, 268.
Ruston, D. F., 166.

Saalas, U., 37, 64. Saeger, H. De, 80, 81, 96, 281. Sakimura, K., 276. Salt, G., 329, 330. Salt, R. W., 204. Sankey, J. H. P., 115. Sauer, H. F., 334. Schaffner jr., J. V., 236. Schärffenberg, B., 92. Schouteden, H., 81, 96. Schread, J. C., 234. Schuh, J., 160, 292. Scott, L. B., 340. Scott, M. H., 11. Seamans, H. L., 174, 183. Semenov, A. E., 253. Seneviratne, L. J. de S., 188. Sergent, Et., 11. Severin, H. H. P., 227. Shands, R. G., 4. Sharpless, N. E., 5. Shaw, H., 34. Shull, W. E., 204. Siegrist, H., 214. Silberman, R., 300. Silva, P., 5. Silvestri, F., 10. Simmonds, F. J., 295. Simpson, A. C., 100. Sinclair, W. B., 150. Singh, Dalbir, 47. Singh, R. N., 228. Singh Sohi, Gurcharn, 296. Singleton, H. P., 74. Sleesman, J. P., 43. Sloan, W. J. S., 288, 387. Slogteren, E. van, 99. Smallman, B. N., 139. Smee, C., 58, 62, 206, 207. Smith, C. C., 377. Smith, C. F., 347. Smith, F. F., 239, 241, 243. Smith, G. L., 311. Smith, H. S., 305. Smith, J. H., 386.

Smith, K. M., 70. Smith, R. F., 111, 322. Snapp, O. I., 335, 336. Snowball, J. G., 123. Snyder, T. E., 32, 380. Sohi, Gurcharn Singh, 296. Sokolov, N. P., 154. Solomon, M. E., 98. Sorenson, C. J., 261. Spawn, G. B., 231. Speyer, E. R., 161. Spiller, D., 210. Spoon, W., **67.** Stafford, E. M., **26.** Stammbach, W., 388. Stampfel, J., 328. Stanley, W. W., 144, 321. Stapel, C., 7. Staples, E. G., 323. Stapley, J. H., 216, 329. Starr, D. F., 320, 343, 376. Steinbauer, G. P., 226. Steiner, H. M., 140, 141, 196. Steinmetz, F. H., 226. Steinweden, J. B., 26. Stellwaag, F., 95. Stiff jr., H. A., 5, 320. Stoa, T. E., 30. Stoffels, E. H. J., 280, 323. Strickland, A. H., 372. Strickland, E. H., 146. Subrahmanyam, T. V., 207. Subramanyam, V. K., 186. Suire, J., **361.** Sun (Yun-pei), 143. Suomalainen, E., 99. Svärdson, G., 36. Swain, R. B., 343. Swanson, C., 145. Swartenbroekx, J., 185. Sweetman, H., 127. Swezey, O. H., 277. Swingle, M. C., 345. Syrjänen, V., 37.

Tams, W. H. T., 252.
Taskhir Ahmad, 284.
Tate, H. D., 28, 192, 312, 322.
Tauber, O. E., 337.
Taylor, T. H. C., 13, 277, 295.
Tejero, F. Domínguez García-, 269.
Telford, H. S., 30.
Tempère, G., 349.
Tenhet, J. N., 338.
ten Houten, J. G., 59.
Tewfik, M., 15.

Thomas, H. R., 182.
Thomas, W. A., 347.
Thompson, B. G., 247.
Thompson, W. R., 96.
Tilden, J. W., 30.
Tilemans, E., 281.
Tompkins, C. M., 182.
Tooke, F. G. C., 11, 333.
Townsend, J. F., 235.
Townsend, L. H., 128.
Trägårdh, I., 40.
Trelles, C. A. Lizer y, 133.
Trujillo Peluffo, A., 104.
Tseng (Sheng), 116.
Turner, N., 235, 236, 310.
Tuurala, O., 42.
Tydeman, H. M., 273.

Urquijo Landaluze, P., 288.

van Asperen, K., 94. Van den Brande, J., 185. van den Bruel, W. E., 161, 212. van der Helm, G. W., 67. van Eyndhoven, G. L., 98. van Marle, G. S., 95. van Ooststroom, S. J., 99. van Slogteren, E., 99. Vappula, N. A., 38, 42, 99. Vayssière, P., 329, 362. Venables, E. P., 23. Vergani, A. R., 298. Vilcot, P., 349. Villiers, A., 102, 361, 362. Vinson, J., 101. Vogel, M. A., 202. Vorhies, C. T., 262. Voûte, A. D., 69.

Wadley, F. M., 28.
Wain, R. L., 254.
Walkden, H. H., 28, 136, 341.
Walker, J. C., 182.
Wallace, C. R., 208, 382.
Wallace, P. P., 236.

Walton, R. R., 339, 340. Warner, C. L., 127. Wason, E. J., 289, 385. Watson, J. R., 146, 356. Watson, M. A., 113. Weatherley, P. E., 166. Webb, J. E., 87. Weber, A., 7. Webster, R. L., 72. Wehrle, L. P., 262. Wellington, W. G., 184, 315. Wells, W. G., **165.** West, T. F., **327.** Weston, W. A. R. D., 381. Wheeler, E. H., 246, 249. White, J. H., 330. Whitehead, F. E., 339, 340. Wigglesworth, V. B., 60. Wilford, B. H., 55. Wilkinson, D. S., 352. Wilson, A. R., 370. Wilson, F., **121, 230.** Wilson, G. F., **34, 184.** Wilson, J. D., 43. Wimshurst, F. M., 90. Winters, H. F., 218. Wolcott, G. N., 172, 320, 344. Woo (Ta-chang), 381. Wood, J., 216. Woodside, A. M., 4. Wray, D. L., **347.** Wright, D. W., 43.

Yakhontov, V., **155.** Yeager, J. F., **5, 193.** Yen (Chia-hsien), **118, 119, 121.** York, G. T., **135.** Young, M. T., **311.** Young, P. F., **195.**

Zabud'ko-Reĭngard, T. N., 381. Zaki, M., 18. Zappe, M. P., 233, 236, 237. Zeller, S. M., 160, 292. Zentmyer, G. A., 236. Zoheiry, M. S. El-, 18.

GENERAL INDEX.

In the case of scientific names, the page reference is cited only under the heading of the generic name.

When a generic name is printed in brackets, it signifies that the name is not the one adopted.

Abacarus, gen. n., 110.

Abacarus hystrix, on Elymus triticoides in California, 110.

abbreviata, Aphis (see A. rhamni).

abbreviatus, Diaprepes; Hypnus (Cryptohypnus).

abdominalis, Olla.

Aberia gardneri, Ceratitis capitata not

ovipositing on, 132.

Abies, pests of, in Canada, 263, 377, 378; Harmologa fumiferana on, in U.S.A., 176; effect of methyl bromide on, as Christmas trees, 316.

Abies alba, new Cecidomyiid on, in France, 214.

Abies amabilis, pests of, in Br. Columbia,

Abies balsamea, pests of, in Canada, 129, 173, 320.

Abies lasiocarpa, Acleris variana on, in Br. Columbia, 378.

Abies nobilis, Urocerus californicus in imported timber of, in Britain, 115.

Abies sibirica, new Torymid in seeds of, in Finland, 271.

abietella, Dioryctria.

abieticola, Gilpinia (Lophyrus).

abietina, Plemeliella.

abietis, Agevillea; Hylobius.

abruptorius, Exenterus.

Abutilon avicennae, Liorhyssus hyalinus on, in Central Asia, **155.**

abutilonea, Tetraleurodes.

abyssinia, Spodoptera.

Acacia, pests of, in India, 285, 287; Coleoptera on log of, in Nyasaland, 63. acalyptus, Matsucoccus.

Acanthopsyche, on tea in Nyasaland, 62. Acanthoscelides (see Bruchus).

Acaphylla steinwedeni, on Camellia in California, 109.

Acarus translucens, Green (see Hemitarsonemus latus).

Aceratagallia uhleri, DDT against, on potato in Nebraska, 224; arsenicals increasing infestation by, 224.

Aceria, gen. n., 110.

Aceria diospyri, sp. n., on Diospyros kaki in California, 111.

Aceria peucedani, on carrot in California, 374.

Aceria sheldoni, sprays against, on Citrus in Australia, **126, 127, 384.**

Aceria tulipae, type of genus, 111. Acetic Acid, in solid baits for Cydia pomonella, 308.

Acetone, dust impregnated with DDT in, 319; uses and effects of, in aerosols, 239, 240, 241, 242, 338.

Achaea spp. (Fruit-piercing Moths), bionomics and control of, in Nigeria, 14, 15.

achatana, Ancylis.

Acheta campestris (see Gryllus).

Achillea spp., tests of preparations of, against Tetranychus turkestani, 49. Achorutes armatus (see Hypogastrura).

Achras sapota, Conotrachelus mamillatus on, in Brazil, 6.

Achrysocharella ruforum, bionomics of, in Holland, 88.

achyloessa, Glyphipteryx. acinaciformis, Coptotermes.

Acleris variana, on conifers in Br. Columbia, 320, 377, 378, 379; bionomics and control of, 378, 379.

acletus, Scapteriscus.

Aclypea turkestanica (see Silpha). acontioides, Lyncestis (Melipotis).

Acroclisoides africanus, parasite Antestia in Uganda, 278.

Acrolepia assectella (on Allium spp.), in England, 69; in Holland, 68, 92; in Switzerland, 214, 215, 216; bionomics of, 69, 70, 92, 215, 216; measures against, 68, 70, 92, 216.

Acropyga, associated with Coccids on cacao in Trinidad, 372.

Acrostalagmus aphidum, infesting Aphids in Argentina, 299.

Acrylonitrile, in aerosol against Ephestia elutella, 339.

Actinoplagia, introduced into U.S.A. against Heliothis armigera, 179.

aculeatus, Chirothrips. acutus, Platymoideus.

Adalia bipunctata, experiment Perilitus coccinellae and, 114.

Adelges (see Chermes).

Adelphocoris lineolatus, on cotton and lucerne in Central Asia, 154, 253.

Adhesives, banding with, **95**, **116**, **259**, **357**; other methods of catching insects with, **118**, **120**, **176**, **331**.

Adiantum, Saissetia coffeae on, in Maryland, 315.

Adirus trimaculatus (see Hartigia).

adonidum, Pseudococcus.

Adoretus sinicus, DDT against, in Hawaii, 275.

adornatus, Calacarus.

adustalis, Epimorius.

adustus, Xylion.

advena, Adasverus.

Aegeria apiformis (see Trochilium).

Aegeria exitiosa (on peach), A. graefi recorded as, in Br. Columbia, 23; in U.S.A., 202, 321, 336, 347; measures against, 202, 336, 347.

Aegeria graefi, on peach in Br. Columbia,

23; characters of, 23.

Aegeria opalescens, fumigation against, on ornamental peach in California, 109.

Aegeria pictipes, on peach in Tennessee, 321.

Aegeria scoliaeformis, on birch in Finland, 41.

Aegeria tipuliformis, on currant and gooseberry in Oregon, 79.

aegopodii, Cavariella.

aeneus, Chalcodermus; Corymbites (Selatosomus); Meligethes.

Aenoplex (see Mastrus).

Aeolothrips fasciatus, possibly predacious on Thrips angusticeps in Finland, 37. aequatus, Rhynchites.

aerivora, Diplogaster.

Aerosols, uses of insecticides as, 239-244, 262, 327, 338, 379; apparatus for testing, 320; risks of, containing DDT, 96.

Aeschynomene sensitiva, practical absence of rotenone in, 218.

Aesiotes notabilis, bionomics and control of, in Queensland, 258-260.

affinis, Bruchus; Dasyneura (Perrisia). Africa, Longicorns recorded from coffee in, 102, 361.

Africa, French Equatorial, *Metamasius* sericeus on banana in, 361; Longicorns on coffee in, 361, 362.

Africa, French West, Pseudococcus njalensis on cacao in, 46; Platyedra gossypiella on cotton in, 96; new insects on other plants in, 14, 96, 188, 206; Rhizopertha dominica boring in wood in, 328; Schistocerca gregaria in, 165, 351; new parasite of Prodenia litura in, 96.

Africa, Portuguese East, parasite of Apanteles diparopsidis in, 63; Rhizopertha dominica boring in wood in, 328.

Africa, South, Jassids on cotton in, 166; Halotydeus destructor on vegetables in, 162; Coleoptera in timber in, 11, 12, 333; Hodotermes in, 32; Braconids in, 82, 296; plant quarantines in, 292; fumigation of apples exported from, against Cydia pomonella, 189; suggested importation of Anaphoidea nitens into Mauritius from, 102.

africanus, Acroclisoides.

Agalliana ensigera, food-plants and relation of, to virus disease in Argentina, 379, 380.

Agapophyta bipunctata, measures against, on beans in Queensland,

126.

agassizi, Xylocrius.

Agastache uricifolia, preferred by Anabrus simplex in California, 25.

Agathis [Conifer], Aesiotes notabilis on, in Queensland, 258.

Agathis pumila [Braconid], utilisation of, against Coleophora laricella in Nova Scotia, 130.

agavis, Chrysomphalus.

Ageniaspis fuscicollis, parasite of Prays oleellus in Greece, 326.

Agevillea abietis, gen. et sp. n., bionomics of, on Abies alba in France, 214.

Aglaope infausta, spray against, on almond in Spain, 266.

agonus, Limonius.

Agral 2, 34.

Agrilus anxius, relation of, to die-back of birch in Canada, 130.

Agrilus ater, on Populus tremula in Finland, 40.

Agrilus derasofasciatus, on Pistacia terebinthus in Jugoslavia, 327; bionomics of, on vines in France and Germany, 328.

Agrilus mendax, on Sorbus aucuparia in Finland, 41.

Agrilus viridis, on Salix caprea in Finland, 41.

Agriotes, survival of, in crop rotations, 157.

Agriotes lineatus, soil factors affecting injury by, in Germany, 92.

Agriotes mancus, on potato in Connecticut, 235.

Agriotes obscurus, soil factors affecting injury by, in Germany, 92.

Agromyza, new parasite of, on coffee in Belgian Congo, 96; key to N. American species of, 57.

Agromyza allia, sp. n., on onion in U.S. A., 57.

Agromyza flaveola, A. orbona confused with, in California, 57.

Agromyza orbona, on peas in California, 57.

INDEX. 399

Agromyza phaseoli, insecticides against, on beans in Australia, 125, 384, 386; variety of bean resistant to, 125.

Agromyza phaseolunata, sp. n., in lima beans in New Jersey, 57.

Agromyza pinguis, bionomics and control of, on chicory in Belgium, 212, 213.

Agromyza pusilla, food-plants of, in U.S.A., 57; species confused with, 57. Agromyza subpusilla, sp. n., food-plants of, in U.S.A., 57.

Agromyza virens, on celery in Hawaii, 274.

Agropyrum repens, Hylemyia coarctata on, in Britain, 368.

Agrostis alba var repens, Hylemyia

coarctata on, in Britain, 368.

Agrothereutes stokesi (parasite of Cydia spp.), introduced into New Jersey from Australia, 138, 139; bionomics and rearing of, 138, 139; not parasitising Pyrausta nubilalis, 139.

Agrotis c-nigrum (see Amathes). Agrotis nigricans (see Euxoa).

Agrotis orthogonia, starvation method of control of, on cereals in Alberta, 183.

Agrotis segetum, experiments with Beauveria bassiana and, in Russia, 131.

Agrotis tokionis, in China, 116; characters of, 116.

Agrotis tritici (see Euxoa).

Agrotis ypsilon, in China, 116; characters of, 116.

Ahasverus advena, in dried bananas in France, 363.

Aircraft, interceptions and risk of spread of pests in, 2, 162, 171, 324; spray against pests in, 2; termite mounds in landing grounds for, 366; insecticidal dusts, baits and sprays distributed from, 89, 109, 176, 177, 178, 247, 340, 369.

Ajuga reptans, Aphid on, in Holland, 367.

ajugae, Myzus.

Alabama, pests of Solidago leavenworthii in, 309.

Alabama argillacea (on cotton), bionomics of, in S. America, 300; in Barbados, 159; DDT against, in Texas, 247; fungus infesting, 300.

Alaus prosectus, predacious on Aesiotes notabilis in Queensland, 260.

Alberta, pests of field crops in, 159, 174, 183, 204; Dendroctonus monticolae on pine in, 360.

albifrons, Decticus.

albilinea, Protoleucania.

Albizzia, Sthenarus on, in Uganda, 296. albohirtum, Dermolepida.

albopictus, Elasmus.

Alchornea cordifolia, Coccid on, in Belgian Congo, 82.

Alcis verrillata (see Thamnonoma ochrifascia).

Alcohol, as solvent for DDT, 58.

Alcohols, action of primary, on blowfly larvae, 85, 86.

Alder (see Alnus).

Aleurites spp., pests of, in Nyasaland, 62, 63.

Aleurocanthus woglumi, establishment of Eretmocerus serius against, in Mexico, 179.

Aleurodes campanulae, sp. n., on Campanula in Finland, 64.

Aleurothrixus howardi, natural enemies and control of, on Citrus in Argentina, 171.

Alfalfa (see Lucerne).

Algeria, Quadraspidiotus perniciosus in, 363, 364; Schistocerca gregaria in, 10, 11.

aliberti, Eloeidiphilos (see Epimorius adustalis); Heterostylus.

allia, Agromyza.

alliorum, Taeniothrips.

Allium, suitability of species of, for Acrolepia assectella, 215. (See Leek, Onion, etc.)

Allotropa, relation of, to Cryptochetum iceryae in Belgian Congo, 82.

Allotropa burrelli, parasite of Pseudo-coccus comstocki in Virginia, 337.

Allotropa utilis, introduced into U.S.A. from Canada against Pseudococcus comstocki, 179.

Almond, Caliroa limacina on, in W. Australia, 365; Capnodis spp. on, in Tadzhikistan and Palestine, 253, 254, 369; Lepidoptera on, in Spain, 266.

alni, Dryocoetes; Cryphalus (Trypophloeus).

Alnus incana, pests of, in Finland, 9, 41.
Alocasia, Pentalonia nigronervosa on, in greenhouses in France, 362.

Alphitobius laevigatus, measures against, in stored rice in Fiji, 101.

alpina, Cephaleia.

Alsophila pometaria, on oak in Nova Scotia, 129.

Alstonia congensis, Lyctid in timber of, in Uganda, 103.

Alternaria solani, DDT ineffective against, 224.

Alumina (see Aluminium Oxide).

Aluminium Arsenate, tests of, against Leptinotarsa decemlineata, 84; nascent form of, 84.

Aluminium Fluoride, unsatisfactory against termites, 172.

Aluminium Gels, effects of, in sprays, 220, 221, 308.

Aluminium Hydroxide (see Aluminium Gels).

Aluminium Oxide, action of, on cuticle of Blatta orientalis, 60; useless as safener for cryolite, 322.

Aluminium Silicate, aluminium gels

(q.v.) formed by, 221, 308.

Aluminium Sulphate, in preparation of nascent aluminium arsenate, 84.

amabilis, Eublemma.

Amarantus (Amaranth), Agalliana ensigera breeding on, in Argentina, 380; Hymenia recurvalis on, 219, 274, 365.

Amaryllis, Trionymus on, in California,

109.

Amathes c-nigrum, in China, 116; characters of, 116.

Amauromorpha metathoracica, parasite of Schoenobius bipunctifer in China, 152.

ambiguella, Clysiana (Clysia).

Amblycerus martorelli, sp. n., in seeds of Prosopis chilensis in W. Indies, 64.

Amblymerus, parasite of Ceuthorrhynchus assimilis in Washington, 76.

Amblyteles montanus, parasite of Autographa egena in California, 30. americana, Larra; Periplaneta.

amictorius, Exenterus.

Ammonium Bifluoride, sap-stream treatment of trees with, as timber preserv-

Ammonium Copper Arsenite, sap-stream treatment of trees with, as timber preservative, 56.

Ammonium Dinitro-o-cresylate under Dinitro-o-cresol).

Ammonium Fluoride, sap-stream treatment of trees with, as timber preservative, 56.

Ammonium Oleate, in formulae for inverted spray mixtures, 50.

Ammonium Phosphate, analysis of, as bait for Dacus oleae, 268.

Ammonium Sulphamate, uses of, to destroy plants, 76, 226.

Ammonium Sulphate, manuring with, reducing infestation of jute by Apion corchori, 117; attractive to Dacus oleae, 268; in sprays against weeds, 226, 227; determination of, in commercial ammonium phosphate, 268.

Ammonium Thiocyanate, uses of, to destroy Prunus nigra, 226.

Ammothamnus lehmanni, sophocarpin prepared from, 49.

Amorpha fruticosa, toxicity of seeds of, to Epilachna varivestis, 303.

Ampeloglypter ater, bionomics and control of, on vines in Massachusetts, 52. Amphimallon majalis, morphology of,

Amphimallon solstitiale, survival of, in crop rotations in U.S.S.R., 157.

amygdali, Anuraphis (see A. schwartzi).

Amyl Acetate, effect of spray containing, on Anastrepha ludens, 376.

Amyl Salicylate, in baits for Protoparce,

Anabrus simplex, in U.S.A., 25, 109, 177, 181; bionomics of, 25, 177; baits for,

Anacardium occidentale, Othreis attacking fruits of, in Ceylon, 282.

Anagrus ovijentatus (see Anaphes).

analis, Bruchus.

Anamirta, larvae of Othreis on, in Ceylon,

Anaphes ovijentatus, parasite of Lygus hesperus in California, 346.

Anaphoidea nitens, suggested introduction of, into Mauritius against Gonipterus scutellatus, 102.

Anaphothrips, on maize in Hawaii, 276,

Anaphothrips obscurus, DDT fog ineffective against, on Citrus in Arizona, 263; on sugar-cane and Sorghum sudanense in Hawaii, 275.

Anaphothrips orchidii, spray against, on Cinchona in Porto Rico, 218.

Anaphothrips speciosus, on sugar-cane in New Caledonia, 275.

Anaphothrips swezeyi, on sugar-cane in New Caledonia, 275.

Anasa tristis (on squash in U.S.A.), bionomics of, 140; utilisation of parasite of, 72; dusts against, 72, 311, 312; resistance of, to DDT aerosols, 244.

Anastatus antestiae, parasite of Antestia in Uganda, 278.

Anastatus bifasciatus, A. antestiae recorded as, 278.

Anastatus disparis, parasite of Lymantria dispar in U.S.A., 193.

nitidulus, Anastoechus parasite Dociostaurus maroccanus in Spain, 265.

Anastrepha, quarantine treatment against, of fruit imported into U.S.A. from Mexico, 112.

Anastrepha fraterculus, cage for distributing parasites of, in Argentina, 291: bionomics of form of, in Mexico, 137.

Anastrepha ludens, investigations re-lating to control of, in mangos in Mexico, 343, 376.

anates, Euphorus.

ancilla, Othreis.

Ancistrotermes latinotus, damage to cotton by, in Tanganyika, 167. ancorifer, Psallus.

Ancylis achatana, food-plants and control of, in Britain, 185.

Ancylis comptana (fragariae), in U.S.A., 141, 142, 199, 322; bionomics and control of, on strawberry, 199; on other plants, 142; as host of parasite of Cydia molesta, 141, 142, 322.

ancylivorus, Macrocentrus.

ancylus, Diaspidiotus.

Andropogon spp., relation of Coccinella septempunctata to Aphids on, India, 228.

Anethol, effect of evaporation on baits containing, 341.

Angitia, parasite of Schoenobius bipunctifer in China, 152.

angustella, Oreopsyche.

angusticeps, Thrips

angustus, Haplothrips.

Anicetus parvus, parasite of Ceroplastes in Belgian Congo, 81.

Anilastus didymator, parasite of Taeniocampa gracilis in Finland, 42.

Anisoplia, survival of, in crop rotations in U.S.S.R., 157.

Annona coriacea, new weevil on, in Brazil, 6.

Annona muricata, Ceroplastes quadrilineatus on, in Belgian Congo, 81.

Annona squamosa, Conotrachelus bondari on, in Brazil, 6; Ceratitis capitata not ovipositing on, in Palestine, 132.

annularis, Chlorophorus.

Anobium nitidum (see Hadrobregmus).

Anobium punctatum, in S. Africa, 12, 333; bionomics of, in New Zealand, 210, 211; in structural timber and furniture, 12, 116, 211, 333; in other materials, 211; measures against, 12, 116.

Anomala, measures against, on Citrus in Costa Rica, 128.

Anomala horticola, on rye in Denmark, 116.

Anomala lucicola, experiments with Bacillus popilliae and, in U.S.A., 51. Anomala nazarena, bionomics and con-

trol of, on wheat in Palestine, 17. Anomala orientalis, on maize in Connecticut, 237; on pineapple and sugarcane in Hawaii, 135, 170;

mixture against, 135.

Anomala sulcatula, intercepted in aircraft in Pacific Is., 2.

Anomis flava (see Cosmophila).

Anomis sabulifera, measures against, on jute in Bengal, 164.

anonae, Conotrachelus.

Anoplonyx laricis, on larch in Quebec,

Antestia, importance and control of, on coffee in Belgian Congo, 280.

Antestia cincticollis, in Uganda, 277, 278; parasite of, 278.

Antestia faceta, value and use of parasites against, on coffee in Uganda, 277-279; bionomics of, 277, 278.

Antestia lineaticollis, value and use of parasites against, on coffee in Uganda, 277-279; bionomics of, 277, 278.

Anastatus; Corioxenos; antestiae, Hadronotus.

Antholcus varinervis, trap for collecting emerging adults of, in New Zealand,

Anthonomus grandis (on cotton in U.S.A.), dusts against, 178, 303, 311, 355, 356; suggested use of defoliants against, 315.

Anthonomus piri, egg stage of, on pear in France, 362.

Anthonomus pomorum, bionomics and control of, on apple in Holland and Britain, 93-95, 273, 274; on pear, 273.

Anthonomus rubi, dusts against, on strawberry in Denmark, 7.

Anthonomus signatus, on strawberry in New Brunswick, 319.

Anthophila pariana (Apple and Thorn Skeletoniser), in California, 373.

Anthores leuconotus, measures against, on coffee in Africa, 102.

anthracina, Hylemyia (Pegohylemyia).

anthracinus, Bracon.

Anthrenus coloratus, distribution of, in stored products, etc., 59.

Anthrenus fuscus, in Britain, 58. Anthrenus museorum, in Britain, 58.

Anthrenus pimpinellae, infesting insect specimens in Britain, 59.

Anthrenus scrophulariae, infesting insect specimens in Britain, 59.

Anthrenus verbasci, in Britain, 58.

Anthrenus vorax, in imported products in Britain, 59.

Anticarsia gemmatalis, bionomics of, on soy bean in Porto Rico, 218, 219. antiqua, Hylemyia (Chortophila); Orgyia (Notolophus).

Antisera, effects of, on Laphygma eridania, 193.

Ants, destroying noxious insects, 284, 379; attacking honey bees, 297; associated with Aphids, 27, 169, 238, 311, 386; associated with Coccids, 33, 46, 238, 372; stored-product pests in nests of, 150; measures and experiments against, 27, 28, 192, 238, 249, 297, 323, 328, 334; Coleoptera infesting stored dry pupae of, 212.

Ants, Carpenter (see Camponotus).

Ants, Leaf-cutting (see Atta).

Ants, Little Fire (see Wasmannia auropunctata).

Anuraphis amygdali (see A. schwartzi). Anuraphis persicae-niger, against, on peach in New South Wales, 385.

Anuraphis schwartzi, on Prunus spp. in Britain, 272.

Anuraphis tulipae, bionomics and control of, in Queensland, 169.

anxia, Lachnosterna (Phyllophaga).

anxius, Agrilus.

Aonidiella aurantii (on Citrus), in Australia, 289, 364; in California, 79, 109, 145, 176, 302, 374; in New Zealand, 21; in Persia, 16; miscellaneous food-plants of, 109, 145, 364; natural enemies and biological control of, 145, 364; fumigants against, 176, 289, 302, 364, 374; resistance to HCN in strains of, 150, 302; sprays against, 21, 79, 289, 364; DDT dust ineffective against, 80.

Aonidiella citrina, on Camellia in California, 109.

Aonidiella orientalis, on Citrus in Iraq and Persia, 16.

Apanteles, parasite of Othreis in Ceylon, 188; of Ethiopian Region, 281; parasite of Plecoptera reflexa in India, 286; Palaearctic species of, 352; review of information on, 281.

Apanteles bredoi, sp. n., parasite of Sylepta derogata in Belgian Congo, 81. Apanteles carpatus, suggested method of studying oviposition by, 315.

Apanteles congoensis, sp. n., parasite of Dichocrocis crocodora in Belgian

Congo, 80.

Apanteles diparopsidis, parasite of, in Nyasaland and Portuguese E. Africa, 63.

Apanteles epiplemicidus, sp. n., parasite of Leucoplema dohertyi in Belgian Congo, 81.

Apanteles ghesquièrei, sp. n., parasite of Eublemma spp. in Belgian Congo, 81.

Apanteles glomeratus, use of, against Pieris rapae in New Zealand, 323; parasite of P. brassicae in Russia, 131; new bacterium pathogenic to, 131.

Aparteles marginiventris, establishment of, in Hawaii, 3, 171, 274; in Texas, 3; hosts of, 3, 171, 274; life-cycle of, 3.

Apanteles risbeci, sp. n., parasite of Prodenia litura in Senegal, 96.

Apanteles ruficrus, hosts of, in China, 152, 381.

Apanteles singaporensis, 62.

Apanteles solitarius, value of, against Stilpnotia salicis in Canada, 129.

Apanteles transvaalensis, host and parasites of, in Nyasaland, 63.

Apanteles uroxys, associated with Myelois ceratoniae in Belgian Congo, 80.

Apanteles wilkinsoni, sp. n., parasite of Lamprosema indicata in Belgian Congo, 81.

Apanteles xanthostigmus, parasite of Prays oleellus in Greece, **326**.

Apantesis blakei, on wheat in Montana, 181.

Apate indistincta, on Aleurites in Nyasaland, 63.

Apateticus nigrolimbatus (see Podisus). Aphelinus diaspidis (see Aphytis).

Aphelinus mali, imported into Mexico against Eriosoma lanigerum, 179.

Aphidius, parasite of Cinara in N. Carolina, 311.

Aphidius granarius, experiments with, against Aphids in Wales, 46.

Aphidius procephali, parasite of Cinara in N. Carolina, 311.

Aphids, and plant virus diseases, 70, 71, 74, 90, 99, 102, 113, 114, 182, 206, 226, 297, 367, 370; ants associated with, 27, 169, 238, 311, 386; natural enemies of, 33, 46, 73, 111, 145, 228, 274, 297, 309, 311; effect of insecticides on reproduction of, 45, 143; classification and new species of, 149, 318, 326, 367.

Aphis abbreviata (see A. rhamni).

Aphis avenae, auct. (see Rhopalosiphum prunifoliae).

Aphis dianthi (see Myzus persicae).

Aphis fabae,* in Britain, 113, 114; in Holland, 99; on beans in Queensland, 125; and virus diseases of Narcissus and beet, 99, 113, 114; winter foodplant of, 113; measures against, 113, 114, 125.

Aphis gossypii (on cotton), sprays against, in China, 293; effects of dusts on infestation by, in U.S.A., 178, 311, 355, 356.

Aphis maidis, DDT aerosol against, on maize in greenhouses in Maryland, 240; on sorghum in Queensland, 387.

Aphis pomi, on Cotoneaster spp. in Britain, 184; on apple in Denmark, 8; insecticides against, 8, 184.

Aphis rhamni, measures against, on potato in U.S.A., 226, 243; foodplants and relation of, to potato leaf-roll, 226.

Aphis rumicis* (Bean Aphis), effects of dusts on mortality and reproduction of, in U.S.A., 143.

Aphis, Cabbage (see Brevicoryne brassicae).

Aphis, Green Peach (see Myzus persicae). Aphis, Pea (see Macrosiphum onobrychis).

^{*}In view of the divergent opinions on the identity of the Aphids of this group, no attempt is here made to show the synonymy.

Aphis, Tobacco (see Myzus persicae).

Aphis, Walnut (see Chromaphis juglandicola).

Aphis, Woolly Apple (see Eriosoma lanigerum).

Aphodius howitti, measures against, on Trifolium subterraneum in New Zealand, 324.

Aphonus castaneus, bionomics of, in turf in Connecticut, 237.

Aphthona bimaculata, on Sesamum in Nyasaland, 63.

Aphthona euphorbiae, on apple, etc., in Britain, 272.

Aphytis chrysomphali, introduced parasite of Aonidiella aurantii in W. Australia, 364.

Aphytis diaspidis, parasite of Chrysomphalus agavis in Bermuda, 190.

apicale, Megacoelum.

apicalis, Calidea dregii; Nephotettix; Trioza.

apiforme, Trochilium (Aegeria).

apii, Trephognathus.

Apion, crop rotations ineffective against, on clover in Russian Union, 157.

Apion corchori, bionomics and control of, on Corchorus spp. in India, 117, 163.

Apion ulicis, utilisation of, against Ulex europaeus in New Zealand, 323.

Apis indica, natural enemies of, in Punjab, 297.

Aplanobacter stewarti, Chaetocnema pulicaria transmitting, to maize in Maryland, 244.

Aplastomorpha, hyperparasite of Plutella maculipennis in Nyasaland, 63.

Aplastomorpha calandrae, parasite of Lasioderma serricorne in Nyasaland, 62.

aplopappi, Ephialtes (Scambus).

Apodemus sylvaticus, predacious on Diprion pini in Holland, 88.

appalachianus, Neopristomerus.

Apple, Quadraspidiotus perniciosus on, in Algeria, 364; Aonidiella aurantii on, in W. Australia, 364; pests of, in Britain, 6, 35, 115, 272, 273; pests of, in Br. Columbia, 175; pests of, in Denmark, 7, 8, 116; pests of, in Finland, 37, 42; pests of, in France, 229, 362, 370; pests of, in Holland, 67, 93, 94, 95; Eriosoma lanigerum on, in India, 228; Aphids on, in Norway, 7; Coccids on, in Persia, 16; Cydia pomonella on, in Spain, 288; Aphids on, in Sweden, 7; pests of, in U.S.S.R., 156, 254, 374; pests of, in U.S.A., 4, 31, 49, 50, 51, 52, 73, 75, 105, 106, 107, 109, 139, 175,

195-199, 202, 203, 219-221, 234, 235, 237, 245, 246, 248, 306, 308, 317, 336, 374; distinct strains of *Eriosoma lanigerum* on *Cotoneaster* and, 184; insecticides and injury to, 7, 50, 51, 52, 83, 107; problem and estimation of DDT spray deposits on, 106, 107, 160.

Apple (Fruit), Pristonychus terricola damaging, 59; effects of fumigating, with methyl bromide against pests, 107, 189, 190, 364; dried refuse from, in baits, 200, 292.

Apple Aphis, Woolly (see Eriosoma lanigerum).

Apple Leafhopper, White (see Typhlocyba pomaria).

Apple Maggot (see Rhagoletis pomonella).
Apple and Thorn Skeletoniser (see
Anthophila pariana).

Apple-blossom Weevil (see Anthonomus pomorum).

approximatus, Pissodes.

Apricot, pests of, in Central Asia, 49, 253; Melanaspis obscura on, in California, 108; Neurotoma nemoralis on, in France, 160; Capnodis tenebrionis on, in Morocco, 364; Coccids on, in Persia, 16; fumigation of fruit of, with methyl bromide, 364.

Apricot Weevil (see Rhynchites auratus). Aprostocetus, hyperparasite of Antestia in Uganda, 278.

Apterona crenulella, on Corylus in California, 374; on apple in U.S.S.R., 374.

Arabia, Schistocerca gregaria in, 15, 16. arabidis, Asterolecanium.

Araecerus fasciculatus, methyl bromide against, in banana flour in Morocco, 65.

Aralia, Halticus bracteatus on, in California, 374.

Araucaria cunninghami, pests of, in Queensland, 258, 260.

araucariae, Rioxa; Syllitus.

Arbor-vitae (see Thuja).

Arbutus, Parlatoria oleae on, in California, 373.

Archips fumiferana (see Harmologa).

Archips obsoletana (see Tortrix).

Archips rosaceana (see Tortrix).

arcticus, Phaeogenes.

Arctium lappa, Thrips nigropilosus on, in Hawaii, 277.

argentifera, Plusia.

Argentina, fruit pests in, 6, 171, 291, 298, 299; pests of field crops and vegetables in, 104, 133, 171, 298, 299, 300, 372; Agallia ensigera and virus

disease of beet, etc., in, 379, 380; Rhyacionia buoliana on pines in, 371; Phylloxera on vines in, 299; beneficial insects and biological control in, 5, 58, 104, 171, 291, 300, 372; hyperparasite of Oeketicus kirbyi in, 300; entomogenous fungi in, 171, 299, 300; beneficial insects imported into U.S.A. from, 104, 179.

argentina, Fundella.

argentinense, Calosoma.

argentinensis, Porizon.

argillacea, Alabama.

Argyresthia, bionomics and control of, on cherry in Holland, 90.

Argyresthia ephippella, 90.

Argyresthia nitidella, 90.

Argyresthia sorbiella, on Cotoneaster in Europe, 185.

Argyria, bionomics of, in sugar-cane in China, 121.

Argyria sticticraspis (see Proceras).

Aridelus spp. (parasites of Antestia), bionomics of, in Uganda, 278.

Arilus cristatus, bionomics of, in Pennsylvania, 145.

Aristotelia fragariae, on strawberry in Montana, 181.

Aristotelia subericinella, on lucerne in Tadzhikistan, 253.

Arizona, pests of Citrus and vines in, 24, 262, 263; miscellaneous pests in, 26, 27, 176, 375; pests of Pinus edulis in, **57.**

Armadillidium, DDT aerosols toxic to, in greenhouses in Maryland, 240.

armata, Hypogastrura (Achorutes).

armida, Arsenura.

Blondeliaarmigera, (Eucelatoria); Heliothis.

Armyworm, Grass (see Laphygma exempta).

Aromia moschata, on Salix caprea in Finland, 41.

Arsenic, in pastes against Melanauster chinensis on Citrus, 270: regulations restricting use of, on apple in Denmark, 8; residues of, on apples, 199.

Arsenic Acid (Pentoxide), in bait for grasshoppers, 387; sap-stream treatment of trees with solution containing, as timber preservative, 56.

Arsenura armida, food-plants and para-

site of, in Brazil, 5, 6.

Artemisia, Parnara guttata on, in China, Taeniocampa gracilis on, in Finland, 42.

Artichoke, Jerusalem (see Helianthus tuberosus).

Artocarpus, Icerya tremae on, in Belgian Congo, 82.

Artona catoxantha, parasite of, in Java,

Asclepias, Agromyza spp. on, in U.S.A.,

Ascogaster, parasite of Acleris variana in Br. Columbia, 379.

Ascogaster quadridentata, establishment of, against Cydia nigricana in Br. Columbia, 73; not spreading into Washington, 73.

Ascotis selenaria (see Boarmia).

Ash (Fraxinus), Coccids on, in California, 112, 373.

Asia, Central, cotton pests in, 49, 154; pests of lucerne in, 131, 155, 156, 253; miscellaneous pests in, 155, pests of stone fruits in, 49, 253.

asiatica, Chionaspis.

Asopus punctatus (see Eysarcoris).

asparagi, Crioceris.

Asparagus, pests of, in U.S.A., 72, 375. Aspen (Populus tremula), pests of, in

Finland, **39, 40.**

protect coconut from Asphalt, to termites, 211.

Aspidaspis oxycoccus, sprays against, on cranberry in Massachusetts, 51.

Aspidiotiphagus citrinus, parasite of Chrysomphalus agavis in Bermuda, 190.

Aspidiotiphagus lounsburyi, parasite of Coccids in Nyasaland, 63.

Aspidiotus perniciosus (see Quadraspidiotus).

aspidoprocti, Cryptochetum (Tritolestes).

Aspidoproctus congolensis, new parasite of, on Copaifera demeusei in Belgian Congo, **82.**

Assam, Apion corchori in, 117.

assectella, Acrolepia. assimilis, Ceuthorrhynchus.

associatus, Chlamydatus.

Aster, Thrips nigropilosus on, in Hawaii, 277; new Agromyzid on, in U.S.A., (See Aster and Callistephus.)

Aster buniceus var. lucidulus, Calomycterus setarius on, in Connecticut, 105. Asterolecanium arabidis, on Pittosporum

and Veronica in California, 109. Asterolecanium bambusae, introduced Coccinellids predactions on, on bamboo in Porto Rico, 219.

miliaris, Asterolecanium introduced Coccinellids predacious on, on bamboo in Porto Rico, 219.

Asterolecanium phoenicis, on date palm in Persia, 16.

Asyncrita, parasite of Hylemyia anthracina in Finland, 64.

Atanycolus heteropus, parasite of Poecilonota variolosa in Finland, 41.

ater, Agrilus; Ampeloglypter; Dermestes.

Athalia rosae (spinarum), dusts against, on turnip in Denmark, 7.

Athous obscurus (haemorrhoidalis), soil factors affecting injury by, in Germany, 92.

atlanticus, Tetranychus.

Atomus pilosus, destroying eggs of Sitodiplosis mosellana in Washington, 251. atra, Phyllotreta; Psyche (see Oreopsyche angustella).

atratus, Taeniothrips.

Atriplex, beet pests on, in France, 10.
Atriplex semibaccata, Xerophloea vanduzeei on, in California, 227.

Atropa belladonna, development of Leptinotarsa decemlineata on, 264.

Atta (Leaf-cutting Ants), fumigation campaign against, in Brazil, 334.

Atta cephalotes, measures against, damaging crops in Br. Honduras, 323; DDT ineffective against, in Ecuador, 249.

attenuatella, Cosmopteryx.

Augomonoctenus, systematic position of, 14.

Aulacophora chinensis, bionomics and control of, on watermelon in China, 344.

Aulacophora hilaris, insecticides against, on cucurbits, etc., in W. Australia, 365.

Aulacorthum solani (see Macrosiphum). Aularches miliaris, damaging coffee and Erythrina in Mysore, 187.

aurantia, Othreis.

aurantii, Aonidiella; Toxoptera.

auricularia, Forficula.

auritus, Conoderus.

auropunctata, Wasmannia.

Austracris guttulosa, in Australia, 365, 387; bait for, on sorghum, 387.

Australia, Caliroa limacina in, 364; Halotydeus destructor in, 162; pests of stored wheat in, 222, 223; parasite of Cryptochetum iceryae in, 82; parasite of Cydia. spp. introduced into New Jersey from, 138. (See also under the various States.)

Australia, South, outbreak areas of Austroicetes cruciata in, 123, 124; pests of pastures in, 324; pests of stored wheat in, 21, 122, 188.

Australia, Western, fruit pests in, 22, 23, 364, 365; grasshoppers in, 123, 124, 365; Halotydeus destructor in, 162, 163; miscellaneous pests in, 23, 365, 366; pests of stored products in, 21, 121, 365; parasites introduced into, against Aonidiella aurantii, 364. australis, Orius (Triphleps).

Austroicetes cruciata, factors affecting distribution of, in Australia, 123, 124.

Autographa (in U.S.A.), dusts against, on leguminous crops, 317; parasite of, 3.

Autographa brassicae (see Plusia).

Autographa californica (see Plusia).
Autographa egena, bionomics of, on beans in California, 30.

Autoplectrus taprobanes, gen. et sp. n., hosts of, in Ceylon, 46.

autumnalis, Musca.

avellanae, Phytoptus.

avenae, Macrosiphum.

avenae, auct., Aphis (see Rhopalosiphum prunifoliae).

Azalea, Coccids on, in California, 109; thrips on, in Hawaii, 277...

azaleae, Eriococcus.

azureus, Torymus (Callimome).

B.

bacchus, Rhynchites.

Bacillus popilliae (causing milky disease in Popillia japonica in U.S.A.), experiments on utilisation of, 234; resistance of Anomala lucicola to, 51; imported and tested against Odontria zealandica in New Zealand, 210.

Bacillus tracheiphilus, Diabrotica melanocephala transmitting, in cucurbits in

U.S.A., 148.

Bacteria (infecting insects), experiments on utilisation of, 10, 11, 51, 131, 177, 210, 234; (infecting plants), relation of insects to, 1, 148, 244, 309.

Bacterial Blight of Walnut (see Pseudo-

monas juglandis).

Bacterial Wilt of Cucurbits (see Bacillus tracheiphilus).

Bacterial Wilt of Maize (see Aplano-bacter stewarti).

Bacterium eurygastoris, sp. n., experiments with, against insects in Russia, 131.

Bacterium prodigiosum, experiments with, against Lepidopterous larvae in Russia, 131.

badia, Orgyia (Notolophus) antiqua.

Bagasse, in baits for locusts, 294, 295. Baikea minor, pests in timber of, in E. Africa, 103.

Bait-sprays, for cotton stainers, 167; for fruit-flies, 133, 276, 385; for Pegomyia hyoscyami betae, 331; for thrips, 24, 25, 78, 240, 243, 338, 383, 384, 385.

Baits, for Anabrus simplex, 25; for fruit-flies, 268, 376; for Coleoptera, 17, 23, 30, 200, 292, 324, 341, 382; for Forficula auricularia, 8; for locusts and grasshoppers, 2, 16, 111,

165, 180, 181, 286, 294, 339, 340, 387; for mole-crickets, 19, 191, 192; for moths, 15, 198, 282, 308, 322, 372; preparation and advantages of solid types of, 309; for Lepidopterous larvae, 179, 340; formulae for, 2, 23, 30, 165, 179, 191, 192, 198, 200, 282, 295, 308, 339, 340, 372, 387; distribution of, by aircraft, 109.

bajulus, Hylotrupes.

Balsam (see Abies). Bamboo, Asterolecanium spp. on, in Porto Rico, 219; factors affecting susceptibility of stored culms of, to Dinoderus minutus, 219; Coleoptera boring in manufactured, 37, 101.

bambusae, Asterolecanium.

Banana, weevils on, in W. Africa, 1, 14, 361; Othreis attacking fruits of, in Ceylon, 282; pests of, in Cuba, 172: Pentalonia nigronervosa on, in greenhouses in France, 362; thrips on, in Hawaii, 274; Heteronychus sanctae-helenae on, in New South Wales, 382; Coccid on, in Punjab, 285; (dried Fruit and Flour), insects in, 65, 363.

Banana Borer, Black (see Cosmopolites sordidus).

bancrofti, Tarsonemus.

Banding, against Anthonomus pomorum, 93, 95; against Cydia pomonella, 73. (See Adhesives.)

banian, Hieroglyphus.

Barbados, miscellaneous pests in, 159; introduced beneficial insects in, 159. barbicornis, Magdalis.

Baris cucurbitae, sp. n., on cucurbits in Kenya, 206.

Baris nigritula, sp. n., on cucurbits in Senegal, 14.

Barium Arsenate, against Leptinotarsa decemlineata, 83.

Barium Fluoride, dusting with, against Leptinotarsa decemlineata, 84.

Barium Fluosilicate, in baits, 191, 192; dusting with, 30, 84, 100, 149, 208, 319, 321, 356; spraying with, 321, 356; formulae containing, 84, 149, 321; and oil emulsion, 321; rotenone, 84; and sodium fluoaluminate (see Dutox); causing injury to plants, 149, 321, 356.

Bark-beetles, relation of, to Ceratostomella ulmi, 54, 55, 236.

Barley, Cephus pygmaeus on, in Belgium, 185; pests of, in Britain, 273, 330, 381; Sesamia vuteria on, in Morocco, 257; pests of, in U.S.A., 157, 181, 203, 250, 251, 340, 375 Hylemyia coarctata probably developing on, 368; resistance of, to

wireworms, 330; varieties of, resistant to Toxoptera graminum, 375, 376. Barriers, for Blissus leucopterus, 250. Baryconus, hyperparasite of Antestia

in Uganda, 278.

basalis, Microphanurus.

Basic Slag, dusting with, to protect crucifers from Phyllotreta, 256.

basifulva, Erycia.

basizonus, Microcryptus.

Bassus (see Agathis).

Bathyplectes curculionis, parasite Hypera variabilis in Montana, 181.

Batrachedra ledereriella, bionomics of, in France, 361.

Bean Beetle, Mexican (see Epilachna varivestis).

Bean Fly (see Agromyza phaseoli).

Beans, pests of, in Australia, 125, 126, 383, 384, 386; Conotrachelus phaseoli on, in Brazil, 6; resistance of, to wireworms in Britain, 330; Bruchus obtectus on, in Chile, 205; thrips on, in Finland, 37; pests of, in Hawaii, 275, 276; damaged by Hylobius abietis in Holland, 93; Pseudococcus citri on, in Nyasaland, 63; new Capsid on, in Peru, 128; pests of, in U.S.A., 27, 30, 54, 138, 144, 183, 236, 240, 267, 313, 317, 318, 374, 375; Dacus tryoni trapped on leaves of, 72; little attacked by Listroderes obliquus, 23; insecticides and injury to, 176, 244, 316.

Beans (Stored), Bruchids in, and their control, 65, 66, 126, 229, 329, 334, 383, 385.

Beans, Broad (see Broad Beans).

Beans, Jack (see Canavalia ensiformis).

Beans, Lima (see Lima Beans).

Beans, Long (see Long Beans).

Beans, Sword (see Canavalia gladiata). Beans, Tonga (see Dolichos lablab).

Beauveria bassiana, value and tests of. against insects in Russia, 131.

Beauveria globulifera, infesting insects in S. America, 299.

beckii, Lepidosaphes (Mytilococcus).

Beech, Lachnosterna defoliating,

Connecticut, 237.
Bees, Honey, Vespa germanica in hive of, in New Zealand, 323; effects of insecticides on, 55, 261, 310. Apis indica.)

Beet, Agalliana ensigera on, in Argentina, 379, 380; pests of, in Australia, 169, 365, 384; pests of, in Britain, 113, 114, 272, 330, 331, 370; pests of, in Denmark, 8, 116; Gryllotalpa on, in Egypt, 19; insects on, in France, 10, 362; pests of, in U.S.S.R., 131, 157, 326; pests of, in U.S.A., 57, 74, 105, 109, 353, 374; insects and virus diseases INDEX. 407

of, 113, 114, 353, 370, 374, 379, 380; varieties of, resistant to curly-top, 354; crucifer viruses experimentally transmitted to, 182; Xerophloea vanduzeei causing virus-like symptoms in, 227, 228.

Beet Leafhopper (see Eutettix tenellus). Beet Webworm (see Hymenia recurvalis). Begonia, Otiorrhynchus sulcatus on, in California, 109; pests of, in Holland, 95.

Belgium, miscellaneous pests in, 161, 185, 212-214; beneficial insects in, 185, 213.

Belladonna (see Atropa belladonna).

bellus, Conoderus.

Bemisia, native cotton resistant to leafcurl transmitted by, in Nigeria, 169; measures against, transmitting crinkle disease of tobacco in Nyasaland, 206. benedettoi, Empoasca (see E. lybica).

beneficiens, Telenomus (Phanurus).

Bengal, pests of jute and their parasites in, 117, 163, 164.

Bentonite, nicotine fixed on, 137, 197, 199, 236, 356; other uses and effects of, in sprays and dusts, 34, 51, 179, 220, 221, 234, 308.

Benzene, as solvent for DDT, 345.

Benzene Hexachloride, tests and uses of, as insecticide, 229, 255, 256, 386; γ isomer of, 256. Benzonitriles, 256.

Benzyl Alcohol, accelerating penetration of insecticide through insect cuticle, 87.

bergi, Spilochalcis. bergrothi, Helopeltis. berlesei, Prospaltella. berlesiana, Lasioptera.

Bermuda, Gnorimoschema operculella in, 103, 300; pests of Juniperus bermudiana in, 103, 190, 191; Halticus citri on Talinum in, 103; beneficial insects and biological control in, 103, 190, 191, 301.

Bermuda Cedar (see Juniperus bermudiana).

bertrandi, Hypurus.

Bessa selecta (see Ptychomyia).

betae, Pegomyia hyoscyami.

betulae, Pulvinaria. bianchii, Organothrips.

bibax, Biprorulus.

bibliophagum, Dorcatoma.

biclavis, Howardia. bicolor, Rhynchites.

bidens, Picromerus. bidentatus, Ips (Pityogenes).

bidentipes, Lobotrachelus. bifasciata, Comperiella.

bifasciatus, Anastatus; Leptomastix. bifasciculatus, Chrysomphalus.

bifida, Cerura.

bifoveolatus, Dinoderus.

bigella, Euzophera.

bigranulatus, Omotrachelus.

Bihar, Apion corchori on jute in, 117; insects associated with lac in, 164; use of Trichogramma against sugarcane borers in, 207.

bilunulata, Eupithecia.

bimaculata, Aphthona.

bimaculatus, Leptomias; Tetranychus.

binotalis, Crocidolomia.

Biological Control, of noxious plants, 33, 165, 323.

bipartitus, Scelio.

biplaga, Earias.

Biprorulus bibax, **127.**

bipunctalis, Psara (Pachyzancla).

bipunctata, Adalia; Agapophyta.

bipunctifer, Schoenobius

bipustulatus, Bracon (Microbracon).

Birch, pests of, in Canada, 129, 130, 173; pests of, in Finland, 41.

Birds, destroying honey bees, destroying noxious insects, 35, 39, 47, 102, 148, 152, 191, 379; encouragement of, in jute fields against Anomis sabulifera, 164; spread of Orthezia insignis by, 33; risk of poisoning of, by arsenicals, 83.

Biscuits, factors affecting insect damage to, **139.**

biselliella, Tineola.

bispinulus, Cryphalus (Trypophloeus).

Bittersweet (see Solanum dulcamara).

bituberculatum, Eulecanium.

bivittatus, Melanoplus.

Bixadus sierricola, on coffee in tropical Africa, 96, 102, 362; on Musanga 362; parasite of, 96; smithii, measures against, 102.

Black Leaf 10, 236, 356.

Black Leaf 40, **356.**

Black Leaf 155, 54, 195, 197, 198, 199, 202, 203, 236, 249, 356.

Black-headed Budworm (see Acleris variana).

Blackberry, Caedicia on, in New South Wales, 208; pests of, in U.S.A., 78, 142, 292.

blackburni, Chelonus.

blakei, Apantesis (Callarctia).

blancardella, Lithocolletis.

blanchardi, Parlatoria.

blanda, Systena.

Blaps, survival of, in crop rotations in U.S.S.R., 157.

Blatta orientalis, effect of inert dusts on transpiration through cuticle of, 60,

Blattella germanica, tests of sabadilla dusts on, 144.

Blissus leucopterus, measures against, on cereals in U.S.A., 250, 312.

blitealis, Noorda.

Blitophaga opaca (see Silpha).

Blondelia, parasite of Pyralids in Porto Rico, 219.

Blondelia armigera, establishment and hosts of, in Hawaii, 171, 274.

Blowflies, structure and permeability of cuticle of larvae and pupae of; 60, 85, 86.

Blowlamp, use of, against Tyrophagus lintneri, 78.

Boarmia selenaria imparata, food-plants and parasite of, in India, 285.

bobana, Eucosma.

Boehmeria nivea, Sylepta sabinusalis on, in Belgian Congo, 81.

boisduvali, Diaspis.

Bolivia, new Coccid on Cinchona in, 112. Bollworms (see Cotton Bollworms).

Bombay, spread of *Icerya purchasi* from Madras into, 287.

Bombomima grossa, predacious on noxious insects in U.S.A., 194.

Bombyx mori, Beauveria globulifera infesting, in Brazil, 299; tests of insecticides on, 91, 304.

bondari, Conotrachelus.

Bondia comonana, bionomics of, on peach in California, 108.

Books, insects damaging, 57, 291.

Borax, soil treatment with, against

plants, 226.

Bordeaux Mixture, against Epitrix cucumeris, 53; against Jassids, 267, 293; formulae for, 52, 53, 199, 235, 239, 247, 249, 294, 384; formulae containing, 52, 235, 239, 247, 249, 294, 384; and arsenicals, 53, 149, 235, 247, 249, 327, 349; as safener in lead arsenate sprays, 199; method of incorporating, in lead arsenate sprays, 204; reducing toxicity of cryolite, 208; and DDT, 106, 157, 239, 245, 247, 249, 255, 384; and Lethane B-72, 247; and nicotine, 52, 53, 54, 247, 249, 267; and oil emulsions, 21, 52, 247, 249, 294, 319, 384; as an emulsifier for oils, 294; and rotenone, 239; and Ryanex, 249; and sulphur, 384; and injury to cucurbits, 149. borealis, 1ps.

bosqi, Lissorhoptrus.

Bostrychoplites cornutus, measures against, in timber in S. Africa, 12.

Bothriophryne ceroplastae, parasite of Ceroplastes in Belgian Congo, 81.

Bothynoderes (see Cleonus).

Bolrytis, sprays against Thrips tabaci favouring, on onion in Maryland, 240. boucheanus, Dibrachys (see D. cavus). bowringi, Colaphellus.

Box (see Buxus).

Boxia khayae, parasite of, on Khaya grandifoliola in Gold Coast, 296.

Brachycolus brassicae (see Brevicoryne). Brachymeria, parasite of Gnorimoschema heliopa in Nyasaland, 62.

Brachymeria obscurata, parasite of Cosmophila flava in China, 381.

Brachynema germari (virens), on cotton in Bokhara, 154.

Brachyrrhinus (see Otiorrhynchus).

Bracon, bionomics and laboratory host of, parasitising Apion corchori in Bengal, 163; parasite of Schoenobius bipunctifer in China, 152; parasite of Cydia strobilella in Finland, 40.

Bracon anthracinus, 40.

Bracon bipustulatus, hosts of, in Africa, 82.

Bracon discoideus, possibly parasitic on Anthonomus pomorum in Holland, 94. Bracon efoveolatus, parasite of Pissodes

notatus in Finland, 39, 40.

Bracon etiellae, sp. n., parasite of
Etiella zinckenella in Belgian Congo, 81.

Bracon gelechiae, introduction of, into
Bermuda against Gnorimoschema

Bermuda against operculella, 301.

Bracon greeni (parasite of Eublemma amabilis), laboratory hosts and utilisation of, in Bihar, 164, 165.

Bracon heteropus (see Atanycolus).

Bracon recessus, B. bipustulatus recorded as, in Tanganyika, 82.

Bracon terastiae, sp. n., parasite of Terastia meticulosalis in Belgian Congo, 81.

bracteatus, Halticus.

Bran, in baits, 2, 16, 23, 25, 30, 191, 192, 294, 324, 339, 340, 387.

294, 324, 339, 340, 387.

Brassica campestris, pests on, in U.S.A., 76, 226; methods of destroying, 226.

Brassica chinensis, resistance of varieties of, to Crocidolomia binotalis in Fiji, 101.

Brassica juncea, Crocidolomia binotalis, on, in Fiji, 101.

Brassica sativus hortensis, Raphanus sativus hortensis erroneously cited as, 101, 366.

brassicae, Brevicoryne (Brachycolus); Hylemyia; Mamestra; Phaedon; Pieris; Plusia (Autographa).

Brazil, Anobiids damaging books in, 57; cotton pests in, 300, 334; miscellaneous pests in, 5, 6, 104, 333, 334; new mite on rubber in, 104; pests of tomato in, 172; parasite of Arsenura armida in, 5, 6; fungus infesting Bombyx mori in, 299; supplement to plant quarantines in, 292; Theresia claripalpis imported into Porto Rico from, 179.

409 INDEX.

Bread, in baits for Forficula auricularia, 8; Tenebrio spp. fed on, 328.

bredoi, Apanteles.

Brenthia pavonacella, on soy bean in Porto Rico, 219.

brevicaudis, Coeloides.

brevicornis, Ephialtes (Pimpla).

Brevicoryne brassicae (on crucifers), in Argentina, 300; in New South Wales, 383, 384, 385, 386; bionomics of, in Palestine, 17, 18; in Washington, 72; experimentally transmitting diseases of crucifers, 71, 182; natural enemies of, 72, 300; tests with DDT against, 45, 383, 384, 385, 386; other insecticides against, 72, 73, 144, 385, 386; effects of dusts on reproduction

Brevipalpus, status and scope of, 104. Brevipalpus obovatus, type of genus, 104. brevipes, Pseudococcus.

brevis, Kalotermes (Cryptotermes).

Bridelia ferruginea, Achaea lienardi on, in Nigeria, 14.

brillians, Harrisina.

britannicus, Dynaspidiotus.

British Columbia, pests of agricultural crops in, 73, 112, 227, 251, 319; forest pests in, 115, 320, 360, 377-379; orchard pests in, 23, 74, 175, 319; beneficial insects and biological control in, 73, 227, 379.

British Isles, cereal pests in, 46, 216, 252, 273, 367, 368, 381; sawfly on cob nuts in, 273; pests of *Cotoneaster* spp. in, 184, 185; pests of hops in, 100, 101, 273; orchard pests in, 6, 34, 101, 115, 272, 273; Carabid damaging stored apples in, 59; pests of small fruits in, 34, 101, 272, 273; pests of vegetable and root crops in, 44, 45, 69, 70, 100, 113, 114, 161, 256, 330, 370: Aphids and virus diseases of lettuce and beet in, 70, 113, 114, 370; investigations on wireworms in, 329, 330; indigenous and imported species of Anthrenus in, 58; Siricids in imported timber in, 115; beneficial insects in, 45, 115, 331, 332; utilisation of Aphidius granarius against Aphids in, 46; parasite of Coccinellids in, 114.

Broad Beans, Bruchids in stored and field, in Morocco and Chile, 65, 205.

Broccoli, Murgantia histrionica on, in Tennessee, 144; viruses in, 182.

Bromoform, aerosol of, 339.

Bromus, relation of Oria musculosa to, 252.

Bruchophagus gibbus, bionomics of, on lucerne in Tadzhikistan, 253.

Bruchus, effects of mercury vapour on, 297.

Bruchus affinis, measures against, in stored pulses in India, 325.

Bruchus analis, bionomics and control of, in stored pulses in India, 229, 325.

Bruchus chinensis, in India, 325; bionomics of, in Queensland, 126; measures against, in stored pulses,

obsoletus, methyl bromide Bruchus against, in stored beans in Morocco, 65.

Bruchus obtectus, DDT against, in stored beans in Brazil and Australia, 334, 383, 385; food-plants of, in Chile, 205.

Bruchus pisi (see B. pisorum).

Bruchus pisorum (on peas), in Montana, 181; in Morocco, 65; in U.S.S.R., crop rotations ineffective 157; 157; against, methyl bromide against, in stored peas, 65.

Bruchus rufimanus, methyl bromide against, in broad beans in Morocco,

65.

brumata, Operophtera (Cheimatobia).

brunnea, Eucolaspis.

brunneus, Heterobostrychus; Lyctus.

bubalus, Ceresa.

Buckthorn (see Rhamnus).

Budworm, Black-headed (see Acleris variana).

Buffalo Treehopper (see Ceresa bubalus).

marinus (destroying noxious insects), establishment and bionomics of, in Hawaii, 170; in Queensland, 1, 2; in Trinidad, 191.

Building Timbers, Coleoptera in, 11, 12, 13, 116, 333; termites in, 49, 116, 366.

buoliana, Rhyacionia.

Bupalus piniarius, experiments with Beauveria bassiana against, on pine in Russia, 131.

Bur Clover (see Medicago hispida).

Burma, Euplectrus parvulus in, 285; Anthrenus vorax imported into Britain in clothing from, 59.

burrelli, Allotropa.

Bursera simaruba, used in tests of preservatives against termites, 172, 345.

buscki, Perissopterus.

Butyl Phthalate, as solubiliser for derris extractives in oils, 80; causing defoliation of orange, 80.

Butyl Thiocyanate, tests of, against pests of stored tobacco, 339.

Buxus sempervirens var. suffruticosus, Coccid on, in California, 109.

Byturus urbanus (tomentosus, auct.), on loganberry in Britain, 34; on raspberry in Denmark, 7; tests of insecticides against, 7, 34.

c-nigrum, Amathes (Agrotis).

Cabbage, Hellula phidilealis on, in Argentina, 372; pests of, in Australia, 231, 382, 383, 384, 385, 386; pests of, in China, 844; Pieris spp. on, in Denmark, 7; new weevil on, in Eritrea, 14; Lepidoptera, on, in Fiji, 101; Lepidoptera on, in Finland, 100, 271; Plutella maculipennis on, in New Zealand, 209; P. maculi-pennis on, in Nyasaland, 63; Brevicoryne brassicae on, in Palestine, 17; Hylemyia brassicae on, in Spain, 266; mole-crickets damaging, in Trinidad, 192; pests of, in U.S.A., 72, 73, 75, 76, 182, 236, 240, 313, 318, 319, 337, 355; Aphids and virus diseases of, 182; Myzus persicae ovipositing on, 367; fumigation of, with methyl bromide, 76.

Cabbage, Chinese (see Brassica chin-

ensis).

Cabbage Aphis (see Brevicoryne brassicae).

Cabbage Bug, Harlequin (see Murgantia histrionica).

Cabbage Maggot (see Hylemyia brassicae).

Cabbage Moth (see Plutella maculipennis).

Cabbage Seed-pod Weevil (see Ceuthor-

vhynchus assimilis).

Cables, Electric, Chlorops causing short circuits in, 286; Lepidoptera damaging lead sheathing of, 271, 272.

Cacao, pests and disease of, in Africa, 46, 102; pests of, in Brazil, 5; Dacus passiflorae on, in Fiji, 101; pests of, in Trinidad, 57, 372.

Cacoecia (see Tortrix).

cacti, Chilocorus.

cadaverinus, Dermestes (see D. ater).

Cadelle (see Tenebroides mauritanicus). Cadmium Compounds, protecting wood from termites, 172.

Caedicia spp., bionomics and control of, on fruit trees in New South Wales, 208.

Cages, for distribution of parasites of Anastrepha, 291.

Cajanus cajan (indicus), Heliothis armigera on, in Tanganyika, 167.

Calacarus adornatus, on Camellia in California, 109.

Calandra, natural sources of, 149.

Calandra granaria (in stored grain), in Australia, 230, 385; in Manitoba, 320; preferring large grains for oviposition, 297; DDT against, 385; effects of mercury vapour on eggs of, 45; parasite of Apion corchori reared on, 163.

Calandra oryzae (in stored cereals), in Australia, 21, 188, 222, 223, 230, 365, 385, 388; in S. Carolina, 354, 355; in China, 120; in Fiji, 101; in India, 296, 325; in Manitoba, 320; in Morocco, 65; in macaroni and spaghetti, 365; on field sorghum, 388; strains of, 21, 22, 188, 222, 230; effects of temperature and humidity on, 21, 22, 122, 188, 222, 223; effect of density on oviposition by, 223; DDT against, 333, 334, 355, 385; fumigants against, 65, 101, 342, 343, 354, 365; other measures and experiments against, 120, 296, 325, 365.

calandrae, Aplastomorpha. Calanthe (see Orchids).

Calcid, 289.

calcitrator, Collyria.

Calcium Arsenate, 270; against cotton pests, 178, 179, 180, 247, 293, 303, 311, 345, 355, 356, 381; favouring infestation of cotton by Aphis gossypii, 178, 311, 356; against pests of crucifers, 72, 73, 75, 76, 209, 231, 313; against pests of cucurbits, 149, 337; against pests of graminaceous plants, 207, 340; against Lepidoptera on jute, 164; against orchard pests, 50, 196, 254, 349; against Lygus on Parthenium, 135; against pests on potato, 53, 72, 83, 84, 100, 238, 327, 358; against Heliothis armigera on tomato, 178, 354; in baits for mole-crickets, 191; soil treatment with, against Paria canella, 78; formulae for, in dusts, 72, 73, 135, 149, 164, 231, 247, 313, 337, 340, 345, 354, 355, 358; carriers for, 149, 164, 231, 303, 354; aeroplane dusting with, 340; spraying with, 50, 51, 53. 83, 84, 100, 149, 164, 196, 209, 238, 293, 327, 349, 381; and other arsenicals, 340, 355; and copper compounds, 53, 149, 327, 349; and DDT, 157, 345; and nicotine, 73, 178, 311, 356; "inverted" mixtures of oils and, 50, 51; and rotenone, 356; and sulphur, 135, 180, 303, 345, 358; and thiocyanate, 356; and zinc sulphate, 50, 51, 196; and injury to plants, 50, 51, 83, 84, 209, 270; effect on plants of soil containing, toxicity of residues of, to domestic animals, 340, 341; commercial forms of, 84; determination of lime in, 388.

Calcium Arsenite, effect of, on Heliothis

armigera and cotton, 179.

Calcium Cyanamide, soil treatment with. ineffective against Anomala nazarena.

411

17; toxicity to insects of dusts of, 85, 89; use of, to defoliate cotton, 178, 315; mixtures of insecticides

and, 89, 315.

Calcium Cyanide, forms of, for fumigating Citrus against Aonidiella aurantii, 289, 290; as a greenhouse fumigant against Thrips tabaci, 240; treatment of nests of wasps and ants with, 229, 323.

Calcium Dinitro-o-cyclohexylphenolate (see under Dinitro-o-cyclohexyl-

phenol).

Calcium Phosphate, useless as safener

for cryolite, 322.

Calidea dregii var. apicalis, bait-spray against, on cotton in Tanganyika, 167.

California, pests of Citrus in, 24, 25, 79, 80, 112, 145, 176, 302, 374, 375; new Coccid on Fraxinus in, 112; pests of leguminous crops in, 25, 30, **57**, **111**, **177**, **227**, **322**, **374**; pests of nut trees in, 108, 145, 309, 374; orchard pests in, 24, 108, 109, 110, 111, 149, 305, 373, 374, 375; pests of Parthenium argentatum in, 135, 149, 318, 346; Cerambycids in Sequoia in, 30, 31; pests of vegetable and root crops in, 25, 30, 57, 109, 134, 149, 177, 374, 375; vine pests in, 109, 112, 227; Anabrus simplex and grasshoppers in, 25, 109, 176; miscellaneous pests in, 26, 108-111, 373-375; beneficial insects and biological control in, 30, 31, 110, 111, 145, 206, 305, 309, 345, 346, 373, 374; new Coccid intercepted on orchids in, 112; Idiopterus formosanus imported into Hawaii from, 274; pests intercepted in Hawaii in aircraft from, 2; beneficial insects introduced into Bermuda from, 301; suggested introduction of parasite of Saissetia oleae into Chile from, 206.

California Red Scale (see Aonidiella

aurantii).

californica, Cerosipha (see C. subterranea); Plusia (Autographa). californicus, Limonius; Urocerus.

Caliroa limacina, measures against, in Australia, 364, 365; bionomics of, in Britain, 272; in France, 160; on fruit trees, 160, 272, 364, 365; other food-plants of, 365.

Calla Lily (see Zantedeschia).
Callarctia blakei (see Apantesis).

Callidium pallidum, bionomics of, in California, 30, 31.

Callimome (see Torymus).

Calliphora erythrocephala, transpiration through cuticle of pupa of, 60. (See Blowflies.)

Calliptamus italicus and var. marginellus, in Spain, 268.

Callistephus chinensis, Xerophloea vanduzeei causing disease-like symptoms in, 227, 228.

Callitris intratropica, resistance of timber of, to termites in Australia, 366. callosus, Sitona.

Callyntrotus hystrix (see Abacarus).

Calomel (see Mercurous Chloride).

Calomycterus setarius, bionomics and control of, in Connecticut, 105.

Calonota festiva, experimentally parasitised by Neoaplectana glaseri in New Zealand, 210.

Calosoma argentinense, imported into U.S.A. against armyworms, 179.

Calosoma sycophanta, predacious on Lymantria dispar in U.S.A., 193.

Calotermes (see Kalotermes).

calyptrata, Helina.

Calystegia (see Convolvulus).

Camellia, fumigation of, against Coccids in California, 109.

camelliae, Hemiberlesia (see H. rapax); Lepidosaphes; Parlatoria.

camelus, Conotrachelus.

Cameroons, Longicorns on coffee in, 361, 362.

Camrula pellucida, measures against, in Br. Columbia, 227; fungous disease of, 227.

Campanula, new Aleurodid on, in Finland, 64.

campanulae, Aleurodes.

campestris, Gryllus (Acheta); Lygus. Camphor (see Cinnamomum camphora).

Camponotus, associated with Pseudococcus njalensis in W. Africa, 46; importance and control of, in houses, etc., in Oregon, 27; measures against, infesting bee-hives in Punjab, 297.

Camponotus herculeanus modoc, in Oregon, 27.

Camponotus herculeanus pennsylvanicus, bibliography on, 128.

Camponotus laevigatus, in Oregon, 27. Camponotus maculatus vicinus, in Oregon, 27.

Camptobrochis punctulatus (see Deraeocoris).

Canada, forest pests in, 159, 263, 377;

Tyrophagus lintneri on mushrooms in, 76; Leptinotarsa decemlineata on potato in, 264; Cephus cinctus on wheat in, 146, 319; pests of stored cereals in, 320; Tachinids in, 263; Sirex juvencus imported into Britain in spruce from, 115; beneficial insects imported into U.S.A. from, 179. (See also under the various Provinces.)

canadensis, Epochra.

Canavalia ensiformis, pests of, in Queensland, 125, 126.

Canavalia gladiata, pests of, in Queensland, 125, 126.

canella, Paria.

Canna, Chilo loftini on, in Arizona, 375. Cannabis sativa (see Hemp).

Cantaloupe (see Melon).

Canthium, Antestia faceta on, in Uganda, 278.

Canton Island, pests intercepted in aircraft at, 2.

canus, Limonius.

Capeweed (see Cryptostemma calendulaceum).

capitata, Ceratitis.

Capitophorus fragariae, virus diseases of strawberry transmitted by, 90.

Capitophorus ribis, on currant and gooseberry in Oregon, 79.

Capnodis, measures against, on fruit trees, 254, 364.

Capnodis carbonaria, bionomics of, on fruit trees in Tadzhikistan and Palestine, 254, 369.

Capnodis cariosa, study on oviposition

by, in Palestine, 369.

Capnodis tenebricosa, in Tadzhikistan, 254; bionomics of, on fruit trees, 254, 364, 369.

Capnodis tenebrionis, in Palestine, 369; in Morocco, 364.

Capsicum, 380; thrips on, in greenhouses in Maryland, 240; Empoasca decipiens on, in Spain, 267; Ceratitis capitata ovipositing on, 132; Leptinotarsa decembineata not feeding on, 264.

Caraway, Anthrenus coloratus in seed of,

in Sudan, 59.

Carbitol (see Diethylene Glycol Monoethyl Ether).

Carbolic Acid (see Phenol).

Carbolineum, types of, 7, 90. (See Tar

Distillates.)

Carbon Bisulphide, 87; fumigation of ants' nests with, 323, 334; soil fumigation with, against Coleoptera, 1, 20, 100, 162, 201, 327; against Zeuzera pyrina, 35; against pests of stored products, 101, 230, 342, 343, 365, 388; fumigation of bee-hives with, against Galleria mellonella, 297; revival of insects exposed to, 343; emulsion of, 201.

Carbon Dioxide, and methyl bromide, fumigation of fruit with, against Quadraspidiotus perniciosus, 363, 364.

Carbon Tetrachloride, alone or in mixtures, fumigation with, against pests of stored products, 180, 230, 320, 342, 343; action of, on insects, 343.

carbonaria, Capnodis. carcharias, Saperda. Cardamom (see *Elettaria cardamomum*). Cardboard, *Anobium punctatum* attacking, **211.**

cardinalis, Dysdercus. caricis, Vesiculaphis.

caridei, Parexorista.

carinatus, Cheiloneurus.

cariosa, Capnodis.

Carnation, Tetranychus on, in greenhouse in Massachusetts, **54.** (See Dianthus.) carolina, Cinara.

Carolina, North, pests of forest and shade trees in, 158, 233, 311; Hypogastrura armata in, 347; Aegeria exitiosa on peach in, 347; tobacco pests in, 178, 315, 347; beneficial insects in, 311, 347.

Carolina, South, Tetralopha scortealis on Lespedeza in, 304; miscellaneous pests in, 354, 355; Aphids on Pinus spp. in, 311; pests of Solidago leavenworthiin, 309; tobacco pests in, 178, 356; beneficial insects and biological control in, 311, 354.

carpatus, Apanteles; Tetrastichus.

Carpenter Ants (see Camponotus).

Carpets (see Textiles). Carpocapsa (see Cydia).

carpocapsae, Mastrus (Aenoplex).

Carpodiplosis papaveris, gen. et sp. n., bionomics of, on poppy in Sweden, 59. Carpophilus dimidiatus, in dried bananas in France, 363.

Carpophilus hemipterus, methyl bromide against, in dried figs in Morocco, 65; on sorghum in Queensland, 388.

Carpophilus obsoletus, in dried bananas in France, 363.

Carrot, pests of, in Australia, 169, 383, 384; Psila rosae on, in Britain, 44; P. rosae on, in Br. Columbia, 319; Trioza apicalis on, in Denmark, 116; Gryllotalpa on, in Egypt, 19; Epermenia chaerophyllella on, in Finland, 38; P. rosae on, in Germany, 68; P. rosae on, in Germany, 68; P. rosae on, in Switzerland, 43; pests of, in U.S.A., 250, 374; (chopped), in bait for Listroderes obliquus, 28.

Carrot Rust Fly (see *Psila rosae*).

Carthamus tinctorius, pests of,

Uzbekistan, 155.

caryaefoliae, Melanocallis. caryana, Enarmonia.

caryophyllacearum, Myzus.

casei, Piophila.

Casein, in food for Anastrepha fraterculus, 138; in spreaders and adhesives for sprays, 175, 220, 221.

Cashew (see Anucardium occidentale).
Cashew-nut Shell Oil, ineffective against
Xylotrechus quadripes, 187.

Casimiroa edulis, Ceratitis capitata on, in Palestine, 133.

Cassava, damaged by Atta cephalotes in Br. Honduras, 323.

Cassia didymobotrya, Icerya purchasi on, in India, 287.

cassiae, Pachymerus.

Cassida nebulosa, crop rotations ineffective against, on beet in U.S.S.R.,

castanea, Diparopsis.

castaneiceps, Spatulifimbria. castaneum, Tribolium.

castaneus, Aphonus.

Castor, Aonidiella aurantii on, in W. Australia, 364; tests of sprays containing extracts of seeds of, 53.

Casuarina, new weevil on, in Annam, 362.

casuarinae, Sepiomus.

Catorama herbarium, damaging books in Brazil, 57; boring in bamboo in Fiji, 101; DDT against, 57.

catoxantha, Artona.

Cattleya (see Orchids).

caudatus, Ephialtes; Torymus (Calli-

mome).

Cauliflower, Hylemyia cilicrura on, in California, 375; pests of, in New South Wales, 208, 382, 384; Plutella maculipennis on, in Nyasaland, 63; mole-crickets attacking, in Trinidad, 192: Aphids and virus diseases of, 71, 182.

cautella, Ephestia.

Cavariella aegopodii, insecticides against, on carrot in New South Wales, 383. cavus, Dibrachys.

Cecidomyiids, books on, 251.

Cecidophyes, status of, 110.

Cecidophyes vermiformis, associated with Phytoptus avellanae on Corylus in Oregon, 111.

Ceiba pentandra, Arsenura armida on, in Brazil, 6; Pseudococcus njalensis on,

in Gold Coast, 46.

Celery, pests of, in Hawaii, 274; Lygus campestris on, in Massachusetts, 53; Listroderes obliquus on, in Queensland, 387; new weevil on, in Senegal, 189.

Cellophane, use of, for obtaining oviposition by Cydia molesta, 234.

Celtis prantlii, Achaea faber on, in Nigeria, 14.

Cephaleia alpina, on larch in Holland, 69.

cephalonica, Corcyra.

Cephalosporium lecanii (infesting Coccids), in Argentina, 299; utilisation of, in Russia, 131.

cephalotes, Atta.

Cephus cinctus (on cereals), in Canada, 146, 174, 319; in U.S.A., 146, 181; on flax, 146, 181; effects of climate

on, 174; duration of development of, 181: cultural measures against, 146. Cephus pygmaeus, parasite and control

of, on cereals and grasses in Belgium,

Ceratitis capitata, in W. Australia, 22; in France, 23, 160; in Palestine, 132; host fruits of, 23, 131-132, 160; bionomics and control of, 23, 131–132. Ceratonia, Pseudococcus filamentosus on,

in Hawaii, 274.

ceratoniae, Myelois.

Ceratostomella ulmi, relation of Scolytids to, in elm in U.S.A., 54, 55, 236. Cercospora, 338.

cerealella, Sitotroga. cerealium, Limothrips.

Cereals, Oria musculosa on, in Britain, 252; pests of, in Canada, 112, 183; Limothrips cerealium on, in Europe, 374; Chilo loftini on, in Mexico, 375; Laphygma exempta on, in Nyasaland, 63; pests of, in U.S.S.R., 157, 326; pests of, in U.S.A., 77, 312. (See Maize, Wheat, etc.)

326, 342, 365, 388; in baits for mole-

crickets, 191, 192.

Ceresa, 317.

Ceresa bubalus, sprays against weeds and, in peach orchards in Pennsylvania, 141.

cerina, Glyptina.

Ceromasia, new species of, in Canada,

Ceromasia senilis, auct. (see Lydella stabulans grisescens).

ceroplastae, Bothriophryne.

Ceroplastes destructor, spray against, on Citrus in New Zealand, 21; Bothriophryne ceroplastae associated with, on coffee in Belgian Congo, 81.

Ceroplastes quadrilineatus, parasite of, on Annona muricata in Belgian Congo,

Ceroplastes rusci, on fig in Iraq, 16.

Ceroplastes sinensis (on Citrus), sprays against, in New Zealand, 21; use of Cephalosporium lecanii against, in Russia, 131.

Cerosipha, key to species of, 149.

Cerosipha californica (see C. subterranea). Cerosipha subterranea, synonymy of, 318; food-plants of, described as C. californica, sp. n., from California and Hawaii, 149.

Cerotoma ruficornis, on soy bean in Porto Rico, 219.

certus, Myzus.

Cerura bifida, damaging lead sheathing of telephone cable in Denmark, 271.

cervina, Thosea.

cervinatus, Oxycanus.

Ceuthorrhynchus assimilis, bionomics and control of, on crucifers in Washington, 72, 76.

Cevadine, toxicity of, to insects, 302.

Cevine, not toxic to insects, 302.

Ceylon, fruit-piercing moths in, 188, 281-282; miscellaneous pests 188; pests of tea in, 71, 211, 324, 325, **366**: beneficial insects and biological control in, 46, 47, 82, 188, 282, 324.

Ceylon Gooseberry (see Aberia gardneri).

chaerophyllella, Epermenia.

Chaetocnema concinna, crop rotations ineffective against, on beet U.S.S.R., 157.

Chaetocnema pulicaria, spray against, on maize in Maryland, 244; bacterial wilt transmitted by, 244.

Chaetodacus (see Dacus).

Chaetogaedia monticola, parasite of Autographa egena in California, 30.

Chaetophleps setosa, bionomics of, parasitising Diabrotica melanocephala in U.S.A., 148.

chalcites, Plusia.

Chalcodermus aeneus, dusts against, on cowpeas in S. Carolina, 355.

Chalepus dorsalis, Reduviid predacious on, in Pennsylvania, 145.

Chalepus scapularis, 27.

Changa (see Scapteriscus vicinus).

Charaeas graminis, fungus infesting, on grasses in Finland, 271.

Charips, parasite of Cinara in N. Carolina, 311.

Chauliognathus pennsylvanicus, dacious on Diabrotica melanocephala in Indiana, 148.

Cheiloneurus carinatus, parasite Pseudococcus citri in Nyasaland, 63.

Cheimatobia brumata (see Operophtera). Chelonella (see Chelonus).

Chelonus, parasite of Gnorimoschema heliopa in Nyasaland, 62.

Chelonus blackburni, ovipositing in eggs of Syrphid in Hawaii, 274.

Chelonus phthorimaeae, introduction of, into Bermuda against Gnorimoschema operculella, 301.

Chelonus rufoscapus, 62.

Chelonus texanus, bionomics of, in Texas,

Chenopodium, beet pests on, 10, 380.

Chermes piceae, on Abies balsamea in Nova Scotia, 129.

Chermes pinifoliae, on conifers in Nova Scotia, 129.

Cherry, pests of, in Britain, 35, 115; pests of, in France, 160, 370; pests of, in Holland, 67, 90; pests of, in U.S.A., 24, 75, 108, 143, 245, 374; not injured by salt of dinitro-o cyclohexylphenol, 52.

Cherry Fruit-flies (see Rhagoletis cingulata and R. fausta).

Cherry Plum (see Prunus divaricata). Chestnut, Anthrenus coloratus on, in

India, 59. Chevalierella elaeidis, gen. et sp. n., on

Elaeis guineënsis in Belgian Congo, 83. Cheyletus, predacious on other mites on Juniperus in Bermuda, 191.

Chick Pea (see Cicer arietinum).

Chicory, Agromyzids on, in Belgium,

Chile, miscellaneous pests in, 205, 206, 352; beneficial insects and biological control in, 206, 352.

Chilo loftini, food-plants of, in U.S.A.

and Mexico, 375.

Chilo oryzae, possibly identical with C. simplex, 269.

Chilo simplex, on rice in Spain, 269; C. oryzae possibly identical with, 269. Chilo zonellus, bionomics of, in Punjab,

117, 118. Chilocorus cacti, predacious on Asterolecanium spp. in Porto Rico, 219.

Chilocorus stigma, killed by DDT sprays in Florida, 238.

Chilosia illustrata, on parsnip in Finland,

China, Melanauster chinensis on Citrus in, 270; cotton pests in, 120, 293, 381; Coleoptera on crucifers in, 118, 119, 344; pests of rice in, 119, 151, 152, 293; Argyria in sugar-cane in, 121; Aulacophora chinensis on watermelon in, 344; Locusta migratoria manilensis in, 293; morphology of injurious cutworms of, 116; new Coccid in, 205; beneficial insects in, 152, 381; establishment of Aphytis chrysomphali in W. Australia from, 364; danger and prevention of introduction of pests into Hawaii by aircraft from, 2.

China Clay, preparation containing DDT and, 34.

Chinch Bug (see Blissus leucopterus).

chinensis, Aulacophora (Raphidopalpa); Bruchus; Melanauster; Parlatoreopsis (Parlatoria).

Chinese Cabbage (see Brassica chinensis). Chinese Radish (see Raphanus sativus hortensis).

Chionaspis asiatica, on plum in Persia, 16.

Chionaspis etrusca, on Tamarix California, 109.

Chionaspis longispina (see Parlatoreopsis chinensis).

Chionaspis pinifoliae, on Pinus edulis in U.S.A., **57.**

415 INDEX.

Chionaspis wistariae, on Wistaria in California, 109.

Chirothrips aculeatus, on sugar-cane in New Caledonia, 275.

Chirothrips spiniceps, on maize Hawaii, 276.

Chlamydatus associatus, insecticides against, on potato in Nebraska, 224.

Chlordifluoromethane, use of, for producing aerosols, 241.

chlorea, Sphingomorpha. Chloridea (see Heliothis).

α-Chlornaphthalene, as solvent for DDT, 157.

Chlorophorus annularis, in bamboo ski sticks in Finland, 37.

Chlorops, studies on, causing short circuits on high tension cables in India, 286.

Chlorpicrin, fumigation with, against pests of stored grain and flour, 180, 320, 354; and carbon tetrachloride, 320; methods of soil treatment with, against pests, 49, 76; soil treated with, increasing resistance of pineapple to mealybug wilt, 135.

Chortoicetes terminifera, bait for, on sorghum in Queensland, 387.

Chortophila (see Hylemyia).

Chromaphis juglandicola, effect of insecticides on natural enemies and control of, on walnut in California, 145, 309.

Fluoride, unsatisfactory Chromium

against termites, 172.

Chrysanthemum, pests of, in greenhouses in U.S.A., 54, 240; spray causing injury to, 54.

Chrysanthenum cinerariaefolium, Thysanoptera on, in Nyasaland, 63.

Chrysobothris mali, on currant and

gooseberry in Oregon, 79.

Chrysocharis laricinellae, utilisation of, against Coleophora laricella in Nova Scotia, 130.

Chrysomela (Chrysolina) hyperici, utilisation of, against Hypericum perforatum in New Zealand, 323.

chrysomphali, Aphytis.

Chrysomphalus agavis, parasites and control of, on Juniperus bermudiana in Bermuda, 190, 191.

bifasciculatus, Chrysomphalus plants of, in California, 109, 112.

Chrysomphalus dictyospermi, in Maryland, 315; in Persia, 16; in Spain, 268, 269; on apricot, 16; on Citrus, 16, 268, 269; on orchid, 315; sprays against, 268, 315.

Chrysomphalus vossi (see Linding-

aspis).

Chrysopa ventralis, predacious on Neodiprion sertifer in Sweden, 130.

chrysorrhoea, auct., Euproctis, Porthesia (see Nygmia phaeorrhoea).

Cicer arietinum (Chick Peas), Bruchus obtectus in, in Chile, 205; (stored), B. analis in, in India, 229; fumigation of, with methyl bromide, 65, 66.

Cidaria fluctuata, on turnip in Finland,

Cigarette Beetle (see Lasioderma serricorne).

cilicrura, Hylemyia.

Cinara spp., on Pinus spp. in U.S.A., 311; natural enemies of, 311.

Cinchona, pests of, in Belgian Congo, 153, 154, 323; new weevil on, in Nigeria, 206.

Cinchona calisaya, new Coccid on, in

Bolivia, 112.

Cinchona calisaya var. ledgeriana, Helopeltis spp. on, in Nyasaland and Belgian Congo, 63, 153, 154.

Cinchona officinalis, pests of, in Porto

Rico, **218.**

cinchonae, Protodiaspis. cincticollis, Antestia.

cinctus, Cephus.

cingulata, Herse; Rhagoletis.

cinnaberinus, Cucujus.

Cinnamomum camphora, Icerya purchasi on, in Persia, 16.

Cipollini (see Muscari comosum). circumflexa, Plusia (Phytometra).

Cirphis unipuncta, in Hawaii, 170, 274; in U.S.A., 3, 340; parasites and biological control of, 3, 170, 274; measures against, on wheat barley, 340.

cistipennis, Fundella (see F. pellucens). citri, Dialeurodes; Halticus; tetranychus; Pseudococcus; Scirto-

thrips; Unaspis (Prontaspis). citrina, Aonidiella.

citrinus, Aspidiotiphagus.

Citrullus vulgaris (see Watermelon).

Citrus, pests of, in Argentina, 171, 298; pests of, in Australia, 23, 72, 126, 127, 207, 208, 289, 364, 365, 383, 384, 385; fruit-piercing moths on, in Ceylon, 281, 282; Saissetia oleae on, in Chile, 206; Melanauster chinensis on, in China, 270; Coccids on, in Belgian Congo, 82; Lamellicorns on, in Costa Rica, 128; mite on, in Cyprus, 371; Coccids on, in Iraq, 16; Anastrepha on, in Mexico, 112, 376; Coccids on, in New Zealand, 21; fruit-piercing moths on, in Nigeria, 14, 15; pests of, in Nyasaland, 63; Ceratitis capitata on, in Palestine, 132, 133; Coccids on, in Persia, 16; Drosichiella tamarindus on, in Punjab, 285; Ceroplastes sinensis on, in Russia, 131; Coccids on, in Spain, 268, 269; pests of, in U.S.A., 24, 25, 79, 80, 112, 145, 176, 238, 262, 263, 374, 375; methyl bromide causing injury to, 374; (fruits), quarantine treatment against introduction of *Anastrepha* into U.S.A. in, 112; in baits for grasshoppers, 339.

Citrus Blackfly (see Aleurocanthus wog-

lumi).

Citrus Bud Mite (see Aceria sheldoni).
Citrus Red Mite (see Paratetranychus citri).

Citrus Red Scale (see Aonidiella

aurantii).

Citrus Thrips (see Scirtothrips citri).
Cladis nitidula, predacious on Asterolecanium spp. in Porto Rico, 219.

Cladosporium aphidis, infesting Aphids in Argentina and Uruguay, 299.

claripalpis, Theresia (Paratheresia). Clausenia purpurea, parasite of Pseudococcus comstocki in Virginia, 337.

Clavaspis herculeana, fungus infesting, on Aleurites in Nyasaland, 63.

Clay, as carrier for dusts, 235, 236, 303, 319; reducing effect of light on rotenone dusts, 50.

Clemora smithi, introduced Elaterid predacious on, on sugar-cane in Barbados, 159; measures against, 159.

Cleome monophylla, race of Myzus persicae on, in Nyasaland, 206.

Cleonus punctiventris, on beet in U.S.S.R., 131, 157; use of Metarrhizium anisopliae against, 131; crop rotations ineffective against, 157.

Clerodendron fragrans pleniflorum, Crocidolomia binotalis on, in Fiji, 101.

Clibadium surinamense, tests of, as an insecticide, 304.

Clinodiplosis mosellana (see Sitodiplosis). Clothes Moths, DDT against, **95.** (See Tinea pellionella, Tineola biselliella and Trichophaga tapetzella.)

Clothing (see Textiles).

Clover (Trifolium), 227; susceptibility of species of, to Halotydeus destructor in W. Australia, 162, 163; resistance of, to Odontria zealandica in New Zealand, 20; Apion on, in U.S.S.R., 157; pests of, in U.S.A., 57, 105, 374.

Clover, Red (see Trifolium pratense). Clover, Subterranean (see Trifolium

subterraneum).

Clover Leaf Weevil (see *Hypera punctata*).

Clysiana (Clysia) ambiguella (Vine Moth), new parasite of, in France, 32. coarctata, Hylemyia (Leptohylemyia).

Cob Nut (see Corylus avellana).

Coccinella septempunctata, parasite of, in Britain, 114; bionomics of, predacious on Aphids in India, 228.

coccinellae, Perilitus.

Coccobacillus acridiorum, imported into Algeria against Schistocerca gregaria, 10; bacteria similar to, in S. gregaria, 11.

Coccomytilus halli (see Nilotaspis).

Coccotrypes dactyliperda, on Phoenix canariensis in California, 375.

Coccus hesperidum, on Cinchona in Belgian Congo, 323; on Citrus in Iraq and Persia, 16; DDT against, on orchids in Maryland, 315.

Coccus pseudohesperidum, DDT against,

on orchids in Maryland, 315.

Coccus pseudomagnoliarum, measures against, on Citrus in U.S.A., 25.

Coccus viridis, on tea in Ceylon, 325; on coffee in Mysore, 187; honeydew of, as food for Neoplectrus maculatus, 325.

cockerelli, Dikraneura; Paratrioza.

Cockroaches, testing of attractiveness of fabrics and papers to, 127, 128; killed by DDT aerosol residue, 240; sites of action of DDT and nicotine in, 5; tests of sabadilla dusts on, 144; effect of inert dusts on transpiration through cuticle of, 60, 61.

Cocksfoot Grass (see Dactylis glomer-

ata).

cocois, Diaspis.

Coconut, Nephantis serinopa on, in Ceylon, 188; termites attacking, in Danger Is., 211; Coleoneura trichogramma on, in Fiji, 366.

Coconut Caterpillar (see Nephantis

serinopa).

Coconut Spike Moth (see Coleoneura trichogramma).

Coconut-oil Soap (see Soap).

Cocos plumosa, Coccid on, in California, 145.

Codling Moth (see Cydia pomonella).

Codophila varia, on lucerne in Tadzhikistan, 253.

Coelocaryon staneri, insects on, in Belgian Congo, 80.

Coeloides brevicaudis, sp. n., parasite of Pissodes spp. in Finland, 39, 40.

Coeloides stigmaticus, parasite of Pissodes notatus in Finland, 40.

coerulea, Zicrona.

coeruleocephala, Diloba.

Coffea (see Coffee).

coffeae, Aridelus (Helorimorpha); Saissetia; Zeuzera.

Coffee, Longicorns on, in tropical Africa, 96, 102, 361, 362; other pests of, in Belgian Congo, 80, 81, 82, 96, 153, 280, 281; pests of, in Mysore, 186, 187; Pseudococcus njalensis on, in Sierra Leone and Br. Togoland, 46; Antestia on, in Uganda, 277-279;

Anastrepha fraterculus not developing in berries of, 138.

Coffee Parchment, experiments with, in baits for locusts, 294, 295.

cognatus, Poeciloscytus.

Colaphellus bowringi, dust against, on cabbage in China, 344.

Cold Storage, against Cydia molesta in fruits, 374.

colemani, Phenacoccus.

Colemania sphenarioides, measures against, on graminaceous crops in Mysore, 207.

Coleoneura trichogramma, biological control of, on coconut in Fiji, 366; systematic position of, 83.

Coleophora laricella, on larch in Canada, 129, 130, 173; biological control of, 130.

Coleus, Orthezia insignis on, in S. Rhodesia, 33; Pseudococcus citri on, in U.S.A., 79.

Colias eurytheme, on lucerne in California, 111.

Colletotrichum, causing decay of fumigated pears, 190.

Collyria calcitrator, parasite of Cephus pygmaeus in Belgium, 185.

Colocasia, Megamelus proserpina on, in Fiji, 101; Organothrips bianchii on, in Hawaii, 276.

Colombia, new Coccid introduced into California on orchids from, 112.

Colorado, Cydia molesta in, 373; Dendroctonus engelmanni on Picea engelmanni in, 176; Vasates destructor on tomato in, 374.

Colorado Potato Beetle (see Leptinotarsa decembineata).

coloratus, Anthrenus.

columbiana, Phyllotreta.

Commelina, as food-plant of Helopeltis orophila in Belgian Congo, 154.

comonana, Bondia.

Comperiella bifasciata, introduced parasite of Aonidiella aurantii in W. Australia, 364.

completa, Rhagoletis suavis.

Compsilura concinnata, parasite of Lymantria dispar in U.S.A., 193. comptana, Ancylis.

comstocki, Pseudococcus.

conchiformis, auct., Mytilococcus (see Lepidosaphes ficus).

concinna, Chaetocnema. concinnata, Compsilura. confusum, Tribolium.

Congo, Belgian, pests of Cinchona in, 153, 323; coffee pests in, 80, 81, 82, 96, 153, 280; miscellaneous pests in, 81-83; beneficial insects in, 80, 81,

82, 96, 281; fungus infesting Zonocerus variegatus in, 280.

congoensis, Apanteles.

congolensis, Aspidoproctus. coniferarum, Diaspidiotus.

Connecticut, forest pests in, 193, 233, 236, 237; pests of maize in, 147, 236, 237; miscellaneous pests in, 105, 235, 237; orchard pests in, 49, 50, 219-221, 234, 235, 237, 308; Popillia japonica in, 194, 233, 234; pests of vegetables in, 105, 235, 236, 310; beneficial insects and biological control in, 193, 194, 234, 237.

Conoderus auritus, bionomics and control of, in Kentucky, 129.

Conoderus bellus, bionomics and control of, in Kentucky, 183.

Conophthorus edulis, on Pinus edulis in U.S.A., 57.

Conotrachelus, on Rheedia brasiliensis in Brazil, 6; on Pinus edulis in U.S.A., 57.

Conotrachelus anonae, sp. n., on Annona coriacea in Brazil, 6.

Conotrachelus bondari, on Annona squamosa in Brazil, 6.

Conotrachelus camelus, 6.

Conotrachelus copaiferae, sp. n., on Copaifera langsdorfii in Brazil, 6.

Conotrachelus licaniae, on Licania rigida in Brazil, 6.

Conotrachelus mamillatus, on Achras sapota in Brazil, 6.

Conotrachelus myrciariae, on Myrciaria jaboticaba in Brazil, 6.

Conotrachelus nenuphar (in U.S.A.), on apple, 51, 196, 220, 221; on peach, 140, 141, 201, 321, 335; on plum, 321; bionomics of, 140, 201, 321; measures against, 51, 141, 196, 201, 220, 221, 321, 325.

Conotrachelus phascoli, on beans in Brazil, 6.

Conotrachelus psidii, on guava in Brazil,

Conotrachelus ubacahy, sp. n., on Eugenia edulis in Argentina, 6.

consobrina, Phyllotreta.

conspersa, Narosa.

Contarinia juniperina, Inostemma associated with, on Juniperus bermudiana in Bermuda, 103, 190; sprays against, 191.

Contarinia sorghicola, on sorghum in Queensland, 387.

Contarinia tritici, bionomics of, on wheat in Sweden, 36.

contorticornis, Hypocampsis.

Convallaria majalis, Sciara on, in Denmark, 160. convergens, Hippodamia.

arvensis, Spermophagus Convolvulus sericeus on, in Uzbekistan, 155.

Convolvulus sepium, infusions of, against Tetranychus turkestani, 49.

Copaifera demeusei, Aspidoproctus congolensis on, in Belgian Congo, 82.

Copaifera langsdorfii, new weevil on, in Brazil, 6.

copaiferae, Conotrachelus.

Copidosoma truncatella (see Litomastix). Copper, treatment of wood with compounds of, against termites, 172, 173; DDT compatible with insoluble compounds of, 239.

Copper Arsenate, sap-stream treatment of trees with, as timber preservative,

Copper Arsenate (Basic), against cotton pests, 179, 356; against pests of leguminous crops, 178, 355; against pests of solanaceous crops, 29, 84, 177, 178, 224, 354, 355, 356, 358; increasing infestation of potato by Aceratagallia uhleri, 224; dusting with, 177, 179, 224, 354, 355, 356, 358; and kaolin, 354; and sulphur, 179, 224, 358; spraying with, 29, 178, 179, 356; adhesive for, 356; formulae containing, 179, 224, 354, 356, 358; effect on plants of soil containing, 355.

Arsenate (Neutral), against Copper Leptinotarsa decemlineata on potato,

84.

Copper Arsenite, against Leptinotarsa

decemlineata on potato, 84.

Copper Chloride, against weeds, 226; and arsenic acid, sap-stream treatment of trees with solution of, as timber preservative, 56.

Copper Fluoride, treatment of wood

with, against termites, 172.

Copper Nitrate, against weeds, 226. Copper Oxychloride (in sprays), repellent to Leptinotarsa decembineata, 85: slightly reducing efficiency of dinitro

compound against mites, 100. Copper Pentachlorphenate, treatment of wood with, against termites, 172.

Copper Phosphates, treatment of wood

with, against termites, 173.

Copper Sulphate, impregnation of trees and timber with, against insects, etc., 56, 172; uses of, to destroy plants, 226. (See Bordeaux Mixture.)

Coptotermes, in timber in E. Africa, 103; use of arsenic against, in tea bushes in Ceylon, 211; influence of reproductive forms on control of, in timber in New Zealand, 169.

Coptotermes acinaciformis, New Zealand, 169.

Coptotermes frenchi, in New Zealand, 169.

Coptotermes lacteus, in New Zealand,

corchori, Apion.

Corchorus spp., pests of, in India, 117, (See Jute.) 163, 164.

Corcyra cephalonica, measures against, in stored cereals in India, 325; Bracon greeni reared on, 165.

Cordia myxa, new Coccid on, in Iraq, 16. Corioxenos antestiae, bionomics and utilisation of, against Antestia in Uganda, 278, 279.

Corizidolon, parasite of, on cotton in

Uganda, 296.

Corizus hyoscyami, on cotton in Bokhara, 154.

Corn Borer, European (see Pyrausta nubilalis).

Corn Earworm (see Heliothis armigera). corni, Eulecanium (Lecanium).

cornicularius, Pemphigus. cornutus, Bostrychoplites.

Sublimate (see Corrosive Mercuric Chloride).

corvina, Musca (see M. autumnalis).

Corylus avellana (Filbert), Croesus septentrionalis on, in Britain, 273; pests of, in U.S.A., 111, 374.

Corymbites, soil factors affecting injury by, in Germany, 92; survival of, in crop rotations in U.S.S.R., 160.

Corymbites aeneus, 92.

Corymbites pectinicornis, 92.

subtilis, pyrethrum Cosmia against, on apricot in Central Asia,

Cosmophila flava, bionomics and control of, on cotton in China, 381.

Cosmophila sabulifera (see Anomis).

Cosmopolites sordidus, on Musa spp. in Cuba, **172.**

Cosmopteryx attenuatella ab. luteoapicalis, n., on sorghum in Senegal, 96. Cossids, immature stages of Indian, 288.

Cossus cossus, trees infested by, in Finland, 41; damaging lead of telephone cables, 272.

Costa Rica, Lamellicorns on Citrus in,

costalis, Myzocallis (Monellia).

costimacula, Eublemma.

Cotoneaster, pests of species of, in Britain, 184; Lepidoptera on, in continental Europe, 185; pests of, in U.S.A., 109, 185, 373; strains of Eriosoma lanigerum on apple and, 184. cotoneastri, Cremona; Depressaria.

Cotton, Platyedra gossypiella on, in Fr. W. Africa, 96; pests of, in S. Africa, 166; Alabama argillacea on, in Argentina, 300; A. argillacea on, in Barbados, 159; Heliothis virescens on, in Brazil, 334; pests of, in China,

419 INDEX.

120, 293, 381; pests of, in Belgian Congo, 81, 280; Gryllotalpa on, in Egypt, 20; pests of, in India, 228, 283, 284; pests of, in Morocco, 66, 257; pests and disease of, in Nigeria, 96, 169; pests of, in Nyasaland, 62, 168, 332; new Coccid on, in Persia, 16; new Capsid on, in Peru, 128; pests of, in Queensland, 165, 166; pests of, in S. Rhodesia, 166; pests of, in Russian Central Asia, 49, 154: pests of, in Sudan, 166; pests of, in Tanganyika, 81, 82, 167; Capsids on, in Uganda, 13, 296; pests of, in U.S.A., 178, 180, 247, 303, 311, 315, 345, 355, 356, 374; varieties and time of planting of, in relation to pests, 13, 14, 120, 166, 167, 169; variety of resistant to leaf-curl virus, 169; effects on, of insecticides in soil, 355; dusts and sprays to cause defoliation of, **178, 315.**

Cotton Aphis (see Aphis gossypii).

Cotton Boll Weevil (see Anthonomus grandis).

Cotton Bollworm, American (see Heliothis armigera).

Cotton Bollworm, Green (see Heliothis

vivescens). Cotton Bollworm, Pink (see Platyedra

gossypiella). Cotton Bollworm, Spotted (see Earias

fabia and E. insulana). Cotton Bollworm, Sudan (see Diparopsis

castanea). Cotton Fleahopper (see Psallus seriatus). Cotton Leaf-roller (see Sylepta derogata). Cotton Leafworm (see Alabama argil-

lacea). Cottonseed, fumigation of, against Platyedra gossypiella in, in Morocco, (husks), in baits for grass-

hoppers, 339. Coumarin, in bait for Sitona cylindri-

collis, 30.

Cowpeas (Vigna unguiculata), Chalcodermus aeneus on, in S. Carolina, 355; Bruchus obtectus in, in Chile, 205; Lamprosema indicata on, in Belgian Congo, 81; Fundella pellucens on, in Porto Rico, 104; new weevil on, in Senegal, 206: effects of insecticides on, 355.

crabro, Vespa.

Crambus, crop rotation against, U.S.S.R., 157.

Crambus hortuellus, on cranberry in Massachusetts, 51.

Cranberry, pests of, in U.S.A., 51, 75, 313: insecticides causing injury to, 51.

Cranberry Fireworm (see Rhopobota naevana).

Cranberry Fruitworm (see vaccinii).

crataegella, Scythropia. crataegi, Glossonotus.

Crataegus (Hawthorn), 90; Caliroa limacina on, in W. Australia, 365; insects on, in Britain, 185, 272; Coccid on, in Iraq, 16.

cremastoides, Nemeritis (Phaedroctonus)

(see N. moderator).

Cremastus shirakii, parasite of Schoenobius bipunctifer in China, 152.

Crematogaster, associated with Pseudococcus njalensis in W. Africa, 46.

Cremona cotoneastri, on Cotoneaster in Oregon, 185.

crenulata, Lachnosterna.

crenulella, Apterona.

Creosote, in dust against Hylemyia brassicae. 319; uses of, treatment of timber, logs and tree wounds against pests, 12, 27, 54, 55, 259; and oil, as repellent for Chlorops, 286; effect of, as a bee-repellent in orchard sprays, 55.

Cresols, treatment of wood with, against termites, 172; accelerating penetration of insecticide through insect

cuticle, 87.

Crested Dogstail Grass (see Cynosurus cristalus).

crinitus, Sitona.

Crioceris asparagi, dusts against, on asparagus in Washington, 72.

cristatus, Arilus. croceus, Nanophyes.

Crocidolomia binotalis (on crucifers), in Fiji, 101, 366; in Queensland, 231; bionomics of, 101; measures against, 101, 231.

crocodora, Dichocrocis.

Croesus septentrionalis, bionomics and control of, on Corylus avellana in Britain, 273.

Crotalaria, recommended use of, to protect coffee from Dasus simplex,

Crotalaria agatifolia, possible food-plant of Helopeltis orophila in Belgian Congo, 153.

Crotalaria anagyroides, Xyleborus morstatti on, in Mysore, 187.

Crotalaria juncea, Etiella zinckenella on, in Belgian Congo, 81.

cruciata, Austroicetes

cruciferae, Phyllotreta.

Cryolite, against caterpillars on cabbage, 101, 231, 310; against cotton pests, 179, 356; against pests of cucurbits, 52, 149; against pests of fruit trees, 73, 74, 175, 208, 321, 322, 383; against pests of leguminous plants, 138, 305, 310, 316, 317, 318; against

pests of solanaceous plants, 29, 72, 84, 178, 224, 236, 354, 356, 358, 359, 382 against pests of other plants, 53, 105, 135, 176, 177, 200; in baits, 179, 191, 200; dusting with, 29, 52, 105, 135, 138, 149, 177, 178, 179, 208, 224, 231, 236, 305, 310, 316, 317, 318, 354, 356, 358, 359, 382; carriers for, 105, 138, 149, 208, 236, 316, 317, 354, 382; factors affecting coverage by dusts of, 310; spraying with, 29, 53, 72, 73, 84, 101, 105, 138, 175, 200, 208, 316, 318, 322, 356, 383; adhesives and spreaders for, 72, 175; formulae containing, 29, 50, 52, 72, 84, 101, 105, 135, 138, 149, 175, 179, 191, 200, 208, 224, 231, 316, 317, 318, 321, 354, 358, 359, 382, 383; application of, from aircraft, 176, 177; and barium fluosilicate (see Dutox); and copper compounds, 52, 149, 208; DDT compatible with, 157; and dinitro compounds, 73, 74; and ferric dimethyl dithiocarbamate, 53; and lime-sulphur, 29; and nicotine, 52; and oil emulsions, 74, 175, 200, 208, 321; "inverted" spray mixture of oils and, 50; and rotenone, 29, 356; and sulphur, 29, 138, 179, 224, 317, 318, 358, 359; and thiocyanate, 316, 317; and zinc sulphate, 50; and injury to plants, 50, 321, 322; aluminium oxide and calcium phosphate useless as safeners for, 322.

Cryphalus (Trypophloeus) alni, bionomics of, on Alnus incana in Finland,

9.

Cryphalus bispinulus, parasite of, on aspen in Finland, 39, 41.

Cryptaspidiotus shastae, spray against, on Juniperus in Kansas, 200.

Crypticerya natalensis, parasite of, on rose in Belgian Congo, 82.

Crypticerya rosae, parasite of, or Grevillea in New South Wales, 82.

Cryptoblabes gnidiella, new parasite of, in Belgian Congo, 81; predacious on Stictococcus, 81.

Cryptochetum, catalogue of, 83.

Cryptochetum (Tritolestes) aspidoprocti, subgen. et sp. n., parasite of Coccid in Belgian Congo, 82.

Cryptochetum curtipenne, parasite of Coccids in Belgian Congo and Ceylon, 82.

Cryptochetum iceryae, bionomics of, in Belgian Congo, 82.

Cryptochetum (Tritolestes) melanum, subgen. et sp. n., parasite of Icerya spp. in Belgian Congo, 82.

eryptographus, Xyleborus.

Cryptohypnus (see Hypnus).

Cryptorrhynchidius (see Cryptorrhyn-chus).

Cryptorrhynchus lapathi, on Salix and Alnus in Finland, 41.

Cryptostemma calendulaceum, Listroderes obliquus on, in W. Australia, 23.

Cryptotermes (see Kalotermes).

Cryptus (parasite of Loxostege sticticalis), factor affecting diapause of, 295.

Cryptus sexannulatus, introduced into U.S.A. from Canada against Cydia pomonella, 179.

Ctenolepisma spp., testing of attractiveness of fabrics and paper to, in U.S.A.,

Cuba, weevils on Musa spp. in, 172.

cubaecola, Sturmia.

Cubé, dusting with, 27, 50, 84, 112, 137, 178, 258; oil-impregnated dust of, 50; carriers for, 50, 112, 258; spraying with, 72, 80, 84, 112, 137, 313; formulae containing, 72, 80, 112, 137, 258, 313; and barium fluosilicate, 84; and nicotine, 137; and oil emulsions, 80; rotenone contents of, 28, 50, 72, 80, 112, 137, 258, 304; other insecticides compared with, 72, 80, 84, 137, 258, 304, 313. (See Lonchocarpus.)

Cucujids, survey of data on, in stored

products, 328.

Cucujus cinnaberinus, in galleries of Xyleborus cryptographus in Finland,

Cucumber, experiments with Dacus cucurbitae and, in Hawaii, 276, 308; pests and disease of, in U.S.A., 148, 355; insecticides and injury to, 149, 239, 240; methyl bromide injuring packed fruit of, 109.

cucumeris, Harr., Epitrix.

cucumeris, auct., Epitrix (see E. tuberis). cucurbitae, Baris; Dacus; Phloeophagosoma.

Cultural Practices, influence of, on insect pests, 74, 147, 156, 183, 216, 232, 235.

cupreoviridis, Earias.

Cupressus, Diaspidiotus coniferarum on, in California, 109.

Cupric Compounds (see under Copper). Cuprous Oxide, effect of, on dinitro compound against mites, 100.

curculionis, Bathyplectes. Currant, pests of, in Oregon, 79.

Currant, Black, pests of, in Britain, 35, 272.

curtipenne, Cryptochetum.

curvipes, Thalessa (Pimpla).

Custard Apple (see Annona squamosa). Cutworm, Pale Western (see Agrotis orthogonia).

cyanea, Scutellista.

cyaneus, Sirex.

Cyanogas, 290, 323.

Cybocephalus, establishment of, against Coccids in California, 145.

Cyclohexane, 87.

Cyclohexanone, as a solvent for DDT, 239, 240, 242, 244, 338, 346; plants injured by, in aerosols, 239, 240, 244.

Cyclohexylamine, in repellent spray against Anastrepha ludens, 376.

Cyclohexylamine Dodecyl Sulphate, as an emulsifier, 44.

Cyclopentane Propionic Acid, treatment of wood with compounds of, against termites, 172.

Cydia, measures against, on peas in Holland, 90.

Cydia illutana, in spruce cones in Finland, 64.

Cydia molesta (Oriental Peach Moth), fungus infesting, in Argentina, 299; in Australia, 138; in Mexico, 110; in U.S.A., 108, 109, 110, 139, 141, 142, 204, 234, 246, 249, 305, 321, 322, 354, 373, 374; on quince, 246, 249; bionomics and miscellaneous foodplants of, 109, 110, 322, 373; relation of, to Sclerotinia fructicola, 321; parasites and biological control of, 110, 138, 139, 141, 142, 204, 234, 305, 322, 354, 373; baits for, 322; treatment of harvested fruits, etc., against, 109, 110, 374; other measures and experiments against, 246, 249, 321, 373.

Cydia nigricana (Pea Moth), in Br. Columbia, 73; in Washington, 73, 76; parasites and biological control of, 73; eradication of vetches against, 76.

Cydia packardi, sprays against, or cherry in Washington, 75.

Cydia pomonella (Codling Moth), in S. Africa, 189; in Australia, 138; in Canada, 175, 179; in Denmark, 8; in France, 349; in New Zealand, 323; in Spain 200; in New Zealand, 323; in Spain, 288; in U.S.A., 4, 50, 51, 73, 74, 75, 106, 107, 109, 139, 175, 179, 195, 196, 197, 198, 202, 203, 221, 248, 308, 309; on pear, 349; on walnut, 309; relation of, to Pseudomonas juglandis, 309; studies on adult activity and oviposition of, 4, 8, 198; other bionomics of, 8, 202, 309; parasites and biological control of, 138, 139, 179, 288, 323; solid baits for, 308; dusts against, 195, 197, 248; sprays against, 4, 8, 50, 51, 52, 73, 74, 75, 106, 107, 175, 195, 196, 197, 198, 199, 202, 203, 221, 248, 309, 349; banding and orchard sanitation against, 73; treatment against hibernating larvae of, 175; fumigation of apples against, 109, 189; toxicity of extract of Erigeron affinis to, 359; strains of, resistant to insecticides, 107, 195, 196.

Cydia strobilella (in spruce cones), in Finland, 39, 40, 64; in Sweden, 40; parasites of, 39, 40.

Cydonia japonica, Cydia molesta on, in California, 109.

cylindricollis, Sitona.

Cyllene robiniae, factors affecting infestation of Robinia pseudacacia by, in N. Dakota, 233.

Cymbidium (see Orchids).

Cynodon plectostachyum, Laphygma exempta on, in Nyasaland, 63.

Cynosurus cristatus, resistance of, to Odontria zealandica in New Zealand, 20.

Cypress (see Cupressus).

Cypripedium (see Orchids).

Cyprus, Phyllocoptruta oleivorus on Citrus in, 371.

Cyrtacanthacris, attacking Cinchona in Belgian Congo, 323.

Cyrtopeltis tenuis, on tomato in Queensland, 288.

Cyrtopeltis varians (see Engytatus geniculatus).

Cyrtopodium (see Orchids).

Cyrtorhinus fulvus, predacious on Megamelus proserpina in Fiji, 101.

Cyrtorhinus riveti, in Fiji, 101.

Cytherea spp., parasites of Dociostaurus maroccanus in Spain, 265.

Czechoslovakia, miscellaneous insects in, 212, 328.

D,

D-4, **52, 55.**

Dacnusa gracilis, parasite of Psila rosae in Britain, 45; killed by DDT, 45.

Dacrydium cupressinum, resistance of timber of, to Anobium punctatum in New Zealand, 211.

dactyliperda, Coccotrypes.

Dactylis glomerata, experiment with Hylemyia coarctata and, in Britain, 368; Glyphipteryx achyloessa on, in New Zealand, 323; resistance of, to Odontria zealandica, 20.

Dactylopius, utilisation of, against

Opuntia in Mauritius, 165.

Dacus cucurbitae (in Hawaii), host-fruits and control of, 276; effect of foods on ovarian development in, 308.

Dacus ferrugineus tryoni (in New South Wales), in loquats, 71; trapped by bean leaves, 71; tests of bait-sprays against, 385.

Dacus oleae (on olive), parasites of, in Greece, 326; in Spain, 267,

422 INDEX.

268; bionomics of, 267; encouragement of Cecidomyiid transmitting *Macrophoma dalmatica* against, 326; baits for, 268.

Dacus passiflorae, on cacao in Fiji, 101; parasite of, 101.

Dadobia immersa, in galleries of Saperda carcharias in Finland, 41.

daghestanica, Podonta.

Dahlbominus, n.n., for Microplectron, Dahlb., 252.

Dahlbominus (Microplectron) fuscipennis, parasite of Neodiprion sertifer in Finland, 39; utilisation of, against Gilpinia hercyniae in Quebec and Connecticut, 173, 237; parasitising G. frutetorum, 237; strains of, 173; type of genus, 252.

Dahlia, Eulia loxonephes on, in Argentina, 298; (host of tobacco virus), modification of restrictions on importation of, into Nyasaland, 62.

Daikon Radish (see Raphanus sativus longipinnatus).

Dakota, North, Sitona cylinaricollis on Melilotus in, 29, 30; Cephus cinctus on wheat and flax in, 146.

Dakota, South, grasshoppers in, 231-233; mite destroying grasshopper eggs in, 232.

Dalbergia sissoo, insects on, in India, 285.

dampfi, Frankliniella.

Dana Derris 10 & 33, 7.

Dandelion (see *Taraxacum officinale*). Danger Islands, *Kalotermes* spp. on coconut in, 211.

Daphne, relation of Myzus persicae to, 367.

darwiniensis, Mastotermes.

Dasus simplex, measures against, on coffee in Belgian Congo, 153.

Dasyneura affinis, bionomics and control of, on Viola odorata in Egypt, 18. Dasyneura ignorata, bionomics and control of, on lucerne in U.S.S.R., 131, 132.

Dasyneura mali, on apple in Connecticut, 237.

Dasyneura rhodophaga (see Neocerata).
Dasyneura strobi, on spruce in Finland,
38, 39, 64; parasites of, 38, 39.

38, 39, 64; parasites of, 38, 39. Date Palm, Coccids on, in Iraq and Persia, 16; Ceratitis capitata on, in Palestine, 133.

Date Palm Ornamental (see *Phoenix* canariensis).

Dates (see Fruit, Dried).

Datura, nectar of, increasing oviposition by Protoparce sexta, 335.

Datura meteloides, Agalliana ensigera on, in Argentina, 380.

Datura stramonium, Agalliana ensigera on, in Argentina, 380; A. ensigera not transmitting curly-top to, 380; experiment with Leptinotarsa decemlineata and, 264.

D-D Mixture (1,2-Dichlorpropane and 1,3-Dichlorpropylene), against *Kalotermes piceatus*, 275; soil treatments with, against pests, 76, 135, 178, 374.

DDT [2,2-bis (parachlorphenyl)-1,1,1trichlorethane], against Aleurodids, 171, 240, 275, 334; against Aphids, 45, 178, 224, 231, 239, 240, 241, 242, 243, 248, 261, 309, 337, 346, 383, 384, 385, 386, 387; favouring infestation by Aphids, 75, 80, 107, 309; against Capsids, 45, 180, 224, 243, 247, 260, 261, 276, 345, 384; against Coccids, 75, 79, 80, 106, 240, 314, 315, 334, 336, 384; against Jassids, 43, 223, 224, 239, 243, 244, 247, 248, 262, 263, 337, 384; against Lygaeids, 250, 301, 302, 382, 384; against Paratrioza cockerelli, 223, 224; against Pentatomids, 244, 355, 383, 386; against other Hemiptera, 72, 243, 244, 246; against ants, 238, 249; against Coleoptera, 7, 29, 34, 57, 58, 72, 93, 94, 176, 208, 223, 224, 238, 239, 243, 244, 245, 246, 256, 275, 327, 333, 334, 337, 339, 346, 355, 359, 382, 383, 385, 386, 387; uses of, as killing agent in traps for Popillia japonica, 245; crickets and cockroaches killed by, against Diptera, 43-45, 143, 221, 235, 240, 250, 276, 384, 385, 386; protection of house-flies pollinating onions from, 240; Forficula auricularia killed by, 249, 254, 255, 276, 305, 309, 318, 319, 337, 339, 354, 355, 357, 373, 382, 383, 384, 385, 386, 387; against mites, 240, 384, 385; favouring infestation by mites, 75, 80, 106, 107, 196, 197, 384, 387; against sawflies, 7; against Scutigerella immaculata, 318; treatment of wood with, against termites, 172, 345; against thrips, 143, 239, 240, 243, 261, 262, 263, 338, 383, 384, 385, 387; protecting fruits from Vespa spp., 229; woodlice killed by, 240, 255; effect of, on insect parasites and predators, 45, 107, 146, 224, 238, 244, 261, 309, 336, 337, 387; risk of insect outbreaks resulting from widespread use of, 146; effects of, on honey-bees, 310; mode of action of, on insects, 5, 44, 58, 106, 255; techniques of testing, on insects, 254, 255, 320; sources of error in field-plot tests of, 43; duration of effectiveness of, on

423

plants, 44, 75, 106, 176, 224, 238, 240, 245, 246, 248, 249, 261, 275, 385, 386; duration of toxicity of, on other surfaces, 44; question of fungicidal action of, 224; and injury to plants, 107, 176, 195, 248, 249, 338; slightly retarding germination of seeds, 250. 318; problems of residues of, on fruits, etc., 106, 107, 160, 249, 338; question of risks and toxicity of, to mammals and birds, 96, 107, 230, 320, 327, 334, 337, 348, 386; in aerosols, 96, 239-244, 320, 327, 338, 348; distribution of, from fog generator, 262; in bait-sprays, 276, 383, 385; in bands and barriers, 238, 250; in 231, 235, 238, 239, 244, 245, 247, 248, 249, 250, 256, 260, 261, 275, 276, 301, 305, 310, 318, 319, 327, 337, 338, 345, 348, 354, 355, 356, 357, 359, 382, 383, 384, 385, 386; carriers for, 143, 157, 180, 197, 224, 238, 239, 244, 245, 247, 249, 250, 338, 345, 382, 386; carrier impregnated with, 319; in sprays, 29, 34, 43, 44, 45, 75, 79, 80, 93, 94, 96, 105, 106, 107, 137, 143, 171, 185, 195, 196, 197, 203, 216, 224, 229, 235, 238, 239, 242, 244, 245, 246, 247, 248, 249, 275, 276, 302, 309, 314, 315, 327, 334, 336, 338, 339, 346, 348, 356, 382, 383, 384, 385, 386, 387; application of, from aircraft, 176, 247; treatment of bean seed with, 318; soil treatment with, 43, 44, 143, 176, 203, 318, 382; treatment of papers, clothes, etc., with, 58, 95, 327; uses of, against pests of stored products, 177, 333, 334, 339, 355, 383, 385; with acetone, 239, 240, 242, 319, 338, 339; with alcohol, 58; with arsenicals, 106, 157, 197, 247, 345; with benzene, 345; with Bordeaux mixture, 106, 157, 239, 245, 247, 249, 255, 384; other copper compounds, 238, 239; with cyclohexanone, 239, 240, 242, 338, 346; with 2,3-dichlor-1,4-naphthoquinone, 157; with disodium ethylene bisdithiocarbamate, 239: with with dinitro compound, 106; ethylene dichloride, 106, 245; with ferric dimethyl dithiocarbamate, 106, 157, 239, 248; with fluorine compounds, 157; with lime, 106, 239, 255; with lime-sulphur, 106, 157, 245, 255; decomposition of, by nicotine, 157; with oils, 58, 75, 79, 80, 106, 143, 185, 195, 238, 239, 240, 242, 247, 248, 255, 256, 262, 275, 302, 310, 338, 339, 346, 383, 384, 385; with orthodichlorbenzene, 242; with pyrethrum, 157, 256; with pyrophyllite in sprays,

224, 238, 244, 245; with rotenone insecticides, 80, 157, 239, 243, 244; with solvent naphtha, 45, 382; with sulphur, 106, 143, 157, 245, 345, 359; with butyl thiocyanate, 339; with trichlorethylene, 242; with Velsicol AR60, 319; with xylene, 44, 244, 245, 314; wetting and spreading agents for, 34, 106, 242, 244, 246, 385: emulsifiers for solutions of, 44, 106, 238, 244, 245, 314; reactions causing incompatibility of, with other chemicals, 157, 255; not affected by ultra-violet rays, 235; methods of determination and analysis of, 5, 160, 320, 356, 359, 360, 388; composition of technical, 255, 359, 388; books and surveys of data on, 43, 160, 327, 348; proprietary preparations of, 7, 34, 43, 80, 93, 94, 105, 106, 143, 171, 247, 248, 327, 333, 334; list of proprietary insecticides containing, 288.

DDT Analogues, tests of toxicity of, to insects, 255, 360; reactions of, to colorimetric test for DDT, 5; in

technical DDT, 255, 360.

DDX (see Parachlorphenyldichlorethylene).

Deccan Grasshopper (see Colemania sphenarioides).

decemlineata, Leptinotarsa. decipiens, Empoasca.

decora, Stenoma. decorus, Oedaleus.

Decticus albifrons, in Spain, 268.

degenerata, Hemiberlesia.

Deilephila nerii, on Cinchona in Belgian Congo, 323.

Delaware, Popillia japonica in, 107.

delicatus, Macrocentrus.

Dendrobium (see Orchids).

Dendroctonus engelmanni, outbreak of, on Picea engelmanni in Colorado, 176. Dendroctonus frontalis, on Pinus spp. in

N. Carolina, 158.

Dendroctonus monticolae, outbreaks and control of, on Pinus contorta latifolia

in Canada, 360.

Dendroctonus piceaperda, on conifers in Canada, 129, 173; measures against,

Dendrolimus pini, bionomics of, on pine in Finland and Germany, 42, 99.

dendrophaga, Pyroderces.

in logs and stumps, 173.

Dendrothripoides ipomeae, on sweet potato in Barbados, 159; food-plants of, in Hawaii, 274.

Denmark, Cerura bifida damaging lead cable in, 271; Hylotrupes bajulus in timber in, 12, 116; miscellaneous pests in, 7, 8, 116; new Nematode infesting Sciara on Convallaria in, 160. denticornis, Hyalochloria.

dentipes, Monodontomerus.

Deobase, 341.

Depressaria cotoneastri, on Cotoneaster in Europe, 185.

depressella, Emmalocera.

Devaeocoris, parasites of, on cotton in Uganda, 296.

Devaeocoris punctulatus, on cotton in Bokhara, **154.**

derasofasciatus, Agrilus.

Dermatea molliuscula, 130.

Dermestes, natural sources of, infesting animal products, 150.

Dermestes ater (cadaverinus), infesting rabbit skins imported into U.S.A. from Hawaii, 274.

Dermestes frischi, methyl bromide against, in dried fish and meat in Morocco, 66.

Dermestes maculatus (vulpinus), festing rabbit skins imported into U.S.A. from Hawaii, 274; inert dusts against, 35.

Dermolepida albohirtum, parasite and control of, on sugar-cane in Queens-

land, 1.

derogata, Sylepta. Derris, against ants, 27, 28, 238; against Aphids, 143, 241, 242, 385; stimulating reproduction of Aphids, 143, 144; against Coccids, 80; against other Hemiptera, 89, 126, 243; against Coleoptera, 7, 23, 84, 138, 149, 178, 235, 258, 273, 274, 310, 365; against Rhagoletis pomonella, 50; against Lepidoptera, 7, 53, 68, 91, 126, 136, 138, 169, 231, 243, 310, 317, 337, 382, 385; against sawflies, 7, 37, 59, 88, 89, 229, 273; against thrips, 53, 67; tests of toxicity of, to insects, 59, 88, 258; resistance of Bombyx mori to, 91; list of insects susceptible to, in Holland, 97; in aerosols, 241, 242, 243, 244; dusting with, 7, 23, 27, 28, 50, 59, 67, 68, 84, 88, 89, 91, 126, 136, 138, 149, 169, 231, 235, 258, 274, 310, 317, 337, 365, 382, 385; carriers for, 23, 50, 89, 136, 149, 235, 258, 274, 317, 382; oil-impregnated dust of, 50; proprietary dust of, 7: factors affecting coverage by dusts of, 310; spraying with, 37, 50, 53, 59, 67, 80, 84, 126, 136, 138, 242, 244, 273, 274, 365; formulae containing, 23, 80, 242, 243, 258, 382; reducing toxicity of DDT aerosol, 243, 244; intermediary solvents for extractives of, when used with oils, 80; and pyrethrum, 89, 317; and soap, 274; and sulphur, 138; and thiocyanate, 317; and Velsicol, 143; rotenone contents of, 7, 28, 50, 59, 67, 68, 88, 89, 91, 126, 136, 138, 143, 169, 217, 229, 231, 235, 242, 258, 317, 337; effect of storage on dusts and roots of, 89, 217, 218.

Derris spp., cultivation and insecticidal constituents of, 217, 281, 288.

Desert Locust (see Schistocerca gregaria). Desmodium, insects on, in U.S.A., 105,

destructor, Ceroplastes; Halotydeus; Mayetiola (Phytophaga); (Phyllocoptes).

Dewberry, Ancylis comptana on, in New

Jersey, 142.

Diabrotica, on cucurbits in Indiana, 148. Diabrotica melanocephala, bionomics and control of, on cucurbits in Indiana, 148; transmitting Bacillus tracheiphilus, 148.

Diabrotica undecimpunctata (soror), on lucerne in California, 111; cepted in aircraft in Hawaii, 2.

Diabrotica vittata (see D. melanocephala). Diacrisia obliqua, measures against, on jute in Bengal, 164.

diadema, Sinea.

diademata, Phyllotreta.

Diaeretus rapae, parasite of Brevicoryne brassicae in Washington, 73.

Dialeurodes citri, eradication of, in California, 109, 373.

Diamond-back Moth (see Plutella maculipennis).

dianthi, Aphis (see Myzus persicae).

Dianthus spp., Aphids on, in Holland, 367. (See Carnation.)

Diaphania nitidalis, DDT against, on cucurbits in S. Carolina, 355.

Diaprepes abbreviatus, measures against, on sugar-cane in Barbados, 159.

Diarthrothrips saccharicolus, sp. n., on sugar-cane in New Caledonia, 275.

Diaspidiotus ancylus, food-plants of, in California, 109, 374.

Diaspidiotus coniferarum, on Cupressus in California, 109.

Diaspidiotus liquidambaris, bionomics of, on Liquidambar in California, 109. diaspidis, Aphytis (Aphelinus).

Diaspis boisduvali (in U.S.A.), introduced Nitidulid predacious on, on Opuntia,

145; DDT against, on orchids, 315. Diaspis cocois, introduced Nitidulid predacious on, on Cocos plumosa in

California, 145. Diaspis pentagona (see Pseudaulacaspis). Diatraea grandiosella, on maize and sorghums in U.S.A., 180, 192.

Diatraea saccharalis (Sugar-cane Moth Borer), dusts against, in Louisiana, 177; in Mexico, 179; in W. Indies, 31, 159, 179, 219, 320; on maize, 159; parasites and biological control of, 31, 159, 179, 219, 320.

INDEX.

Diatraea venosata (see Proceras).

diatraeae, Lixophaga.

Dibrachys cavus (boucheanus), parasite of Neodiprion sertifer in Finland, 39; capsules for studying oviposition and development of, 315.

Dichlordifluoromethane, use of, for producing aerosols, 239, 241, 242, 243,

338.

Dichlorethane (see Ethylene Dichlo-

ride).

Dichlorethyl Ether, and oils, treatment of maize ears with, against *Heliothis armigera*, 147; soil treatments with solutions and emulsions of, against pests, 78, 201, 336; causing injury to strawberry, 201.

Dichlorethylene, 98.

2,3-Dichlor-1,4-naphthoquinone, DDT compatible with, 157.

1,1-Dichlor-1-nitroethane, fumigation tests with, 342, 343.

1,2-Dichlorpropane (Propylene Dichloride), emulsion of, against *Aegeria exitiosa*, 336; and dichlorpropylene (see D-D Mixture).

1,3-Dichlorpropylene (Dichlorpropene), and dichlorpropane (see D-D Mixture). *Dichocrocis crocodora*, new parasite of,

on coffee in Belgian Congo, 80.

Dichocrocis punctiferalis, bionomics and control of, on sorghum in Queensland, 387.

Dicranura vinula, damaging lead of telephone cables, 272.

dictyospermi, Chrysomphalus.

Dicyclohexylamine, effect of spray containing, on Anastrepha ludens, 376.

Dicyclohexylamine Di - dinitrocyclohexylphenolate (see under Dinitro-ocyclohexylphenol).

Dicyphus, DDT against, on rhubarb in

New South Wales, 384.

Dicyphus minimus, Engytatus geniculatus predacious on, on tobacco in N. Carolina, 347.

Didyma setifacies (see Lypha).

didymator, Anilastus.

Diethylene Glycol Monoethyl Ether, not accelerating penetration of diphenylamine through insect cuticle, 87.

differentialis, Melanoplus.

dignus, Telenomus (Phanurus).

Dikraneura cockerelli, DDT fog against, on vines in Arizona, 262.

Diloba coeruleocephala, bionomics and control of, on almond in Spain, 266.

Dimethyl Ether, for producing aerosols, 241.

dimidiatus, Carpophilus; Tyrophagus (Tyroglyphus).

Dinitro-o-cresol, in dust barrier against Blissus leucopterus, 250; dusting with, against Eurydema oleraceum, 89; in sprays against orchard pests, 7, 73, 74, 90, 93, 94, 95, 175; ovicidal effect of, 7, 73, 96; causing injury to plants, 74, 89, 100; in sprays against weeds, 226; and lead arsenate or cryolite, 73, 74; and oil emulsions, 74, 93; salts of, 73, 74, 90, 100, 226; formulae containing, 7, 73, 226, 227, 250; proprietary preparations containing, 90, 93.

Dinitro - o - cyclohexylphenol, against Coleoptera, 52, 141, 316; ineffective against Cydia pomonella, 74, 106, 203; against Heliothis armigera, 179, 316; against Jassids, 52, 313; against other insects, 141, 178, 238; toxicity of, to honey bees, 55; against mites, 52, 54, 73, 100, 233; against weeds, 141; in dusts, 52, 55, 100, 178, 179, 313, 316; in sprays, 52, 54, 73, 74, 100, 106, 141, 203, 238; calcium salt of, 316; dicyclohexylamine salt of, 52, 54, 55, 73, 74, 100, 106, 203; and copper fungicides, 54, 100; and cryolite, 73, 74; reducing effectiveness of DDT, 106; and lead arsenate, 52, 73, 74, 203; and oil emulsions, 74, 141; and sulphur, 52, 179; and injury to plants, 52, 54, 74, 100; formulae containing, 52, 54, 73, 74, 100, 141, 179, 203, 313, 316; proprietary preparations containing, 52, 55, 106, 203.

Dinoderus bifoveolatus, fumigation against, in banana flour in Morocco, 65.

Dinoderus minutus, measures against, in timber in S. Africa, 12; factors affecting susceptibility of bamboo culms to, in Porto Rico, 219.

Dioryctria, in cones of Pinus edulis in U.S.A., 57.

Dioryctria abietella, in spruce cones in Finland, 40, 64.

Dioryctria splendidella, on pine in France, 257, 329, 371; oviposition by, 257.

diospyri, Aceria.

Diospyros kaki, new mite on, in California, 111; Cydia molesta developing on, 109.

Diotimana (Diotima) undulata, on Araucaria cunninghami in Queensland, 260. diparopsidis, Apanteles.

Diparopsis castanea (on cotton), in Nyasaland, 62, 168, 332, 333; in S. Rhodesia, 166; bionomics of, 168; measures against, 168, 332, 333.

Diphenylamine, effect of solvents on penetration of insect cuticle by, 87.

Diplogaster spp., bionomics of, infesting Coleoptera in U.S.A., 343, 344.

Diprion, 14.

Diprion frutetorum (see Gilpinia). Diprion hercyniae (see Gilpinia).

Diprion pini, bionomics and control of, on pine in Holland, 87, 88.

Diprion sertifer (see Neodiprion).

Diprionids, classification of, 14.

Dirphya princeps, measures against, on coffee in tropical Africa, 102.

disciger, Helopeltis bergrothi.

discoideus, Bracon.

Disodium Ethylene Bisdithiocarbamate, in mixed sprays, 239, 359.

dispar, Lymantria (Porthetria).

āisparis, Anastatus.

disputaria, Tephrina.

disstria, Malacosoma.

ditella, Tinea.

Dithane, 239, 359.

β-β-Dithiocyanodiethylether (see Thiocyanates).

divisus, Macrosteles.

Dixanthogen, effect of solvents on penetration of insect cuticle by, 87.

DN-111, 52, 55, 106, 203.

Dociostaurus maroccanus, bionomics of, in Spain, 265, 266, 268; parasites of, 265; bait for, 268.

Dock (see Rumex).

dohertyi, Leucoplema (Epiplema).

Dolichos lablab, Bruchus obtectus in, in Chile, 205; pests of, in Queensland, 125, 126.

Dolomite, against pests in stored beans and wheat, 126, 230; as a carrier for dusts, 89.

Dolycoris penicillatus, on cotton and lucerne in Central Asia, 154, 253.

domestica, Musca; Thermobia.

Dominica, summary of plant quarantine restrictions in, 292.

dominica, Rhizopertha.

Dominican Republic, new Bruchid in seeds of *Prosopis* in, **64**; summary of plant quarantine restrictions in, **112**. *Doralis* (see *Aphis*).

Dorcadion, effect of crop rotation on, in U.S.S.R., 157.

Dorcatoma bibliophagum, DDT against, in books in Brazil, 57.

dorsalis, Chalepus; Shirakia (see S. schoenobii).

dorsopictus, Spermophagus.

Dorymyrmex pyramicus, associated with Cinara carolina in S. Carolina, 311.

Douglas Fir (see *Pseudotsuga taxifolia*). Draeculacephala minerva, transmitting lucerne dwarf virus in California, 111. Dreft, 199.

dregii, Calidea.

Drosichiella tamarindus, bionomics of, in Punjab, 285.

drosophilae, Spalangia.

Dryocoetes alni, on Alnus incana in Finland, 9, 41.

dubia, Lypha.

duplicatus, Necremnus.

Dusts, use of oil-impregnated, 50, 195, 197, 248, 338; factors affecting coverage by, 310; equipment for applying, 138, 261, 358, 359, 379; application of, from aircraft, 89, 177, 178, 340; (inert), uses of, against pests in stored products, 35, 126, 188, 230; mode of action of, on insects, 60, 61, 86.

Dutch Elm Disease (see Ceratostomella

ulmi).

Dutox (barium fluosilicate and sodium fluoaluminate), dusting with, 29, 358, 359; and sulphur, 29, 358, 359; spraying with, 29, 201, 321; causing injury to peach, 321.

Dynaspidiotus britannicus, on Buxus in

California, 109.

Dysdercus (on cotton), native cotton susceptible to, in Nigeria, 169; in Nyasaland, 332; in Sudan, 166; bait-spray against, in Tanganyika, 167.

Dysdercus cardinalis, on cotton in

Tanganyika, 167.

Dysdercus fasciatus (on cotton), in S. Rhodesia, 166; in Tanganyika, 167.

Dysdercus intermedius, on cotton in S. Rhodesia, 166; erroneous record of, in Tanganyika, 167.

Dysdercus migratorius, on cotton in Tanganyika, 167.

Dysdercus nigrofasciatus, on cotton in Tanganyika, 167.

Dysdercus sidae, in W. Australia, 366. Dysdercus superstitiosus (on cotton), in S. Rhodesia, 166; in Tanganyika, 167.

E,

Ealincola, gen. n., 81.

Ealincola modesta, type of genus, 81.

Ealincola thoracica, 81.

Earias (on cotton), parasite of, in India, 284; in Tanganyika, 167.

Earias biplaga (on cotton), new parasite of, in Belgian Congo, 81; in Tanganyika, 167.

Earias cupreoviridis, experimental pairing of other species with, in India, 287.

Earias fabia, on cotton in Punjab, 282, 283, 287; effect of climate on distribution of, 283; crossing of, with other species, 287.

Earias huegeli, in Australia, 166, 366; on cotton, 166.

Earias insulana (on cotton), in Punjab, 282, 283, 287; in Sudan, 166; in Tanganyika, 167; distribution and resistance to climatic factors of, 283; crossing of, with other species, 287.

Earwig (see Forficula auricularia).

Easter Lily (see Lilium longiflorum). Echinochloa crusgalli, thrips on, in Hawaii, 274.

echinopus, Rhizoglypus.

Echium, Phenacoccus solani on, in California, 109.

Economic Entomology, book on, 10. (See Forest Entomology.)

Ecuador, Atta cephalotes in, 249.

edulis, Conophthorus.

efoveolatus, Bracon (Orthobracon).

egena, Autographa.

Egg-plant, Leptinotarsa decemlineata developing on, in Europe, 9, 10, 264; pests of, in U.S.A., 109, 240, 243; Ceratitis capitata not ovipositing on, 132; methyl bromide causing injury to packed fruit of, 109.

Egius platycephalus, predacious on Asterolecanium spp. in Porto Rico,

219.

Egypt, miscellaneous pests in, 18, 19. Elachertus (Elachistus) leucogramma,

parasite of Scolytids in Holland, 67. Elachertus (Elachistus) nigritulus, parasite of Cydia strobilella in Finland, 39,

elaeidis, Chevalierella.

Elaeis guineënsis (Oil Palm), pests of, in Belgian Congo, 82, 83.

Elasmus albopictus, parasite of Schoenobius bipunctifer in China, 152.

Elasmus flabellatus, parasite of Prays oleellus in Greece, 326.

Elasmus masii var. natalensis, parasite of Apanteles diparopsidis in E. Africa,

Electricity, apparatus for use of, against Lamellicorns in Uzbekistan, 155; barriers charged with, in rearing trays for *Gnorimoschema*, 345.

Electro-magnetic Radiation (see Light and Ultra-violet Rays).

Electrostatic Charges, of dusts, 310.

Elettaria cardamonum, Prodioctes haematicus on, in Travancore, 283.

Eleusine, Lygus simonyi on, in Uganda,

Eleusine coracana, Colemania sphenazioides attacking, in Mysore, 207.

Elgetol, 90.

elisus, Lygus.
Ellopia fiscellaria lugubrosa, bionomics and control of, in Br. Columbia, 377, 378, 379.

Elm, pests of, in Canada, 130; pests of, in U.S.A., 54, 236, 237; relation of Scolytids to Ceratostomella ulmi in, 54, 236.

Elm Leaf Beetle (see Galerucella luteola). Eloeidiphilos aliberti (see Epimorius adustalis).

elutella, Ephestia.

Elymus triticoides, Abacarus hystrix on, in California, 110.

emissaria, Scopula.

Emmalocera depressella, on sugar-cane in India, 188; failure to rear Bracon greeni on, 165.

Empoasca, on cotton in Africa, 166, 167, 169; DDT against, on tomato in New South Wales, 384.

Empoasca benedettoi (see E. lybica).

Empoasca decipiens, bionomics and control of, on Capsicum in Spain, 267.

Empoasca fabae (in U.S.A.), on cotton, 180; on leguminous plants, 244, 267, 312, 313, 338; on potato, 43, 53, 223, 224, 239, 337; dusts and sprays against, 43, 224, 239, 244, 267, 312, 313, 337, 338.

Empoasca lybica, bionomics and control of, on vines in Spain, 267; synonym

of, **267**.

Empoasca maculata, on cotton in Queensland, 166.

Empoasca mali (see E. fabae).

Empoasca solana, on sweet potato in Barbados, 159.

Empusa, infesting Aphids in Argentina and Uruguay, 300.

Empusa americana, infesting Parexorista caridei in Argentina, 300.

Empusa aphidis (see Entomophthora).

Empusa aulicae, infesting Charaeas graminis in Finland, 271.

Empusa grylli, observations on, infesting Zonocerus variegatus in Belgian Congo, 280.

Enarmonia caryana, measures against, on pecan in Georgia, 176.

Encyrtus, associated with Trioza erytreae in Nyasaland, 63.

engelmanni, Dendroctonus.

Engytatus geniculatus, bionomics of, attacking other insects on tobacco in N. Carolina, 347; insecticides against, on tomato in Hawaii, 276.

ensigera, Agalliana.

Entada gigas, Icerya on, in Belgian Congo, 82.

Entedon fuscipennis (see Dahlbominus). Entomology, book on, 113, (See Economic Entomology.)

Entomophthora aphidis, infesting Aphids in Argentina and Uruguay, 300.

Entomophthora sphaerosperma, infesting Pieris brassicae in Finland, 271.

Eotetranychus (see Tetranychus).

Epermenia chaerophyllella, on carrot in Finland, 38.

Ephestia, methyl bromide against, in dried figs, in Morocco, 65.

Ephestia cautella, measures against, in dried fruit, etc., in W. Australia, 365. Ephestia elutella, insecticides against, in

stored tobacco in Virginia, 338, 339;

synonymy of, 252.

Ephestia kuehniella, measures against, in dried fruit, etc., in W. Australia, 365; Chelonus texanus reared on, 3; nomenclature of, 3, 252; review of literature on, 288.

Ephestia sericaria (see E. elutella).

Ephialtes, parasite of Hylemyia laricicola in Finland. 38.

Ephialtes aplopappi, possible parasite of Cydia nigricana in Washington, 73.

Ephialtes brevicornis, possible parasite of Anthonomus pomorum in Holland, 94.

Ephialtes caudatus, failure to establish, against Cydia pomonella in New Zealand, 323.

Ephialtes glabratus (see E. strobilellae).

Ephialtes hispae, possible parasite of Cydia nigricana in Washington, 73. Ephialtes kangasi, sp. n., parasite of Pissodes spp. in Finland, 39, 40.

parasite Ephialtes | manifestator,

Necydalis major in Finland, 41. Ephialtes pictipes, parasite of Taenio-

campa gracilis in Finland, 42. Ephialtes pterophori, possible parasite of Cydia nigricana in Washington, 73.

Ephialtes sanguinipes (see Pimpla).

Ephialtes strobilellae, parasite of Cydia strobilella in Finland, 40.

ephippella, Argyresthia. Epiblema (see Eucosma).

Epidendrum (see Orchids).

Epilachna varivestis, measures and experiments against, on beans in U.S.A., 53, 138, 178, 182, 244, 310, 316, 317; on lima beans, 316, 317; relation between degree of infestation and damage by, 236; test of plant insecticides on, 144, 303, 304; resistance of, to DDT aerosols, 24.

Epilachna vigintioctopunctata, insecticides against, on cucurbits, etc., in W.

Australia, 365.

Epimorius adustalis, on Elaeis guineënsis in Belgian Congo, 83; synonym of,

Epineura rubens (parasite of Antestia), bionomics and value of, in Uganda, 278, 279.

Epiplagiops littoralis, introduced into U.S.A. against Listroderes obliquus, 179.

Epiplema dohertyi (see Leucoplema).

epiblemicidus, Apanteles.

Epitrimerus hystrix (see Abacarus).

Epitrix, insecticides against, on potato in Washington, 72.

Epitrix cucumeris, Harr. (in U.S.A.), on potato, 53, 181, 235, 236, 239, 243, 337; on tomato, 204; measures and experiments against, 53, 204, 235, 236, 239, 243, 337.

Epitrix cucumeris, auct. (see E. tuberis). Epitrix hirtipennis (parvula, auct.), measures against, on tobacco in N. and S. Carolina, 315, 356; on potato in Nebraska, 28; food-plants of, in Virginia, 128.

Epitrix subcrinita, food-plants of, in Montana, 181.

Epitrix tuberis, measures against, potato in U.S.A., 28, 29, 223, 224, 358, 359; bionomics of, 28, 29.

Epiurus (see Ephialtes).

Epochra canadensis, on currant and gooseberry in Oregon, 79.

eques, Torocampus (Lamachus).

Erannis tiliaria, trees attacked by, in Quebec, 173.

Eretmocerus serius, establishment of, against Aleurocanthus woglumi Mexico, 179.

ericae, Nysius.

erichsoni, Pristiphora (Nematus).

eridania, Laphygma (Prodenia).

Erigeron affinis, insecticidal constituent of, 359.

Eriocampoides (see Caliroa).

Eriococcus azaleae, fumigation against, on azalea in California, 109.

Eriophyes, 191; on olive in Spain, 267; scope of, 110.

Eriophyes mangiferae, measures against, on mango in Egypt, 19.

Eriophyes oleae, on olive in Greece, 326. Eriophyes pistaciae, on Pistacia terebinthus in Jugoslavia, 328.

Eriophyes pyri, sprays against, on pear in Washington, 73; systematic position of, 111.

Eriophyes sheldoni (see Aceria).

Eriophyes tulipae (see Aceria).

Eriophyes vermiformis (see Cecidophyes). Eriophyes vitis, on vines in California,

109; type of genus, 110.

Eriosoma lanigerum (Woolly Apple Aphis), in Britain, 184; in Denmark, bionomics of, in India, 228: DDT spray inin Mexico, 179; creasing infestation by, in U.S.A., 75, 107; biological race and control of, on Cotoneaster, 184: natural enemies and biological control of, 179, 228.

Eritrea, pests of vegetables in, 14, 154. Ernobius mollis, measures against, in timber in S. Africa, 12.

Erodium botrys, 163.

erosus, Ips.

Erwinia tracheiphila (see Bacillus).

Erycia basifulva, establishment of, against Coleoneura trichogramma in Fiji, 366.

Erycia rutila (see Masicera).

Erynnia nitida, parasite of Galerucella luteola in California, 374.

Erysiphe polygoni, 138, 317.

Erythrina, Aularches miliaris on, in Mysore, 187.

Erythrina abyssinica, Terastia meticulosalis on, in Belgian Congo, 81. erythrocephala, Calliphora.

Erythroneura, sprays against, on vines in Pennsylvania, 247, 248.

Erythroneura variabilis, DDT fog against, on vines in Arizona, 262.

erytreae, Trioza (Spanioza). Ethyl Cinnamate, in bait for Cydia

molesta, 322.

Ethyl Formate fumigation of wheat

Ethyl Formate, fumigation of wheat with, against insects, 231.

Ethylene Dibromide, fumigation tests with, against pests of stored products, 342, 343.

Ethylene Dichloride, emulsion of, against Aegeria exitiosa, 202, 336, 347, 348; alone and in fumigant mixtures against pests of stored cereals, 180, 230, 342; extracts of Achillea spp. prepared with, 49; as a solvent for DDT, 106, 245.

Etiella zinckenella, new parasite of, on Tephrosia and Crotalaria in Belgian Congo, 81.

etiellae, Bracon (Microbracon).

etrusca, Chionaspis.

Eublemma amabilis, parasite and control of, predacious on Laccifer lacca in Bihar, 164, 165.

Eublemma costimacula, new parasite of, predacious on Coccids in Belgian Congo, 81.

Eucalyptus, Gonipterus scutellatus on, in Mauritius, 101; Caedicia on, in New South Wales, 208.

Eucalyptus microtheca, resistance of timber of, to termites in W. Australia, 366.

Eucalyptus robusta, preferred by Gonipterus scutellatus in Mauritius, 102.

Eucalyptus Snout Beetle (see Gonipterus scutellatus).

Eucelatoria (see Blondelia).

Eucolaspis brunnea, apparatus for collecting emerging adults of, in New Zealand, 58.

Eucosma bobana, on Pinus edulis in U.S.A., 57.

Eucosma (Epiblema) strenuana, parasite of, in New Jersey, 142.

Eugenia edulis, new weevil on, in Argentina, 6.

Eugenia jambos, Anastrepha fraterculus on, in Mexico, 137.

Eugenia malaccensis, leaves of, as protective collars against molecrickets, 191.

Eugenia uniflora, Achaea attacking fruits of, in Nigeria, 15; Ceratitis capitata on, in Palestine, 133.

Eugenol, effect of evaporation on baits containing, 341.

Eulecanium bituberculatum, on fruit trees and Crataegus in Iraq and Persia, 16.

Eulecanium corni, 200; bionomics and control of, on Cotoneaster in Britain, 184.

Eulecanium fletcheri, sprays against, on Juniperus and Thuja in U.S.A., 200. Eulecanium (Lecanium) kunoense, on fruit trees and walnut in California, 374.

Eulecanium nigrofasciatum, on peach in Tennessee, 321.

Eulia loxonephes, bionomics of, on flax and Dahlia in Argentina, 298.

Eulimneria, parasite of Plutella maculipennis in Nyasaland, 63; parasite of, 63.

Eulimneria geniculata, parasite of Taeniocampa gracilis in Finland, **42.** Eulophus longulus, parasite of Dacus

oleae in Greece, 326.

Eulophus viridulus, spread of, parasitising Pyrausta nubilalis in Ontario, **320.** Eumaenas salaminia, attacking fruits in

Ceylon, **281**.

Eumerus spp. (Bulb Flies), effects of temperature on development of, in Finland, 42; keys to larvae and pupae of. 42.

Euonymus, winter food-plant of Aphis fabae in Britain, 113; Chrysomphalus bifasciculatus on, in California, 109.

Eupatorium, new Agromyzid on, in U.S.A., 57.

Eupelmus urozonus, parasite of Dacus oleae in Greece, **326**.

Euphorbia, Aphthona euphorbiae on, in Britain, 272.

euphorbiae, Aphthona.

euphorbiae, auct., Macrosiphum (see M. solanifolii).

Euphoriella, key distinguishing Euphorus from, 296.

Euphoriella marica, sp. n., hosts of, in Uganda, 296.

Euphorus, studies on, parasitising Capsids in Uganda, 13, 296; revision of African species of, 295, 296.

Euphorus anates, sp. n., parasite of Helopeltis in W. Africa, 296.

Euphorus nigricarpus, hosts of, in Uganda, **296.**

Euphorus praetor, sp. n., parasite of Lygus in Uganda, 296.

Euphorus sahlbergellae, hosts of, in W. Africa, 296.

Euphyllura olivina, on olive in Greece, 326.

Eupithecia, descriptions of larvae of Canadian species of, 159.

Eupithecia bilunulata, in spruce cones in Finland, 64.

Eupithecia filmata, 159. Eupithecia fletcherata, 159. Eupithecia gibsonata, 159. Eupithecia luteata, 159.

Eupithecia palpata, 159.

Eupithecia pini, in spruce cones in Finland, 64.

Euplectrus gopimohani, bionomics of, in Bengal, 164.

Euplectrus laphygmae, parasite of Laphygma exempta in Nyasaland, 63.

Euplectrus parvulus, bionomics of, in India, 285; synonymy of, 285.

Euplectrus platyhypenae, parasite of Cirphis unipuncta in Hawaii, 274. Euplectrus plecopterae (see E. parvulus).

Euproctis chrysorrhoea, auct. (see Nygmia phaeorrhoea).

Eupteromalus, parasite of Schoenobius bipunctifer in China, 152.

Eurhodope suavella, bionomics and control of, on Cotoneaster spp. in Britain, 185.

Eurydema festivum, on cotton in Bokhara, 154.

Eurydema oleraceum, insecticides against, on swedes in Germany, 89.

Eurygaster, diseases of, and their utilisation in Russia, 131.

Eurygaster integriceps (on wheat), effect of temperature on, in Russia, 48; apparatus for collecting, in Turkey, 48.

eurygraphus, Xyleborus.

Euryischia lestophoni, hosts of, in Australia, 82.

eurynotus, Vasates (Phyllocoptes).

Eurystylus risbeci, sp. n., on sorghum in Senegal, 96.

eurytheme, Colias.

Eurytoma, parasite of Hylemyia laricicola in Finland, 38; parasite of Apanteles transvaalensis in Nyasaland, 63; parasite of Ceuthorrhynchus assimilis in Washington, 76.

Eurytoma rosae, parasite of Dacus olear in Greece, 326.

Eurytoma syleptae, 63.

Euspermophagus (see Spermophagus). Eutelus strobicola, parasite of Dasyneura strobi in Finland, 39.

Eutermes triodiae, mounds of, in airfields in W. Australia, 366.

Eutetrapha tridentata, effect of sprays on, in elm logs in Massachusetts, 54, 55.

Eutettix tenellus, and curly-top of beet, etc., in U.S.A., 109, 353, 354, 374, 379, 380; failure of, to transmit Argentine curly-top virus, 379; wild food-plants and ecology of, 109, 353, 374; other bionomics of, 353, 354; measures and experiments against, 109, 374, 377.

Eutrombidium trigonum, destroying grasshopper eggs in S. Dakota, 232.

Euxoa messoria, in China, 116; larva of, 116.

Euxoa nigricans, on onion in Finland, 38. Euxoa tritici, on tomato and onion in Finland, 99, 100.

Euzophera bigella, on apple and vine in Bokhara, 156.

Evotomys glareolus, destroying Diprion pini in Holland, 88.

exempta, Laphygma.

Exenterus, utilisation of, against Gilpinia hercyniae in Canada, 130, 173. Exenterus abruptorius, parasite of Neodiprion sertifer in Finland, 9, 39.

Exenterus amictorius (marginatorius), parasite of Neodiprion sertifer in Finland, 39.

exigua, Laphygma.

exitiabilis, Pseudococcus (see P. njalensis).

exitiosa, Aegeria (Sanninoidea).

Exorista (see Zenillia).

Exosoma lusitanica, fumigation against, in cipollini in Morocco, 65.

Explosives, use of, to apply dusts in forests, 379.

exul, Sciopus.

Eysarcoris punctatus, experimentally predacious on Leptinotarsa decenlineata in Belgium, 213.

F.

fabae, Aphis (Doralis); Empoasca.
faber, Achaea.
fabia, Earias.
faceta, Antestia.
famelicus, Leskiopalpus.
farinae, Tyroglyphus.
fascialis, Hymenia (see H. recurvalis).
fasciata, Mordella.

fasciatus, Aeolothrips; Dysdercus; Oncopeltus; Trichomalus.

fasciculata, Oncopera. fasciculatus, Araecerus.

Fatty Acids, action of, on blowfly larvae, 85, 86.

fausta, Rhagoletis.

Feltia subterranea, DDT against, on melon in S. Carolina, 355.

femoralis, Hercinothrips; Tychius. femur-rubrum, Melanoplus.

fenestratus, Thyridanthrax.

ferganensis, Rhynchites auratus.

Fermate (see Ferric Dimethyldithio-carbamate).

Ferric Arsenate, effect on plants of soil containing, 355; of little value against Anthonomus grandis, 356.

Ferric Chloride, effect of, in DDT

solutions, 157.

Ferric Dimethyldithiocarbamate, and DDT, 106, 157, 229, 248; and other insecticides, 53, 106, 203, 220, 221, 308.

Ferric Fluoride, unsatisfactory against termites, 172.

Ferric Oxide, reducing effect on plants of calcium arsenate in soil, 355.

Ferrisia (Ferrisiana) virgata, experimentally transmitting swollen-shoot virus of cacao in Gold Coast, 102.

ferrugineus, Daeus (Strumeta); Laemophloeus.

festiva, Calonota (Pyronota).

festivum, Eurydema.

festucae, Macrosiphum (Myzus).

feytaudi, Matsucoccus.

ficorum, Gynaikothrips (Phloeothrips). Ficus anomani, Achaea faber piercing fruit of, in Nigeria, 15.

Figure carica (see Fig).

Ficus nitida, thrips on, in Egypt, 18, 19. ficus, Lepidosaphes.

Fig, Ceratitis capitata on, in Palestine, 133; Coccids on, in other countries, 16, 63, 285, 364.

Figs, Dried (see Fruit, Dried).

Fiji, miscellaneous pests in, 101, 366; list of Coccids of, 388; beneficial insects and biological control in, 101, 366.

filamentosus, Pseudococcus.

Filao (see Casuarina).

Filbert (see Corylus avellana).

Filipendula ulmaria, Taeniocampa gracilis on, in Finland, 42.

Filippia gossypii, sp. n., on cotton in Persia, 16.

Filippia oleae, on olive in Greece, 326. filmata, Eupithecia.

finitella, Tlascala.

Finland, new Aleurodid on Campanula in, 64; forest pests in, 9, 38, 39, 40,

41, 42, 64, 99, 271; miscellaneous pests in, 42, 271; pests of vegetables and root crops in, 37, 38, 99, 100, 271; studies on Thysanoptera of, 37, 64; Chlorophorus annularis in imported ski sticks in, 37; beneficial insects in, 9, 38, 39, 40, 41, 42; entomogenous fungi in, 271.

fiscellaria, Ellopia.

Fish, Dried, Dermestes frischi in, in Morocco, 65.

Fish Oil, uses of, in sprays, **50**, **51**, **220**, **304**.

Fish-oil Soap (see Soap).

flabellatus, Elasmus.

Flame-throwers, against locusts, 2. flava, Cosmophila (Anomis); Hoplo-

campa.

flavens, Frankliniella.

flaveola, Agromyza.

flavescens, Pulvinaria.

flavipes, Reticulitermes.

flavus, Thrips.

Flax, Aphthona euphorbiae on, in Britain, 272; Eulia loxonephes on, in Argentina, 298; Tipula paludosa on, in Denmark, 116; Thrips lini on, in Holland, 67; Nysius ericae on, in Russia, 36; relation of Cephus cinctus to, in U.S.A., 146, 147, 181; grasshoppers damaging, in S. Dakota, 232; preventing oviposition by Hylemyia coarctata, 216; resistance of, to wireworms, 330.

Flea-beetle, Potato (see Epitrix cucumeris and E. tuberis).

Flea-beetle, Tobacco (see *Epitrix* hirtipennis).

Fleahopper, Cotton (see Psallus seriatus). fletcherata, Eupithecia.

fletcheri, Eulecanium (Lecanium).

floccifera, Pulvinaria. floralis, Nemorilla.

Florida, Wasmannia auropunctata associated with other insects on Citrus in, 238; pests of field crops in, 104, 146, 147, 177, 335; parasite introduced against pineapple mealybug into, 179, 219; Lyncestis acontioides on Poinciana regia in, 356; pests of pecan in, 373; Scapteriscus spp. in seed-beds in, 192.

Flour, pests in, and their control, 63, 179, 180, 287; relation of insect damage to thiamine content of, 139, 140; methods for estimating populations of mites in, 98; effect of fumigants on baking qualities of, 66, 177; spraying with, against Tetranychus, 293; in mixed sprays, 50, 220, 356. Flour Mills, fumigation of, 179, 180.

Flour Mite (see Tyroglyphus farinae).

Flour Moth, Mediterranean (see Ephestia kuehniella).

fluctuata, Cidaria (Larentia).

Fluoranthene, treatment of wood with, against termites, 172.

Fluted Scale (see Icerya purchasi).

foedus, Melanoplus.

Fog Generator, DDT applied from, 262. Forest Entomology, book on, in Central Europe, 69.

Forests, Rhyacionia buoliana in, in Argentina, 371; Aphids in, in Central Asia, 156; pests of, in Canada, 129, 130, 159, 173, 263, 320, 360, 377; pests of, in Finland, 9, 38, 39, 40, 41, 64, 99; pests of, in France, 214, 257, 329, 371; Dendrolimus pini in, in Germany, 99; pests of, in Holland, 69, 87, 88; Sivex noctilio in, in New Zealand, 323; pests of, in Nyasaland, 63; pests of, in Queensland, 258, 260; Neodiprion sertifer in, in Sweden, 130, 131; pests of, in U.S.A., 54, 76, 158, 160, 176, 193, 233, 236, 237, 311.

Forficula auricularia, in Britain, 45; in Denmark, 8, 116; damaging drying tobacco leaves, 116; bionomics of, 8; baits for, 8; killed by DDT spray, 45.

Formaldehyde, uses of, against Tyrophagus lintneri, 77.

Formica rufa, destroying Diprion pini in Holland, 88.

formiciformis, Theocolax.

formosanus, Idiopterus (Micromyzus). fornication, Xyleborus fornicatus.

Fornicia, lists of species and hosts of, 80. Fornicia ghesquièrei, sp. n., parasite of Phorma pepon in Belgian Congo, 80.

forsteri, Lachnosterna. fortunana, Zeiraphera.

Fowls, Tenebrionids as food for, 328: risk of poisoning of, by arsenicals, 83. fragariae, Ancylis (see A. comptana); Aristotelia; Capitophorus; derma.

France, forest pests in, 214, 257, 329, 371; orchard pests in, 23, 160, 229, 349, 362, 370; pests of ornamental plants in, 348, 349, 350; vine pests in, 32, 33, 328, 362; Bruchids of, 329; Leptinotarsa decembineata in, 9, 10, 32, 83, 84, 327, 363; Locusta migratoria in, 34; miscellaneous pests in, 10, 349, 362; beneficial insects in, 10, 214, 361, 362, 363; fungous and microsporidian diseases of insects in, 10, 32, 33; book on insecticides in, 66; Coleoptera imported into, in stored products, 329, 363.

Frankliniella dampfi, on .Chrvsanthemum cinerariae folium in Nyasaland,

63.

Frankliniella flavens, bionomics of, in Hawaii, 276, 277.

Frankliniella fusca, insecticides against, on ground-nuts in Maryland, 338.

Frankliniella intonsa, on apple in Finland, 37.

Franklinothrips vespiformis, associated with other thrips on Cinchona in Porto Rico, 218.

fraterculus, Anastrepha.

fraterna, Lachnosterna.

fraxini, Hemiberlesia.

Fraxinus (see Ash).

frenchi. Coptotermes: Lebidiota. Freon-12, 239, 240, 241, 242, 243.

Freon-22, 241.

Frianite, as carrier for dust, 112.

frischi, Dermestes.

frit, Oscinella (Oscinosoma).

Frogs, destroying noxious insects, 152. frontalis, Dendroctonus.

Frontiniella parancilla, parasite of Tetralopha scortealis in Virginia, 305.

fronto, Sitona.

frugiperda, Laphygma.

Fruit (Dried), pests in, and their control, 65, 363, 365.

Fruit-flies, Cherry (see Rhagoletis cingulata and R. fausta).

Fruit-fly, Mediterranean (see Ceratitis capitata).

Fruit-fly, Mexican (see Anastrepha ludens).

frutetorum, Gilpinia (Diprion); Lamachus.

fullonia (fullonica), Othreis.

fulvicornis, Perilampus.

fulvipes, Triclistus. fulvus, Cyrtorhinus.

fumiferana, Harmologa (Archips).

Fumigants, equipment for applying: to stored products, 65; to field crops, 370; to soil, 318.

Fundella, revision of, 104.

Fundella argentina, distribution of, 104; confused with F. pellucens, 104.

Fundella pellucens (cistipennis), synonymy and distribution of, on pulse crops, 104.

Fungi, Entomogenous, 10, 30, 31, 33, 39, 63, 73, 102, 111, 171, 181, 187, 213, 219, 227, 237, 260, 271, 280, 284, 293, 299, 300, 322; utilisation of, 33, 131.

Fungi, Injurious, 5, 41, 59, 121, 130, 135, 138, 190, 203, 220, 221, 224, 240, 248, 293, 317, 338; relation of insects to, 33, 54, 55, 171, 236, 246, 321, 370, 388. (See also Macrophoma.)

furcifer, Hieroglyphus (see H. banian). Furniture, Coleoptera in, 11, 12, 103,

116, 333.

Furniture Beetle (see Anobium punctatum).

Furs (see Rabbit Skins).

Fusarium lateritium, infesting Phylloxera vitifoliae in France, 33; utilisation of, 33.

fusca, Frankliniella; Lachnosterna.

fuscicollis, Ageniaspis.

fuscipennis, Dahlbominus (Entedon, Microplectron).

fuscus, Anthrenus; Ptilinus.

Fygran A, 188.

G.

Galeopsis tetrahit, destruction of, against Aphids in Maine, 226.

Galerucella luteola, parasite of, on elm in California, 374.

Galiniera, Antestia lineaticollis on, in

Uganda, 278.

Galinsoga parviflora, encouragement of, in coffee plantations against Helopeltis, 154.

Galleria mellonella, measures against, in bee-hives in Punjab, 297.

Gambrus stokesi (see Agrothereutes).

Gammexane, y isomer of benzene hexachloride (q.v.), 256.

Gardenia, Orthezia insignis on, in S. Rhodesia, 33; Pseudococcus citri on, in U.S.A., 79.

Gargaphia solani, aerosols against, on egg-plant in U.S.A., 243.

Garlic, Thrips tabaci on, in India, 118; Acrolepia assectella on, in Switzerland,

214, 216. Gastrimargus musicus, in Australia, 1, 2, 365, 387; bionomics of, 1, 2;

measures against, 2, 387. Gecinus viridis, destroying Zeuzera

pyrina in England, 35.

Gelatin, in adhesives for lead arsenate sprays, 220.

Gelatine Capsules, use of, for studying insect parasites, 315.

Gelatinosporium fulvum, 130.

Gelechia vepretella, on Cotoneaster in Europe, **185**.

gelechiae, Bracon (Microbracon).

gemmatalis, Anticarsia.

Genaparlatoria pseudaspidiotus, synonymy of, 205.

geniculata, Eulimneria (Limnerium).

geniculatus, Engytatus.

Georgia, miscellaneous pests in, 176, 194, 304, 309; pests of peach in, 335, 336.

Geranium, Phenacoccus colemani on, in Virginia, 31.

germanica, Blattella; Vespa.

Germany, Psila rosae on carrot in, 68; pests of crucifers in, 89, 97; miscellaneous pests in, 99, 328; wireworms in, 92.

germari, Brachynema.

Gesapon (Guesapon), 43, 44.

Gesapon no. 18, 143.

Gesarol, 7, 93, 94, 95, 171, 327, 334.

Gesarol A, **334.** Gesarol A-3, **247, 248.**

Gesarol A-20, 105.

Gesarol AK-20, 80, 105, 106, 247, 248.

Gesarol P, 333, 334.

ghesquièrei, Apanteles; Fornicia.

gibberus, Gonipterus.

gibbosus, Ligyrus.

gibbus, Bruchophagus. gibsonata, Eupithecia.

gifuensis, Macrocentrus.

gigas, Urocerus.

Gilpinia, parasite of, on spruce in Finland, 39; systematic position of,

Gilpinia abieticola, on spruce in Finland,

Gilpinia frutetorum, bionomics of, on Pinus spp. in Connecticut, 236, 237. Gilpinia hercyniae (on spruce), in Canada,

129, 130, 173; in U.S.A., 237; parasites and biological control of, 130, 173, 237.

Gilpinia polytoma, on spruce in Finland,

Gilpinia variegata, systematic position of, 14.

Gipsy Moth (see Lymantria dispar). glabratus, Ephialtes (see E. strobilellae). Gladiolus, Thrips simplex on, in Hawaii, 277; pests of, in New South Wales, 385.

glaseri, Neoaplectana.

Gliricidia, Pseudococcus brevipes on, in Hawaii, 274.

glomeratus, Apanteles.

Glossista infuscata (see Cytherea).

Glossonotus crataegi, sprays against, on quince in New York, 246.

gloveri, Lepidosaphes (Mytilococcus).

Glue, in mercurous chloride seed dressing,

Glyceryl Oleate, as an emulsifier, 238.

Glyphipteryx achyloessa, measures against, on Dactylis glomerata in New Zealand, 323.

Glyphodes unionalis (see Margaronia). Glypta haesitator, established against Cydia nigricana in Br. Columbia, 73;

not spreading into Washington, 73. Glypta rufiscutellaris, reared on Gnorimoschema operculella for use against Cydia molesta in California, 110.

Glyptina cerina, intercepted at Hawaii in aircraft from California, 2.

Glyptoscelis squamulata, on vines in California, 109.

gnidiella, Cryptoblabes.

Gnorimoschema heliopa, parasites of, on

tobacco in Nyasaland, 62.

operculella (Potato Gnorimoschema Moth), in Australia, 382, 387; in Bermuda, 103, 300, 301; in Eritrea, 154; in Morocco, 65; insecticides against, on field potatoes, 382, 387; methyl bromide against, in stored potatoes, 65; on tobacco and tomato, 301; parasites and biological control of, 301; rearing and use of, as host for parasites, 110, 142, 305, 306, 314, 345, 354; tests of insecticides on, 134. godmani, Pantomorus.

Gold Coast, parasites of Capsids on cacao in, 296; Coccids transmitting swollenshoot of cacao in, 46, 102; Pseudococcus njalensis on other plants in, 46.

Goldenrod (see Solidago). Golf Courses (see Turf)

Gonipterus gibberus, characters distinguishing G. scutellatus from, 102.

Gonipterus scutellatus, bionomics of, on Eucalyptus in Mauritius, 101; natural enemies and biological control of, 102; characters of, 102.

Gooseberry, pests of, in Britain, 35, 273; Nematus ribesii on, in Holland and Denmark, 59, 116; pests of, in Oregon, 79.

Gooseberry Sawfly (see Nematus

vibesii).

gopimohani, Euplectrus. Gorse (see *Ulex europaeus*).

Goryphus nursei, bionomics of, in India,

gossypiella, Platyedra (Pectinophora).

gossypii, Aphis; Filippia. Gossypium (see Cotton).

gowdeyi, Haplothrips.

gracilis, Dacnusa; Phyllocoptes; Taeniocampa.

gradatus, Systoechus.

graefi, Aegeria (Sanninoidea).

Grain Beetle, Rust-red (see Laemophloeus ferrugineus).

Grain Borer, Lesser (see Rhizopertha dominica)

graminis, Charaeas.

graminum, Pediculoides (Pediculopsis); Toxoptera.

granaria, Calandra.

granarium, Macrosiphum (see M. avenae); Trogoderma.

granarius, Aphidius.

Granary Weevil (see Calandra granaria). grandiosella, Diatraea.

grandis, Anthonomus.

Grape Cane Girdler (see Ampeloglypter ater).

Grape Hyacinth (see Muscari).

Grape Leafhoppers (see Erythroneura).

Grapefruit, fruit-piercing moths attacking, in Nigeria, 14, 15; Ceratitis capitata on, in Palestine, 132, 133; Aceria sheldoni on, in Queensland, 126; pests of, in U.S.A., 24, 238; quarantine treatment of fruit of, against introduction of Anastrepha into U.S.A. from Mexico, 112.

Graphognathus (see Pantomorus).

Grapholitha (see Cydia).

Graphops pubescens, bionomics of, on strawberry in Kansas, 200, 201.

Grass, Cocksfoot (see Dactylis glomerata). Grass, Crested Dogstail (see Cynosurus

cristalus).

Grass, Sudan (see Sorghum sudanense). Grass, Timothy (see Phleum pratense).

Grasses, pests of, in S. Australia, 324: Charaeas graminis on, in Finland, 271; insects damaging, in France, 349; Anaphothrips obscurus on, in Hawaii, 275; pests of, in New Zealand, 20, 209, 210; Crambid destroying thatching of, in Nyasaland, 207; Conoderus spp. on, in U.S.A., 129, 183; cereal and sugar-cane pests on, 63, 119, 157, 171, 185, 275, 312, 368, 375, 381; relation of locusts and grasshoppers to, 1, 34, 177; suggested use of, to foster Coccinella septempunctata, 228. (See Turf.)

Grasshopper, Jola or Deccan Colemania sphenarioides).

Grasshopper, Red-legged (see Melanoplus femur-rubrum).

Grasshoppers (see Locusts).

Greece, pests of olive and their natural enemies in, 326.

greeni, Bracon (Microbracon).

gregaria, Schistocerca.

Grevillea, Crypticerya rosae on, in New South Wales, 82.

Grevillea robusta, Drosichiella tamarindus on, in Punjab, 285.

grisea, Macquartia.

grisescens, Lydella stabulans.

grönblomi, Megastigmus.

grossa, Bombomima.

Ground-nuts, pests of, in U.S.A., 244, bromide (stored), methyl against pests in, 65, 66; (shells), unsatisfactory in bait for locusts, 294.

Gryllodes sigillatus, killed by residue of DDT aerosol, in greenhouses Maryland, 240.

Gryllotalpa, plants attacked by, Egypt, 19, 20; bait for, 19.

Gryllotalpa hexadactyla, measures against, in Trinidad, 191.

Gryllus campestris, damaging pastures in France, 349.

Guam, Nephotettix apicalis intercepted in aircraft from, 2.

Guatemala, new Coccid introduced into California on orchids from, 112.

Guava, Conotrachelus psidii on, in Brazil, 6; Cydia molesta larvae experimentally developing on, in California, 109; Anastrepha fraterculus on, in Mexico, 137, 138; Achaea attacking fruits of, in Nigeria, 15; Ceratitis capitata on, in Palestine, 132, 133.

Guayule (see Parthenium argentatum).

Guesapon (see Gesapon).

Gum Arabic, in solid bait for *Cydia* pomonella, 308; as emulsifier for dichlorethyl ether, 78.

guttata, Parnara.

guttulosa, Austracris.

Gynaikothrips ficorum, causing galls on Ficus nitida in Egypt, 18, 19; insects predacious on, 19.

Gynaikothrips uzeli, characters distinguishing G. ficorum from, 19.

Gynandropsis pentaphylla, race of Myzus persicae on, in Nyasaland, 62, 206. Gypsum, as a carrier for dusts, 73, 89.

H.

Habrobracon (see Bracon).

Habrocytus tenuicornis, possibly parasitic on Anthonomus pomorum in Holland, 94.

Habrolepis, parasite of Coccids on Aleurites in Nyasaland, 63.

Hadrobregmus nitidus, on Sorbus aucuparia in Finland, 41.

Hadronotus antestiae, parasite of Pentatomids in Uganda, 278.

haematicus, Prodioctes.

haemorrhoidalis, Athous (see A. obscurus); Heliothrips.

haesitator, Glypta.

Haiti, new Bruchid on *Prosopis* in, 64. Hakea arborescens, resistance of timber of, to termites in W. Australia, 366.

halli, Nilotaspis (Coccomytilus).

Halotydeus destructor, in S. África and Australia, 162; bionomics of, 162; effect of, on yield of clovers, 162, 163. Haltichella, possibly hyperparasitic on

Tetralopha scortealis in Virginia, 305. Halticus bracteatus, on Aralia in Cali-

fornia, 374.

Halticus citri, on Talinum in Bermuda, 103.

Hambletonia pseudococcina, introduced into Florida from Porto Rico against Pseudococcus brevipes, 179, 219.

Hamitermes rubriceps, mounds of, in airfields in W. Australia, 366.

Haplothrips, on Chrysanthemum cinerariaefolium in Nyasaland, 63.

Haplothrips angustus, on sugar-cane in New Caledonia, 275.

Haplothrips gowdeyi, on sugar-cane in New Caledonia, 275.

Harlequin Cabbage Bug (see Murgantia histrionica).

Harmolita tritici, measures against, on wheat in Montana, 181.

Harmologa fumiferana (on conifers), forms of, in Canada, 129, 173, 263, 320; sprays against, in U.S.A., 176; parasite of, 263.

Harrisina, of U.S.A., 112.

Harrisina brillians, bionomics and control of, on vines in California, 112.

Hartigia trimaculata, measures against, on greenhouse rose in Ohio, 79. Harvester Termites (see Hodotermes).

Hawaii, pests of pineapple in, 134, 135, 307; sugar-cane pests in, 170, 275; pests of vegetables in, 149, 274, 275, 276, 277, 308; miscellaneous pests in, 274–277; natural enemies and biological control of pests in, 2, 3, 171, 274; pests intercepted in aircraft in, 2, 171; other pests intercepted in, 183; summary of plant quarantines in, 292; pests imported into U.S.A. in rabbit skins from, 274, 275; quarantine treatment against introduction of fruit-flies into U.S.A. from, 190.

hawaiiensis, Thrips.

Hawthorn (see Crataegus).

Heat, uses of, against pests, 165, 180, 325, 365; (vapour), treatment of fruit with, against Anastrepha, 112. (See Solar Heat, Steam and Water, Hot.)

Hedychium gardnerianum, Pentalonia nigronervosa on, in greenhouses in

France, 362.

Hedylepta indicata (see Lamprosema). Helianthus tuberosus, Pseudaulacaspis pentagona on, in France, 362.

Helina calyptrata, predacious on Aesiotes notabilis in Queensland, **260.** heliopa, Gnorimoschema (Phthorimaea).

Heliothis armigera (obsoleta), in Australia, 126, 165, 231, 366, 387; in Barbados, 159; in Hawaii, 3, 274, 276; in S. Rhodesia, 166; in Tanganyika, 167, 168; in U.S.A., 3, 138, 147, 177, 178, 179, 316, 317, 354, 355, 357; on cotton, 165, 166, 167, 168, 178, 179; favouring infestation of cotton by Earias, 167; on leguminous plants, 126, 138, 167, 316, 317; on maize, 3, 147, 159, 167, 177, 355; susceptibility of strains of maize to, 357; on tomato, 177, 276, 354; on other plants, 231, 274, 387; bionomics of, 357, 387; parasites and biological

control of, 3, 179, 274; use of bacterial disease of, 177; other measures and experiments against, 126, 138, 144, 177, 178, 179, 231, 247, 276, 316, 317, 354, 355, 387.

Heliothis peltigera, on Carthamus tinctorius in Uzbekistan, 155.

Heliothis virescens, measures against, on cotton in Brazil, 334.

Heliothrips haemorrhoidalis (on ornamental plants), in Hawaii, 277; sprays against, in Ohio, 78.

Hellula phidilealis, bionomics and control of, on cabbage in Argentina, 372. Hellula undalis, dusts against, on

cabbage in Queensland, 231.

Helopeltis, new parasite of, on cacao in Gold Coast, 296; on cotton in Sudan and Nigeria, 166, 169.

Helopeltis bergrothi vars. disciger and rubrinervis, on Cinchona in Nyasaland, 63.

Helopeltis orophila, on Cinchona in Belgian Congo, 153, 323; relation of, to weeds and hedge plants, 153.

Helorimorpha coffeae (see Aridelus).

helvolus, Metaphycus.

Hemerocampa leucostigma, on elm and maple in Nova Scotia, 130.

Hemiberlesia camelliae (see H. rapax). Hemiberlesia degenerata, on Camellia in California, 109.

Hemiberlesia fraxini, sp. n., on Fraxinus velutina in California, 112.

Hemiberlesia lataniae, on Camellia in California, 109.

Hemiberlesia rapax, on Camellia California, 109.

hemipterus, Carpophilus.

hemisphaerica, Saissetia (see S. coffeae). Hemitarsonemus (Tarsonemus) bionomics of, on tea in Ceylon, 366; measures against, on Begonia in Holland, 95; characters and synonymy of, 95, 366.

Hemlock (see Tsuga).

Hemlock, Mountain (see Tsuga mertensiana).

Hemlock, Western (see Tsuga hetero-

phylla).

Hemp, Tipula paludosa on, in Denmark, Hieroglyphus nigrorepletus on, in India, 47; Icerya purchasi on, in Persia, 16; Nysius ericae on, in Persia, 16; Russia, 36.

Hemp Nettle (see Galeopsis tetrahit).

herbarium, Catorama.

Hercinothrips femoralis, on banana in Hawaii, 274; DDT aerosols against, on greenhouse plants in Maryland, 240.

herculeana, Clavaspis. herculeanus, Camponotus. hercyniae, Gilpinia (Diprion).

Herse cingulata, armyworm movement of, on sweet potato in Florida, 146. hesperidum, Coccus.

Hesperis matronalis, experimentally infected with turnip virus, 182.

hesperus, Lygus.

Hessian Fly (see Mayetiola destructor). Heterobostrychus brunneus, measures against, in timber in S. Africa, 12.

Heterodera, D-D mixture against, in pineapple fields in Hawaii, 135.

Heteronychus sanctae-helenae, plants attacked by, in New South Wales, 208, 382, 386; measures and experiments against, 208, 382.

heteropus, Atanycolus (Bracon).

Heterostylus aliberti, sp. n., on Cola in Ivory Coast, 14.

heveae, Tenuipalpus.

Hexachlorethane, ineffective in repellent spray against Anthonomus pomorum,

Hexachlorphenol, treatment of wood with, against termites, 172.

hexadactyla, Gryllotalpa.

Hibiscus, Flatid on, in Bermuda, 190. Hibiscus cannabinus, Hemiptera on, in

Uzbekistan, 155.

Hibiscus esculentus, Sylepta derogata on, in Belgian Congo, 82; legislation against introduction of Platyedra gossypiella in seed of, into U.S.A., 291.

Hickory Shuckworm (see Enarmonia caryana).

Hide Beetle (see Dermestes maculatus). Hides, Dermestes maculatus attacking,

Hieroglyphus banian (furcifer), nigrorepletus confused with, 47.

Hieroglyphus nigrorepletus, bionomics of, in India, 47.

hilaris, Aulacophora.

Himacerus lativentris, predacious on Pentatoma rufipes in Britain, 115.

Hippodamia, effect on, of nicotine dust against Chromaphis juglandicola California, 145.

Hippodamia convergens, predacious on Macrosiphum onobrychis in California, 111.

hirticula, Lachnosterna (Phyllophaga). hirtipennis, Epitrix.

hispae, Ephialtes (Scambus).

hispanica, Ramburiella.

hispidulus, Sitona.

histrionica, Murgantia.

Hodotermes, biology and control of, in S. Africa, 32.

Holcocera pulverea, experiments with heat against, predactious on Laccifer lacca in Bihar, 165.

Holland, pests of conifers in, 69, 87, 88, 93; pests of greenhouse and ornamental plants in, 95, 99, 367; orchard pests in, 67, 90, 91, 93, 94, 367; pests of vegetables in, 67, 68, 90, 92, 93; miscellaneous pests in, 59, 67, 95; list of insects susceptible to derris in, 97; Aphids and virus diseases of plants in, 90, 99, 367; beneficial insects in, 67, 88, 94, 95, 99; Rhizoglyphus echinopus on hyacinth bulbs imported into France from, 350.

Homoeosoma, bionomics and control of, on sorghum in Queensland, 387.

Honduras, British, Atta cephalotes in, 323.

Honey, used for feeding insects, 138, 314.

Hoop Pine (see Araucaria cunninghami). Hoplia, measures against, on Citrus in Costa Rica, 128.

Hoplocampa flava (on plum), bionomics of, in France, 229.

Hoplocampa minuta, in Sweden, 37; sprays against, 37, 229.

Hoplocampa testudinea, effect of wetting agents in sprays against, on apple in Britain 6

Hops, Tetranychus telarius on, in Britain, 100, 273; book on pests of, in Britain, 101.

Hornet (see Vespa crabro).

horni, Inostemma.

horticola, Anomala (Phyllopertha).

hortuellus, Crambus.

Houseflies (see Musca domestica).

howardi, Aleurothrixus.

Howardia biclavis, fungus infesting, on tea and Aleurites in Nyasaland, 62.

howitti, Aphodius. huegeli, Earias. humeralis, Sitona.

Humidity, relations of locusts to, 124; relations of other insects to, 8, 12, 17, 21, 22, 62, 121, 122, 123, 132, 156, 184, 222, 223, 259, 265, 284, 306; appartus for studying reactions of insects to, 154.

Hyacinth, Rhizoglyphus echinopus on imported bulbs of, in France, 350.

hyalinus, Liorhyssus.

Hyalochloria denticornis, sp. n., on cotton and beans in Peru, 128.

hybneri, Piezodorus.

Hydrangea, Diaspidiotus ancylus on, in

California, 109.

Hydrocyanic Acid Gas, against Aceria sheldoni, 127; against Coccids, 26, 108, 150, 289-291, 302, 363, 364, 373, 374; use of oil sprays together with, 289-291, 364; studies on resistance of stages and races of Coccids to, 26, 150, 302; against pests in mushroom

houses and greenhouses, 77, 78; against *Camponotus* spp. in houses, 27; againsts pests of stored grain and flour, 179, 180; fumigation of harvested fruits with, 363, 364; recovery of, from fumigated insects, 150, 151. (See Calcium and Potassium Cyanides.)

Hydrogen Chloride, liberation of, by DDT and benzene hexachloride, 255. Hylastes rufipes, relation of, to Cerato-

stomella ulmi in elm in U.S.A., 54, 55, 236; measures against, 54, 55.

Hylemyia anthracina, in spruce cones in Finland, 38, 40, 64; bionomics of, 64; natural enemies of, 38, 64.

Hylemyia antiqua, bionomics and control of, on leek in Holland, 67, 68, 92.

Hylemyia brassicae (on cabbage), in Canada, 319; in Spain, 266; bionomics of, in Washington, 73; measures against, 73, 319.

Hylemyia cilicrura (in U.S.A.), on conifer seedlings, 76, 160; on vegetables and maize, 375; soil fumigation

against, 76.

Hylenyia coarctata, bionomics of, on cereals in Britain, 216, 367, 368; on grasses, 368; measures against, 216, 217.

Hylemyia laricicola, parasites of, on Larix sibirica in Finland, 38.

Hylobius abietis, measures against, on Pinus pinaster in France, 371; damaging beans in Holland, 93.

Hylotrupes bajulus (in building timber), bionomics of, in S. Africa, 12; in Denmark, 12, 116.

Hylurdrectonus piniarius, bionomics of, on Araucaria in Queensland, 260.

Hylurgopinus (see Hylastes).

Hymenia recurvalis (fascialis), in Australia, 169, 365, 386; in Hawaii, 274; in Porto Rico, 219; on beet or allied plants, 169, 219, 274, 365, 386; on soy bean, 219; bionomics of, 169; parasites of, 219, 274; insecticides against, 169, 386.

hyoscyami, Corizus; Pegomyia.

Hyocyamus niger, Leptinotarsa decemlineata developing on, 264; infusion of, against Tetranychus turkestani, 49. Hypera, on lucerne in Tadzhikistan, 253. Hypera postica (see H. variabilis). Hypera punctata (in U.S.A.), on lucerne,

111, 322, 346; on Medicago hispida, 322; natural enemies of, 194, 322;

DDT against, 346.

Hypera variabilis (on lucerne), in U.S.S.R., 157, 253; in U.S.A., 111, 181; parasite of, 181; crop rotations ineffective against, 157.

hyperici, Chrysomela (Chrysolina).

D

Hypericum perforatum, establishment of Chrysomela hyperici against, in New Zealand, 323.

hypermnestra, Othreis.

Hyphaene miyaa, locust flails made of, 280.

Hyphantidium sericarium (see Ephestia elutella).

Hypnus abbreviatus, cultural measures against, in Connecticut, 235.

Hypocampsis contorticornis, parasite of Dasyneura strobi in Finland, 39.

Hypogastrura armata, in tobacco plant beds in N. Carolina, 347.

Hypurus bertrandi, bionomics and control of, in France, 349.

Hyra polyandra, spray of sap of, against Aphids, 304.

hystrix, Abacarus (Callyntrotus, Epitrimerus).

I.

Icerya, parasites of, on orange, Tephrosia and Entada in Belgian Congo, 82.

Icerya nigroareolata, parasite of, on coffee and Alchornea in Belgian Congo,

Icerya purchasi, in Argentina, 299; in Belgian Congo, 82; in France, 361; in India, 387; in Nyasaland, 63; in Persia, 16; on Aleurites fordi, 63; on Citrus, 16, 82; other food-plants of, 16, 287; natural enemies of, 82, 299, 361.

Icerya tremae, new parasite of, on Artocarpus in Belgian Congo, 82.

iceryae, Cryptochetum.

Ichneumon moderator, L. (see Nemeritis). Idaho, Cydia molesta in, 373; Eutettix tenellus and curly-top of beet in, 353, 354; Lygus spp. in, 204.

Idechthis (see Nemeritis). Idioaspis (see Yangambia).

Idioaspis macarangae (see Yangambia vesiculata).

Idiopterus formosanus, on onions in Hawaii, 274; probably imported from California, 274.

idiota, Pentodon.

ignorata, Dasyneura (Perrisia).

Ilacuris laticollis, effect of, on *Aesiotes* notabilis in *Araucaria* in Queensland, 260.

ilicis, Lachnosterna.

Illinois, miscellaneous pests in, 105, 292, 318.

illustvata, Chilosia.

illutana, Cydia (Laspeyresia). imbilensis, Temnosternus. immaculata, Scutigerella.

immersa, Dadobia.

imparata, Boarmia (Ascotis) selenaria. Imperial Institute of Entomology, directorship of, 225.

incertellus, Schoenobius (see S. bipunctifer).

inconsequens, Taeniothrips.

inconspicua, Sturmia.

inconstans, Sciara (Lycoria).

Incurvaria quadripunctella (morosa), bionomics and control of, on rose in France. 348.

India, immature stages of Cossids and Metarbelids of, 288; pests and disease of sugar-cane in, 297; parasites of Lepidoptera in, 284, 285; experiments with Teleonemia scrupulosa in, 287; Anthrenus spp. imported into Britain in myrobalans from, 59. (See also under the various Provinces.)

Indiana, Cydia pomonella on apple in, 175; cereal pests in, 4, 5, 177, 250; Diabrotica melanocephala and its natural enemies in, 148.

indica, Apis.

indicata, Lamprosema (Hedylepta).

Indigofera arrecta, possible food-plant of Helopeltis orophila in Belgian Congo, 153.

indistincta, Apate.

Indo-China, new weevil on Casuarina in, 362.

infausta, Aglaope.

infuscata, Ĉytherea (Glossista); Ormenis. inops, Parlagena.

Inostemma, parasite of Contarinia juniperina in Bermuda, 103, 191.

Inostemma horni, parasite of Sitodiplosis mosellana in Washington, 251.

inquilina, Spermatolonchaea.

Insect Cuticle, factors affecting penetration of, by insecticides, etc., 60, 61, 85-87; action of inert dusts on, 60, 61, 86; transpiration through, 60, 61.

insectella, Tinea.

Insecticides, books on composition, analysis and use of, 20, 66, 151; tests of organic compounds as potential, 43, 160, 255, 256, 359; studies on plants as sources of, 137, 217, 218, 301, 303, 304, 359, 381; bibliography of plants as sources of, 32, 192; methods of testing, 43, 254, 255; factors affecting permeability of insect cuticle by, 60, 85–87; principles of action of, 85–87.

Insects, book on food habits of, 256; factors influencing increase of, 128; conditions governing distribution of, in air, 184; principles of action of insecticides on, 85–87. (See Insect Cuticle.)

insignis, Orthezia.
insulana, Earias.
integriceps, Eurygaster.
intermedius, Dysdercus.
interpunctella, Plodia.
intonsa, Frankliniella.

Inula viscosa, Empoasca lybica overwintering on, in Spain, 267.

inunctus, Tetrastichus.

Iowa, Calomycterus setarius in, 105; Cydia molesta in, 373; Pyrausta nubilalis on maize in, 177, 323; pests of vegetables in, 57, 337.

ipomeae, Dendrothripoides.

Ips (on pines), in Connecticut, 237; measures against, in France, 371.

Ips bidentatus, on Pinus pinaster in France, 371.

Ips borealis, measures against, on conifers in Quebec, 173.

Ips erosus, on Pinus pinaster in France, 371.

Ips lecontei, on Pinus edulis in U.S.A., 57.

Ips sexdentatus, on Pinus pinaster in France, 371.

Iraq, Coccids in, 16; migration of Schistocerca gregaria into Arabia from, 15.

Iris, tolerant to methyl bromide, 110.

irregularis, Valanga.

Isaria, infesting Aesiotes notabilis in Queensland, 260.

iselyi, Melanoplus foedus.

isis, Tetrastichus. italicus, Calliptamus.

Itoplectis obesus, parasite of Acleris variana in Br. Colombia, 379.

Ivy, Chrysomphalus bifasciculatus on, in California, 109.

J.

Jaapiella medicaginis, on lucerne in Denmark, 116.

Jack Beans (see Canavalia ensiformis). Jack Pine (see Pinus banksiana).

Jack Pine (see *Pinus banksıana*).

Jack Pine Budworm (see *Harmologa fumiferana*).

jacksoni, Pulvinaria.

Jamides phaseli (see Lampides).

Japan, Calomycterus setarius in, 105; pests imported into other countries from, 37, 350.

Japanese Beetle (see Popillia japonica). japonica, Popillia.

japonicum, Trichogramma.

Jarring, to dislodge insects from trees, 93, 95, 120, 128, 201, 336.

Jasminum sambac, new Coccid on, in Iraq, 16.

Jerusalem Artichoke (see Helianthus tuberosus).

Jobo (see Spondias).

Jola Grasshopper (see Colemania sphenarioides).

jonesi, Setoptus (Platyphytoptus).

juglandicola, Chromaphis.

Juglans nigra, Diaspidiotus ancylus on, in California, 374.

Juglans regia (see Walnut).

Jugoslavia, pests of Pistacia terebinthus in, 327, 328.

Jujube (see Zizyphus jujuba).

juniperi, Pseudococcus.

juniperina, Contarinia.

Juniperus, Coccids on, in Kansas, 200; Monoctenus on, 14.

Juniperus bermudiana, pests of, in Bermuda, 103, 190, 191.

Jute (fibres), unsatisfactory as bait for locusts, 294; (plants) (see *Corchorus* spp.).

Jute Stem Weevil (see Apion corchori). juvencus, Sirex.

K.

Kahikatea (see *Podocarpus dacrydioides*) Kale, *Phyllotreta* spp. on, in Britain, **114**. Kalo, **200**.

Kalotermes, in tea bushes in Ceylon, 211; measures against, infesting plants, 211.

Kalotermes brevis, tests of timber preservatives against, in Porto Rico, 172, 345.

Kalotermes piceatus, measures against, in house in Hawaii, 275.

Kalotermes rainbowi, in coconut palms in Danger Is., 211.

Kalotermes samoanus, in coconut palms in Danger Is., 211.

kangasi, Ephialtes (Pimpla).

Kansas, cereal pests in, 192, 323, 375; Coccids on *Juniperus* in, 200; new Agromyzid on onion in, 57; orchard pests in, 198, 373; pests of strawberry in, 199, 200.

Kaolin, as carrier for dusts, 23, 100,

169, 208, 231, 354, 382, 386. Kapok (see Ceiba pentandra).

Kashmir, pests of peach in, 293.

kellyi, Sarcophaga.

Kentucky, Conoderus spp. in, 129, 183; Pyrausta nubilalis on maize in, 177; Tyloderma fragariae on strawberry in, 292.

Kenya Colony, new weevil on cucurbits in, 206; locusts in, 294, 352; pests of timber in, 103; precautions against

introduction of Cydia pomonella into, 189.

Kerosene, and creosote, spraying with, against Coleoptera in elm logs, 54, 55; atomised sprays of insecticides in, 301, 302, 377; uses of, with DDT, 58, 245, 262, 302, 310, 385; as a solvent, 58, 301, 302, 310, 385; effect of evaporation on baits containing, 341.

Kerosene Emulsion (see Oil Emulsions). Khaya grandifoliola, Boxia khayae on, in Gold Coast, 296.

khayae, Boxia.

kirbyi, Oeketicus (Oeceticus).

Kirganelia, Icerya purchasi on, in India, 287.

Kohlrabi, experimentally infected with turnip virus, 182.

Kok-saghyz (see Taraxacum kok-saghyz). Krym-saghyz (see Taraxacum megalorrhizon).

Kryocide, 358.

kuehniella, Ephestia.

kunoense, Eulecanium (Lecanium). kuvanae, Ooencyrtus.

L.

Laccifer lacca (Lac Insect), parasites and control of predators of, in Bihar, 164, 165.

Lachnosterna, adult activity of species of, in Connecticut, 237.

Lachnosterna anxia, in U.S.A., 181, 237; Xylaria infesting, 181.

Lachnosterna crenulata, 237.

Lachnosterna forsteri, 237.

Lachnosterna fraterna, 237. Lachnosterna fusca, 237.

Lachnosterna hirticula, in U.S.A., 210, 237; bacillus causing milky disease of Odontria not infecting, 210.

Lachnosterna ilicis, 237.

Lachnosterna marginalis, 237.

Lachnosterna micans, 237.

Lachnosterna smithi (see Clemora).

Lachnosterna tristis, in Connecticut, 237.

lacteus, Coptotermes.

Lactuca, Agromyza pusilla on, in U.S.A., 57. (See Lettuce.)

Laemophloeus, bionomics of, in flour in India, 288.

Laemophloeus ferrugineus, fumigation against, in stored grain in Canada, 320.
Laemophloeus minutus, in dried bananas

in France, 363; in bulk wheat in Victoria, 230; resistant to carbon bisulphide, 230.

Laemostenus terricola (see Pristonychus). laevigatus, Alphitobius; Camponotus. Lamachus eques (see Torocampus).

Lamachus frutetorum, parasite of Neodiprion sertifer in Finland, 9, 39.

Lamia textor, on Salix caprea in Finland,

Lampides phaseli, dust against, on Canavalia and Dolichos in Queensland, 126.

Lampronia quadripunctella (see Incurvaria).

Lamprosema indicata, on cowpeas in Belgian Congo, 81; on soy bean in Porto Rico, 218; parasites of, 81, 219. lamprosemae, Stantonia.

lanigerum, Eriosoma (Schizoneura).

Lantana, experiments with Teleonemia scrupulosa and, in India, 287; value of T. scrupulosa against, in Queensland, 33; measures against, as food-plant of Orthezia insignis in S. Rhodesia, 33.

lapathi, Cryptorrhynchus (Crypto-

rrhynchidius).

Laphygma eridania, on sweet potato in Barbados, 159; effects of antisera on larvae of, 193.

Laphygma exempta, in Hawaii, 2, 3, 170; on cereals and Cynodon in Nyasaland, 63; biochemical differences between phases of, 47; parasites and biological control of, 3, 63,

Laphygma exigua, bionomics and control of, on Corchorus spp. in Bengal, 163, 164; in Hawaii, 3; on maize and lucerne in U.S.A., 3, 111; parasites and biological control of, 3.

Laphygma frugiperda (on maize), in Barbados, 159; in U.S.A., 3, 357; strains of maize in relation to, 357; parasites of, 3.

laphygmae, Euplectrus; Meteorus;

Rogas.
Larch, pests of, in Canada, 129, 173;
Hylemyia laricicola on, in Finland,
38; sawflies on, in Holland, 69.

Larentia fluctuata (see Cidaria).

laricella, Coleophora.

laricicola, Hylemyia (Chortophila).

laricinellae, Chrysocharis.

laricis, Anoplonyx; Pristiphora (Lygae-onematus).

Larix (see Larch).

Larra, parasite of mole-crickets in Trinidad, 191.

Larra americana, establishment of, against Scapteriscus vicinus in Porto Rico, 32.

Lasioderma serricorne, damaging antimosquito smoke spirals in Fiji, 101; measures and experiments against, in stored tobacco in Nyasaland and Virginia, 62, 338, 339; natural enemies of, 62.

441

Lasioptera berlesiana, transmitting fungus affecting Dacus oleae in olives in Greece, 326.

Laspeyresia (see Cydia). lataniae, Hemiberlesia.

lateralis, Phytomyza (Napomyza).

Latheticus oryzae, measures against, in bulk wheat in Victoria, 230.

Lathyrus odoratus, experiments with Bruchus obtectus and, in Chile, 205. laticollis, Ilacuris.

latinotus, Ancistrotermes.

lativentris, Himacerus (Nabis).

latro, Lixus.

latus, Hemitarsonemus (Tarsonemus).

Lawns (see Turf).

Lead, Lepidoptera damaging, 271, 272. Lead Acetate, and sodium arsenate, in preparation of nascent lead arsenate,

Lead Arsenate, against Citrus pests, 208, 282; against cotton pests, 293, 334, 355, 356; against pests of forest and shade trees, 176, 191, 372; against orchard pests, 4, 8, 20, 37, 51, 52, 73, 75, 90, 106, 107, 175, 195, 196, 197, 198, 199, 201, 202, 203, 220, 221, 229, 248, 249, 266, 308, 321, 322, 336, 364, 365; strains of Cydia pomonella resistant to, 107; against pests of small fruits, 59, 199, 200, 272, 273, 292; against pests of nut trees, 273, 309, 365; against pests of vegetables, 23, 72, 84, 100, 126, 169, 204, 209, 231, 313, 327, 354, 372, 383, 385, 386; against vine pests, 54, 234, 235, 247; against pests on other plants, 78, 105, 164, 356; against Lamellicorns in soil, 20, 176; use of, against Alphitobius laevigatus, 101; in baits, 8, 191, 200, 282, 292; dip containing, 204; dusting with, 23, 72, 78, 84, 126, 164, 169, 195, 197, 208, 231, 354, 383, 385, 386; carriers for, 164, 195, 208, 231, 354, 383, 386; oil-impregnated dusts of, 195, 197; formulae for, in sprays, 4, 20, 23, 50, 52, 59, 73, 75, 90, 101, 106, 107, 164, 175, 195, 196, 198, 199, 200, 201, 202, 203, 208, 209, 220, 221, 234, 235, 247, 248, 249, 266, 304, 308, 309, 313, 322, 334, 336, 356, 365; aeroplane distribution of spray containing, 176; adhesives and spreaders for, 50, 164, 175, 196, 199, 219–221, 248, 266, 304, 308, 334, 356; and Bordeaux mixture, 53, 199, 204, 235, 247, 249; and calcium arsenate, 355; and DDT, 106, 157, 197, 247; and dinitro-compounds, 52, 73, 74, 203; and ferric dimethyl dithiocarbamate, 203, 220, 308; and Lethane B-72, 247; and lime, 20, 23, 59, 164, 196, 197, 198, 201, 202, 203, 204, 248,

322, 336, 364, 383; and lime-sulphur, 249, 309; "inverted" spray mixtures of oils and, 50, 51; and Paris green, 334; and rotenone, 356; and sulphur, 52, 78, 197, 198, 202, 203, 221, 248, 322; and zinc sulphate, 50, 196, 199, 201, 202, 203, 204, 309, 336; and injury to plants, 50, 51, 197, 199, 201, 204, 209, 321, 322, 336; effects of fertilisers on injury to peach by, 322; problems of residues of, 59, 199, 203, 372; basic form of, 201, 309, 354; preparation of nascent form of, 84; other insecticides compared with, 72, 84, 106, 203, 231, 313, 321.

Lead Oxide, reducing effect on plants of calcium arsenate in soil, 355.

Leaf-cutting Ants (see Atta).

Leafhopper, Beet (see Eutettix tenellus). Leafhopper, Grape (see Erythroneura). Potato (see Empoasca Leafhopper,

fabae). Leafhopper, White Apple (see Typhlo-

cyba pomaria). Leather, Anobium punctatum attacking,

Lecanium, auct. (see Eulecanium). lecontei, Ips; Neodiprion.

ledereriella, Batrachedra.

Leek, Acrolepia assectella on, in Britain, 69; pests of, in Holland, 67, 68, 92; A. assectella on, in Switzerland, 214, 215, 216.

Leek Moth (see Acrolepia assectella). Lemon, pests of, in California, 24, 25, 79, 80; Pseudaonidia trilobitiformis on, in Nyasaland, 63; Aceria sheldoni on, in Queensland, 126; Ceratitis capitata not ovipositing on, 132.

with Lentils, experiments Bruchus obtectus and, in Chile, 205.

Leopard Moth (see Zeuzera pyrina).

lepesmei, Ochropyga. lepida, Parasa.

Lepidiota frenchi, factors favouring, on sugar-cane in Queensland, 1.

Lepidosaphes beckii (on Citrus), in Belgian Congo, 82; sprays against, in New South Wales, 384; in Persia, 16; in Spain, 269.

Lepidosaphes camelliae, fumigation against, on Camellia in California, 109. Lepidosaphes ficus, on fig in Iraq, 16.

Lepidosaphes gloveri (on Citrus), fumigation against, in California, 110; intercepted in Belgian Congo, 82; in Persia, 16; in Spain, 269; Uganda, 82.

Lepidosaphes machili, on orchids in U.S.A., 109, 315; spray against, 315.

Lepidosaphes newsteadi, on Juniperus bermudiana in Bermuda, 190; spray against, 191.

Lepidosaphes pistaciae, on Pistacia vera in Persia, 16.

Lepidosaphes tuberculatus (see L. machil). Lepidosaphes ulmi, spray against, on Cotoneaster horizontalis in Britain, 184, 185; on apple in Persia, 16.

Lepisma saccharina, testing attractiveness of fabrics and paper to, in

U.S.A., 127.

Leptinotarsa decemlineata (Potato Beetle), in Belgium, 213; in Britain, 161; in Canada, 264, 319; in France, 9, 10, 32, 83, 84, 327, 363; in Spain, 100, 263, 264; in Switzerland, 327; in U.S.A., 100, 178, 204, 238, 264, 355; on egg-plant, 9, 264; on tomato, 10, 204, 264; experiments with other plants and, 264; bionomics of, 9, 214, 265; effect of climate on distribution of, 264; natural enemies of, 10, 213; resistance of, to certain Tachinids, 363; measures and experiments against, 83, 84, 100, 161, 178, 204, 238, 258, 304, 319, 327, 355.

Leptohylemyia (see Hylemyia).

Leptomastix bifasciatus, parasite of Pseudococcus citri in Nyasaland, 63.

Leptomias bimaculatus, on lucerne in Tadzhikistan, 253.

Leptostylus praemorsus, on Juniperus bermudiana in Bermuda, 191.

Leskiopalpus famelicus (parasite of Diatraea saccharalis), imported into Porto Rico from Brazil, 219; in Trinidad, 219.

Lespedeza spp., pests of, in U.S.A., 105, 183, 304; susceptibility of, to Tetralopha scortealis, 304.

lestophoni, Euryischia.

Lethane Preparations, 199, 247, 248, 313, 356. (See Thiocyanates.)

Lettuce, Aphids and virus diseases of, in Britain, 70; Thrips angusticeps on, in Finland, 37; Thysanoptera on, in Hawaii, 274, 277; Listroderes obliquus on, in Queensland, 387; Macrosteles divisus on, in U.S.A., 243; effect of naphthalene on germination of, 192.

leucogramma, Elachertus (Elachistus).

leuconotus, Anthores.

leucophthalmus, Philaenus.

Leucopis, predacious on Cinara carolina in S. Carolina, 311.

Leucoplema dohertyi, new parasite of, on coffee in Belgian Congo, 81.

leucopterus, Blissus.

leucostigma, Hemerocampa.

Leurostethus raphiae, gen. et sp. n., on Raphia in Ivory Coast, 14.

libocedrus, Augonomoctenus on, 14.

Licania rigida, Conotrachelus spp. on, in Brazil, 6.

licaniae, Conotrachelus.

lienardi, Achaea.

Light, effect of, on rotenone, 50; (artificial), fruit-piercing moths attracted to, 14.

Light-traps, for Lachnosterna, 237; Coleoptera not attracted to, 20, 187; for moths, 152, 153, 209, 262, 372, 381; ineffective against fruit-piercing moths, 282.

ligneus, Semanotus.

Ligyrus gibbosus, on Parthenium argen-

tatum in Arizona, 26.

Lilies (Lilium), Rhizoglyphus echinopus on imported bulbs of, in France, 350; thrips on, in Hawaii, 277; tolerant to methyl bromide fumigation, 110.

Lily of the Valley (see Convallaria

majalis).

Lima Bean Vine Borer (see Monoptilota

pergratialis).

Lima Beans, pests of, in U.S.A., 57, 74, 104, 316, 317; cyclohexanone causing injury to, in aerosols, 244.

limacina, Caliroa (Eriocampoides). Lime, dusting with, against Caliroa limacina, 365; ineffective against Calandra oryzae in stored cereals, 121: as a carrier for dusts, 23, 54, 73, 126, 127, 149, 164, 231, 236, 239, 301, 303, 383; sabadilla dusts prepared with, 301, 312, 313; addition of, to arsenical sprays, 20, 59, 164, 196, 198, 201, 202, 203, 204, 220, 221, 248, 322, 336; paint of lead arsenate and, against *Capnodis* on apricot, 364; addition of, to other sprays (see also Bordeaux Mixture), 106, 199; question of compatibility of DDT with, 106, 239, 255; infestation of jute by Apion reduced by manuring with superphosphates and, cucurbits scorched by, 149; formulae containing, 20, 23, 126, 127, 164, 175, 196, 197, 198, 199, 202, 236, 248, 312, 313, 322, 336, 383.

Lime-sulphur, against Coccids, 25, 51, 73, 127; against Dasyneura affinis, 18; against Jassids, 293; against Paratrioza cocherelli, 29, 358, 359; against mites, 73, 77, 91, 92, 100, 127, 161, 371, 384; against Scirtothrips citri, 25; ineffective against Anthonomus pomorum, 274; formulae for, 29, 51, 77, 100, 127, 358, 359, 384; formulae containing, 29, 73, 358, 359; and arsenicals, 29, 73, 358, 359; increasing water-soluble

arsenic in lead-arsenate sprays, 204; and DDT, 106, 157, 245, 255; and Dithane, 359; and oil emulsion, spraying with, 73; fruit trees injured by association of oils with, 91.

Limnerium geniculatum (see Eulim-

neria).

Limonius agonus, bionomics of, on potato and tobacco in Connecticut, 235.

Limonius californicus, populations of, in relation to crops in Washington, 74; naphthalene against, 74.

Limonius canus, populations of, in relation to crops in Washington, 74; naphthalene against, 74.

Limothrips cerealium, on tomato and orchids in California, 374.

orchids in California, 374.

Lindingaspis rossi, spray against, on

Citrus in New Zealand, 21.

linearis, Oeme (see O. rigida).

lineaticollis, Antestia.

lineatus, Agriotes.

lineellus, Sitona.

Linen Bags, penetration of, by Calandra oryzae, 121.

lineolatus, Adelphocoris.

lini, Thrips.

Linseed Oil, in cryolite spray, 72.

lintneri, Tyrophagus.

Liorhyssus hyalinus, on Hibiscus and Abutilon in Uzbekistan, 155.

Liquidambar styraciflua, Coccid on, in California, 109.

liquidambaris, Diaspidiotus.

Liriodendron tulipifera, injection of chemicals into, for timber preservation, 56.

Lissorhoptrus bosqi, sp. n., bionomics of, on rice, etc., in Argentina, 171.

Lissorhoptrus oryzae and simplex, characters distinguishing L. bosqi from, 171.

Listroderes, natural enemies of, in Argentina and Uruguay, 104, 299.

Listroderes obliquus (on vegetables), measures and experiments against, in Australia, 23, 383, 386, 387; in Oklahoma, 310; introduction of parasites into U.S.A. against, 104, 179; bionomics of, 23.

Lithocolletis blancardella, bionomics and control of, on apple in Massachusetts,

52.

Litomastix truncatella, parasite of Autographa egena in California, 30.

littoralis, Epiplagiops. litura, Prodenia.

Lixophaga diatraeae (parasite of Diatraea saccharalis), failure to establish, in Barbados, 159; imported into Mexico, 179.

Lixus latro, sp. n., on cabbage in Eritrea, 14.

Lizards, destroying mole-crickets, 191. Lobotrachelus bidentipes, sp. n., on mango in Senegal, 206.

Locust, Desert (see Schistocerca gregaria). Locusta migratoria, in W. Australia, 365; swarms of, in France, 34.

Locusta migratoria manilensis, breeding centres of, in China, 293.

Locusta migratoria migratorioides, biochemical differences between phases of, 47.

Locustana pardalina, biochemical differ-

ences between phases of, 47.

Locusts and Grasshoppers, in Africa, 10, 63, 165, 265, 280, 294, 351, 352; in S.-W. Asia, 15, 16, 352; in Australia, 1, 123, 124, 365, 387; in Canada, 227; in China, 293; in France, 34; in India, 47, 187, 207; in Spain, 265, 266, 268; in U.S.A., 111, 141, 176, 180, 231, 232, 301, 312, 339, 340; breeding centres and ecology of, 124, 265, 293; phases of, 15, 34, 265, 268; biochemical and physiological studies on phases of, 47, 350, 351; influence of weather, etc., on feeding by, 340; migrations of, 1, 2, 15, 34; periodicity of, 15, 207, 227; bacterial and fungous diseases of, 10, 11, 124, 227, 280; insect enemies of, 2, 16, 180, 227, 265; mite destroying eggs of, 232; birds destroying, 47; toad destroying, 2; measures and experiments against (see also under Baits), 2, 11, 16, 47, 177, 180, 207, 227, 232, 280, 301, 302.

loftini, Chilo.

Loganberry, Byturus urbanus on, in Britain, 34.

Lolium perenne, relation of Diptera on cereals to, in Britain, 368, 381; Odontria zealandica on, in New Zealand, 20.

Lolium rigidum, experiments with Halotydeus destructor and, in W. Australia,

162.

Lonchocarpus, DDT compared with, in sprays against Byturus, 34; insecticidal action and constituents of species of, 217, 281, 303; cultivation of, 281, 303. (See Cubé and Timbo.)

Lonchocarpus chrysophyllus, 217. Lonchocarpus urucu, 303.

Lonchocarpus utilis, 217, 303.

Long Beans (Vigna unguiculata sesquipedalis), pests of, in Queensland, 125, 126.

longior, Tyrophagus (Tyroglyphus). longipennis, Scirtothrips.

longispina, Parlatoreopsis (Chionaspis) (see P. chinensis).

longulus, Eulophus; Sitona.

Lophocateres pusillus, in dried bananas in France, 363.

Lophyroplectus luteator, parasite of Neodiprion sertifer in Finland, 9, 39.

Lophyrus abieticola (see Gilpinia). Lophyrus polytomus (see Gilpinia). Lophyrus sertifer (see Neodiprion).

Loquat, Cydia molesta on, in California, 109; Anastrepha fraterculus experimentally infesting, in Mexico, 138; Dacus tryoni in, in New South Wales,

Louisiana, cotton pests in, 178, 303, 311; pests of Solidago leavenworthii in, 309; Diatraea saccharalis on sugar-cane in, 177; virus diseases of turnip in, 182.

lounsburyi, Aspidioti phycus; Trionymus. 'Aspidiotiphagus; Meta-

loxonephes, Eulia.

Loxostege similalis, aerosols and dust against, on spinach in U.S.A., 243.

Loxostege sticticalis, experiments with fungous and bacterial diseases of, in Russia, 131; parasite of, 295.

Loxotropa melanagromyzae, sp. n., parasite of Agromyza in Belgian Congo, 96.

Lucerne, 304; Lygus spp. on, in Alberta, 204; pests of, in W. Australia 265. tralia, 365; Jaapiella medicaginis on, in Denmark, 116; Taeniocampa gracilis on, in Finland, 42; pest of, in U.S.S.R., 36, 131, 132, 156, 157, 253; pests of, in U.S.A., 74, 105, 111, 177, 180, 181, 204, 227, 242, 244, 260, 261, 312, 322, 346, 374; insects affecting seed production of, 131, 132, 253, 260, 261; Jassid transmitting virus disease of, 74; pollination of, by wild bees, 260; unsatisfactory in baits for grasshoppers, 339.

lucicola, Anomala.

luctuosus, Oxycarenus. ludens, Anastrepha.

lugubrosa, Ellopia fiscellaria.

luminosus, Pyrophorus.

luniger, Syrphus.

Lupin (Lupinus), Nysius ericae on, in Russia, 36; preferred food-plant of Anabrus simplex, 25; recommended to protect coffee from Dasus simplex,

luridus, Troilus.

lusitanica, Exosoma (Malacosoma).

luteata, Eupithecia.

luteator, Lophyroplectus.

luteoapicalis, Cosmopteryx attenuatella.

luteola, Galerucella. lybica, Empoasca.

Lycidocoris, on Cinchona in Belgian

Congo, 323. Lycidocoris modestus (see Ealincola). Lycidocoris thoracicus (see Ealincola). lycopersici, Phyllocoptes.

Lycoria inconstans (see Sciara).

Lyctus brunneus, measures against, in timber in S. Africa, 12.

Lydella stabulans var. grisescens, spread of, parasitising Pyrausta nubilalis in Ontario, 320.

Lygaeonematus (see Pristiphora).

Lygus, on coffee in Belgian Congo, 280, 281; varieties of cotton resistant to, in Uganda, 166; on lucerne in Utah, 260, 261; measures and experiments against, 260, 261, 281.

Lygus campestris, bionomics and control of, on celery in Massachusetts, 53.

Lygus elisus, in Alberta, 204; U.S.A., 111, 204, 224, 346; on lucerne, 111, 204; on Parthenium argentatum, 346: insecticides against, on potato, 224; bionomics of, 204.

Lygus hesperus, in Alberta, 204; in U.S.A., 111, 135, 204, 346; on lucerne, 111, 204; on Parthenium argentatum, 135, 346; bionomics of, 204, 346; parasite of, 346; dusts against, 135.

Lygus oblineatus, 191; in U.S.A., 141, 243, 312; sprays against weeds and, in peach orchards, 141; experiments against, on potato and lucerne, 243.

Lygus pabulinus, killed by DDT on

carrots in Britain, 45.

Lygus pratensis, L., on cotton and Hibiscus in Central Asia, 154, 155.

Lygus pratensis, auct. (see L. oblineatus). Lygus sallei, on Parthenium argentatum in California, 135.

Lygus simonyi (vosseleri), on cotton in Uganda, 13, 14, 296; on grain crops, 13; factors affecting infestation by, 13, 14; parasites of, 13, 296; measures against, 14.

Lymantria dispar (in U.S.A.), natural enemies and disease of, 193, 194; effects of temperature on eggs of, 233; aeroplane spraying against, 176.

Lyncestis acontioides, bionomics and control of, on Poinciana regia in Florida, 356.

Lypha, new species of, in Canada, 263. Lypha dubia, not found in Canada, 263. Lypha maculipennis, in Canada, 263.

Lypha setifacies, bionomics of, in Canada, 263.

Lysol, spraying with, against Eurydema oleraceum, 89.

Μ.

macarangae, Idioaspis (see Yangambia vesiculata).

Macaroni, Calandra oryzae in, in W. Australia, 365.

machili, Lepidosaphes.

Macquartia grisea, resistance of Leptinotarsa decemlineata to parasitism by,

in France, 363.

Macrocentrus ancylivorus (in U.S.A.), parasitising Ancylis comptana, 141, 142, 322; utilisation and value of, against Cydia molesta, 110, 141, 142, 204, 234, 305, 322, 373; failure to establish, against C. nigricana, 73; rearing of, on Gnorimoschema operculella, 110, 142, 305, 306, 314, 345; investigations on host relations and overwintering of, 141, 142; microsporidian infecting, 314.

Macrocentrus delicatus, hosts of, in U.S.A., 305, 354; rearing of, on Gnorimoschema operculella, 354.

Macrocentrus gifuensis, introduced parasite of Pyrausta nubilalis in Canada, 320.

Macrodactylus suavis, bionomics and control of, on Citrus in Costa Rica, 128.

Macrodactylus subspinosus, Bombomima grossa predacious on, in U.S.A., 194. Macrodiprion, systematic position of, 14. Macrophoma dalmatica, Cecidomyiid transmitting, in olives in Greece, 326; value of, against Dacus oleae, 326.

Macrosiphum avenae, experiments with Aphidius granarius against, on wheat in Wales, 46.

Macrosiphum euphorbiae, auct. (see M. solanifolii).

Macrosiphum festucae, experiments with Aphidius granarius against, on wheat in Wales, 46.

Macrosiphum granarium (see M. avenae). Macrosiphum onobrychis (in U.S.A.), on lucerne, 111, 177, 241, 346; on peas, 133, 177, 241; biological races of, 133, 134; natural enemies of, 111; insecticides against, 177, 241, 242, 346; effects of dusts on reproduction of, 144.

Macrosiphum pisi (see M. onobrychis). Macrosiphum rosae, measures against, on greenhouse roses in Ohio, 78.

Macrosiphum solani, in Britain, 70; in Holland, 99; in Maine, 226; transmitting virus diseases of plants, 70, 99, 226; weed eradication against, 226.

Macrosiphum solanifolii, in Holland, 99; on gladiolus in New South Wales, 385; on potato in U.S.A., 226, 239, 253, 337; transmitting virus diseases of Narcissus and potato, 99, 226;

measures and experiments against, 226, 239, 243, 337, 385.

Macrosteles divisus, insecticides against, on potato and lettuce in U.S.A., 224, 243.

Macrozamia, Aonidiella aurantii on, in W. Australia, 364.

maculata, Empoasca.

maculatus, Camponotus; Dermestes; Neoplectrus.

maculipennis, Lypha; Plutella.

Madagascar, Bruchid imported into France in beans from, 329.

Madras, Apion corchori in, 117; Icerya purchasi in, 287; termites in, 291. (See Travancore.)

Magdalis, in elm logs in U.S.A., 55,

236; sprays against, 55.

Magdalis barbicornis, on pear and medlar in Britain, 272.

Magdalis ruficornis, bionomics of, on plum in Britain, 272.

Magnesite, against pests of stored products, 126, 230, 388; against cabbage pests, 231; and nicotine sulphate, 231.

Magnesium Arsenate, effect on plants of soil containing, 355; of little value against Anthonomus grandis, 356.

Magnesium Oxide, dust of, protecting stored wheat from insects, 177.

Mahogany, use of, in tests of termite repellents, 172, 345.

Maidenhair Fern (see Adiantum).

maidis, Aphis.

Maine, Rhagoletis pomonella on apple in, 306; pests of vegetables in, 182, 243, 244; Aphids and virus disease

of potato in, 226.

Maize, pests of, in Barbados, 159; Pyrausta nubilalis on, in Canada, 320; pests of, in Hawaii, 3, 274, 277; effect of Orius majusculus on, in Holland, 95; pests of, in India, 47, 117, 118; Sesamia vuteria on, in Morocco, 257; Heteronychus sanctaehelenae on, in New South Wales, 382; studies on interplanting of cotton with, in relation to Diparopsis castanea in Nyasaland, 168, 333; spread of Heliothis armigera to cotton from, in Tanganyika, 167, 168; pests of, in U.S.A., 3, 53, 74, 129, 136, 137, 147, 177, 180, 183, 192, 221, 232, 236, 237, 240, 244, 250, 312, 323, 355, 357, 375; treatment of ears of, against H. armigera, 147, 177; resistance to pests of strains and hybrids of, 221, 222, 357; time of sowing, in relation to pests, 118, 180; refuse of, as carrier for locust baits, 294.

Maize (Stored), pests in, and their control, 21, 252, 333, 334, 341, 342, 354, 355; effects of fumigants on germination of, 342; in baits for molecrickets, 191; (meal), in baits for Heliothis armigera, 179.

majalis, Amphimallon.

major, Necydalis. majusculus, Orius.

Malacosoma disstria, trees defoliated by, in Nova Scotia, 129.

Malacosoma lusitanica (see Exosoma).

Malacosoma neustria, bionomics and control of, on Cotoneaster in Britain, 185.

mali, Aphelinus; Chrysobothris; Dasyneura; Empoasca (see E. fabae); Psylla; Scolytus.

Malva, Listroderes obliquus on, in W.

Australia, 23.

Mamestra brassicae, experiments with predacious Pentatomids and, in Belgium, 213; polyhedral disease of, 213; tests of insecticides on, 254, 255.

Mamestra oleracea (see Polia). Mamey (see Mammea americana).

mamillatus, Conotrachelus.

Mammea americana, leaves of, as protective collars against mole-crickets, 32, 191; insecticidal properties of, 218.

mancus, Agriotes.

Manganese Salts, effects of, in leadarsenate sprays, 204, 220.

Mangel, insects and virus diseases of, in Britain and Argentina, 113, 379, 380. mangiferae, Eriophyes; Parlatoria (see Genaparlatoria pseudaspidiotus).

Mango, Eriophyes mangiferae on, in Egypt, 19; Anastrepha ludens in, in Mexico, 343; moths attacking fruit of, in Nigeria and Ceylon, 14, 15, 381; new Curculionid on, in Senegal, 206; vapour-heat treatment of fruit of, against introduction of Anastrepha into U.S.A. from Mexico, 112; poison baits for moths prepared from fruit or juice of, 15.

manifestator, Ephialtes.

Manila Hemp (see Musa textilis).

manilensis, Locusta migratoria.

Manitoba, pests of field crops in, 159, 174; pests of stored grain in, 320; Lypha setifacies in, 263.

Mantis religiosa, use of, against forest pests in Nova Scotia, 130.

Maple, pests of, in Canada, 130, 173. *Margarodes*, on wheat in Persia, 16.

Margarodes tritici, sp. n., on wheat in Turkey, 256.

Margaronia unionalis, on olive in Greece, 326.

marginalis, Lachnosterna.

marginatorius, Exenterus (see E. amictorius).

marginatus, Melanoplus.

marginellus, Calliptamus italicus.

marginiventris, Apanteles.

Margosa (see Melia azadirachta). marica, Euphoriella.

Marigold, Heteronychus sanctae-helenae on, in New South Wales, 208.

maritimus, Pseudococcus.

marlatti, Phoenicococcus.

maroccanus, Dociostaurus.

Marsh Copal (see Copaifera demeusei).

Marsh Mallow (see Malva).

martorelli, Amblycerus.

Maruca testulalis, bionomics and control of, in Queensland, 125, 126.

Maryland, pests of leguminous plants in, 240, 241, 242, 244, 304, 338, 346; Tyrophagus lintneri on mushrooms in, 76; orchard pests in, 106, 107, 245; miscellaneous pests in, 105, 239-245, 314.

Masicera rutila, erroneous records of, in N. America, 263.

masii, Elasmus.

Massachusetts, miscellaneous pests in, 51-55, 105, 193, 194; beneficial insects in, 193, 194.

Mastotermes darwiniensis, damaging plants and buildings in W. Australia,

366.

Mastrus carpocapsae, utilisation of, against Cydia pomonella in New Zealand, 323.

Matai (see Podocarpus spicatus).

materna, Othreis.

Matricaria inodora, as a source of insecticides, 381.

Matsucoccus acalyptus, on Pinus edulis in U.S.A., 57.

Matsucoccus feytaudi, sp. n., bionomics of, on Pinus pinaster in France, 329. Matthiola incana, thrips on, in green-

houses in Maryland, 240.

Maurandia antirrhiniflora, tests of toxicity of, to insects, 304.

mauritanicus, Tenebroides.

Mauritius, Gonipterus scutellatus in, 101; review of economic entomology in, 165.

Mayetiola destructor, on wheat in U.S.A., 4, 28, 136, 157, 177, 181, 375; in U.S.S.R., 157; bionomics of, 136, 158; on barley and grasses, 157; varieties and time of sowing of wheat in relation to, 4, 28, 158, 177, 375; other cultural measures against, 28, 157, 158.

Mealybug, Pineapple (see *Pseudococcus* brevipes).

Meat, Dried, methyl bromide against pests in, in Morocco, 66.

Mecistorhinus, habits of, on cacao in Trinidad, 57.

medicaginis, Jaapiella.

Medicago arborea, 11.

Medicago hispida, Hypera punctata on, in California, 322.

Medicago sativa (see Lucerne).

Medlar, Magdalis barbicornis on, Britain, 272.

Megacoelum apicale, on cotton Uganda, **13.**

proserpina, Megamelus Cyrtorhinus fulvus predacious on, on Colocasia in Fiji, 101.

Megaselia, parasite of Schoenobius bipunctifer in China, 152.

Megastigmus, key to Finnish species of,

Megastigmus grönblomi, sp. n., in seeds of Abies sibirica in Finland, 271.

Megastigmus nigrovariegatus, bionomics of, on rose in Illinois, 292, 293.

Meigenia mutabilis, failure of, to parasitise Leptinotarsa decemlineata France, 363.

Melaleuca leucadendron, resistance of timber of, to termites in W. Australia, 366.

Melanagromyza (see Agromyza).

melanagromyzae, Loxotropa.

Melanaspis obscura, eradication of, on nuts and stone fruits in California, 108, 373.

Melanaspis tenax, sp. n., intercepted in California on orchids from Colombia and Guatemala, 112.

Melanauster chinensis, bionomics and control of, on Citrus in China, 270.

melanderi, Nomia.

Melanocallis caryaefoliae, on pecan in Arizona, 27.

melanocephala, Diabrotica.

Melanoplus (in U.S.A.), plants damaged by, 111, 180, 232; Arthropods destroying eggs of, 180, 232; baits for, 111, 180, 181, 339, 340; effect of weather, etc., on feeding by, 340; cultural measures against, 231-233; sprays killing weeds and, in peach orchards, 141.

Melanoplus bivittatus, in U.S.A., 180, 232, 339; bionomics of, 232.

Melanoplus differentialis, in U.S.A., 111, 180, 339, 340; bionomics of 232, 340.

Melanoplus femur-rubrum, in U.S.A., 111, 180, 232, 301, 302, 312; bionomics of, 232; sabadilla and its alkaloids in dusts against, 301, 302, 312.

Melanoplus foedus subsp. iselyi, Oklahoma, 339.

Melanoplus marginatus, in California,

Melanoplus mexicanus, outbreak of, in Br. Columbia, 227; in U.S.A., 180, 232, 339; bionomics of, 227, 232.

Melanoplus packardi, in U.S.A., 180, 339. melanum, Cryptochetum (Tritolestes).

Melcha nursei (see Goryphus).

Melia azadirachta, in smoke mixture against moths, 282; Geometrid on, in India, 285; new Coccid on, in Iraq, 16.

Melia azedarach, extract of, as repellent

against locusts, 11.

Melia dubia, Aonidiella aurantii on, in

W. Australia, 364.

Meligethes aeneus, dusts against, in Denmark, 7; cultural measures reducing damage to rape by, in Germany,

Melilotus, effect of sowing date on damage to, by Sitona cylindricollis in N. Dakota, 29, 30.

Melipotis acontioides (see Lyncestis).

satyriniformis, insecticides Melittia against, on squash in U.S.A., 52, 337. mellonella, Galleria.

Melolontha, survival of, in crop rotations

in U.S.S.R., 157.

Melon, Gryllotalpa on, in Egypt, 20; pests and disease of, in U.S.A., 148, 355; insecticides causing injury to, 149. (See Watermelon.)

Melon Fly (see Dacus cucurbitae).

Melophagus ovinus, permeability of cuticle of, by insecticide solutions, 87. mendax, Agrilus.

Mentha rotundifolia, Empoasca lybica overwintering on, in Spain, 267.

Mercuric Chloride, 319; tests of, against Hylemyia antiqua and Acrolepia assectella, 68; unsatisfactory for sapstream treatment of trees, 56.

Mercurous Chloride (Calomel), uses of, against *Hylemyia* spp., **68, 73, 92,** 319; dusting with, unsatisfactory against Acrolepia assectella, 68; used in determination of lime in calcium arsenate, 388.

Mercury, vapour of, against pests of stored grain and pulses, 45, 296, 326; treatment of wood with salts of, against termites, 172.

Merops orientalis, destroying bees in Punjab, 297.

mesembryanthemi, Pulvinaria.

Mesembryanthemum, Pulvinaria mesembryanthemi on, in France, 349.

Mesnilia legeri, gen. et sp. n., Clysiana ambiguella in France, 32.

Mesohomotoma tessmanni, not transmitting swollen-shoot disease of cacao in Gold Coast, 102.

Mesoleius tenthredinis, parasite of Pristiphora erichsoni in Quebec, 173.

messoria, Euxoa.

Metagonistylum minense (parasite of Diatraea saccharalis), apparent failure to establish, in Barbados, 159; imported into Porto Rico from Brazil, 219.

Metaldehyde, ineffective in baits for mole-crickets, 192.

Metamasius sericeus (in banana) in Fr. Cameroons, 361; in Cuba, 172.

Metaphycus helvolus (parasite of Saissetia oleae), in California, 206; suggested introduction of, into Chile, 206.

Metaphycus lounsburyi (parasite Saissetia oleae), in California, 206; finding and utilisation of, in Chile, 206.

Metaplectrus solitarius, sp. n., parasite of Limacodids in Ceylon, 47.

Metaplectrus thoseae, parasite of Thosea cervina in Ceylon, 47.

Metarbelids, immature stages of Indian,

Metarrhizium anisopliae, value against Cleonus punctiventris on beet in Russia, 131.

Metasyrphus, predacious on Cinara in N. Carolina, 311.

Metatetranychus ulmi. auct. (see Paratetranychus pilosus)

metathoracica, Amauromorpha.

Meteorus laphygmae (parasite of Lepidoptera), establishment of, in Hawaii, 3, 171, 274; in Texas, 3; bionomics of, 3.

Methallyl Chloride, as a soil fumigant against Dermolepida albohirtum, 1; against pests in stored cereals, 341, 342; and carbon tetrachloride, 342.

Methyl Benzoate, not accelerating penetration of diphenylamine through

insect cuticle, 87.

Methyl Bromide, fumigation of plants and planting material with, 26, 65, 76, 95, 109, 110, 292, 374; effects of, on plants, 65, 110, 374; unsatisfactory against mushroom mites, 77: against pests in fruit and vegetables, 107, 109, 110, 189, 190, 243, 363, 364, 374; effects of, on fruits and vegetables, 107, 109, 189, 190, 363, 364; against pests in other stored products, 65, 66, 180, 342, 343, 354; retarding germination of certain seeds, 66; improving quality of Christmas trees, 316; residual effect of adsorbed, 343; and carbon dioxide, 363, 364; and carbon tetrachloride, 342; laboratory dispenser for, 320; determination of, in air, 128; for producing aerosols, 320.

Methyl Chloride, for producing aerosols, 241, 242, 243, 338; risks of, 241.

Methyl Salicylate, compost treatment with, against Tyrophagus lintneri, 78.

4 - Methylcyclohexanol, accelerating penetration of insecticide through insect cuticle, 87.

Methylnaphthalenes, tests of, against pests of stored tobacco, 339; Velsicol AR 60 (q.v.) containing, 143, 319.

meticulosalis, Terastia.

Mexican Bean Beetle (see Epilachna varivestis)

Fruitfly (see Anastrepha Mexican ludens).

mexicanus, Melanoplus.

Mexico, fruit pests in, 110, 137, 179, 343, 359, 376; Fundella argentina in, 104; pests of sugar-cane and cereals in, 179, 375; introduction of beneficial insects into, 179; Coccobacillus acridiorum imported into from, 10.

micans, Lachnosterna.

Mice. destroying Diabrotica cephala, 148.

Michigan, Cydia pomonella in, 195; new Agromyzid on onion in, 57.

Microbracon (see Bracon).

Microcryptus basizonus, parasite of Neodiprion sertifer in Finland, 9, 39.

Microdiprion, systematic position of,

Microgaster, of Ethiopian Region, 81. Microgaster plecopterae, parasite Plecoptera reflexa in India, 286.

Microgaster sylleptae, sp. n., parasite of Sylepta sabinusalis in Belgian Congo,

Micromyzus (see Idiopterus).

Microphanurus basalis, introduced parasite of Nezara viridula in Queensland,

Microphanurus mopsus, parasite Antestia in Uganda, 278.

Microphanurus seychellensis, parasite of Antestia in Uganda, 278.

Microphanurus suranus, parasite Antestia in Uganda, 278.

Microplectron, Dahlb. (see Dahlbominus).

Midway Island, pests intercepted in aircraft at, 2.

migratoria, Locusta.

migratorioides, Locusta migratoria.

migratorius, Dysdercus.

miliaris, Asterolecanium; Aularches.

militaris, Stinoplus.

Milk, and bentonite, as adhesive in leadarsenate spray, 220, 221, 308; (dried), Hylemyia coarctata unable to survive on, 369.

Milky Disease, bacillus causing, in Odontria zealandica in New Zealand,

failure to infect American Lamellicorns with New Zealand bacillus of, 210. (See Bacillus popilliae.)

minense, Metagonistylum.

Mineola vaccinii, on cranberry in U.S.A., 51, 75; sprays against, 75.

minerva, Draeculacephala.

minimus, Dicyphus.

Minnesota, Noctuids in, 262.

minor, Myelophilus.

minuta, Hoplocampa.

minutum, Írichogramma. minutus, Dinoderus; Laemophloeus. Mississippi, cotton pests in, 178; pests

of Solidago leavenworthii in, 309. Missouri, pests of cereals in, 157, 177, 323; new Agromyzid on onion in,

57; Reticulitermes flavipes in, 49. Mites, methods of estimating populations of, in stored products, 98; attacking insects, 211, 232, 251, 260, 288, 354; attacking other mites, 92, 191; classification and new species

of, 104, 110, 111, 161; enemies of, 91, 92, 191.

mitis, Zenillia (Exorista). moderator, L., Nemeritis (Phaedroctonus, Ichneumon).

moderator, F., Porizon (Ophion).

modesta, Ealincola (Lycidocoris, lioforma).

modoc, Camponotus herculeanus.

Molasses, in baits, 2, 8, 16, 23, 294, 322, in bait-spray, 331; 372, 387; other sprays, 164, 266; feeding Dacus cucurbitae on, 308. Mole-crickets (see Gryllotalpa and Scap-

teriscus).

molesta, Cydia (Grapholitha, Laspeyresia).

molitor, Tenebrio. mollis, Ernobius.

Monellia (see Myzocallis).

Monilia fructigena (see Sclerotinia).

scutellatus, Monochamus measures against, in conifer logs in Quebec, 173. Monoctenus, systematic position of, 14. Monodontomerus dentipes, parasite of Gilpinia in Finland, 39.

Monodora tenuifolia, Achaea lienardi on,

in Nigeria, 14.

Monolepta russica, on strawberry in Uzbekistan, 155.

Monomorium pharaonis, bait for, buildings in Czechoslovakia, 328.

Monophlebus tamarindus (see Drosichiella).

Monoptilota pergratialis (nubilella), bionomics and control of, on lima beans in Virginia, **317, 318.**

Montana, grasshoppers in, 177, 180; miscellaneous pests in, 146, 181.

montanus, Amblyteles. monticola, Chaetogaedia. monticolae, Dendroctonus.

mopsus, Microphanurus.

Mordella fasciata, bionomics of, on plum in Britain, **272.**

Mordwilkoja vesicalis, on Populus suaveolens in Central Asia, 156.

mori, Bombyx.

Moringa oleifera (pterygosperma), pests of, in Travancore, 288.

mormoides, Achaea.

Mormon Cricket (see Anabrus simplex). Morocco, Sesamia vuteria in, miscellaneous pests in, 65, 364.

morosa, Incurvaria (see I. quadri-

opunctella).

morrisoni, Parlatoria.

morstatti, Xyleborus.

Mortegg, 187.

natural

moruliferus, Rhizoecus.

moschata, Aromia.

mosellana, Sitodiplosis (Clinodiplosis). Mountain Ash (see Sorbus aucuparia). Mountain Fir (see Abies lasiocarpa).

Mountain Pine Beetle (see Dendroctonus monticolae).

Mountain Poplar (see Populus suaveolens).

Mulberry, Coccids on: in W. Australia, 364; in France, 362; in Iraq, 16.

Mullein (see Verbascum). multistriatus, Scolytus.

Murgantia histrionica (Harlequin Cabbage Bug), tests of insecticides against, in U.S.A., 144, 244, 355.

Musa (see Banana).

Musa martinii, Cosmopolites sordidus on, in Cuba, **172.**

Musa textilis, Cosmopolites sordidus on, in Cuba, 172.

Musanga smithii, Bixadus sierricola on, in Fr. Equatorial Africa, 362.

Musca autumnalis (corvina), toxicity of powder of Matricaria inodora to, 381.

Musca domestica, tests of insecticides on, 134, 217, 301; pollinating onions, protection of, from DDT aerosols, 240.

Muscari (Grape Hyacinth), tolerant to methyl bromide, 110.

Muscari comosum (Cipollini), methyl bromide against Exosoma lusitanica in, in Morocco, 65.

musculosa, Oria.

museorum. Anthrenus.

Mushroom Mite (see Tyrophagus lintneri).

Mushrooms, Tyrophagus lintneri on, in Canada and U.S.A., 76.

musicus, Gastrimargus.

Mustard, effect of, on infestation of wheat by Hylemyia coarctata Britain, 216, 217; Phyllotreta on, in Denmark, 7.

Mustard, Indian (see Brassica juncea). mutabilis, Meigenia.

Myelois, methyl bromide against, in dried figs in Morocco, 65.

Myelois ceratoniae, Apanteles uroxys associated with, on Coelocaryon in Belgian Congo, 81.

Myelois venipars, in mummified fruits in California, 375.

Myelophilus spp., measures against, on Pinus pinaster in France, 371.

Myrciaria jaboticaba, Conotrachelus myrciariae on, in Brazil, 6.

myrciariae, Conotrachelus.

Myriangium, infesting Clavaspis herculeana in Nyasaland, 63.

Myrobalans (see Terminalia).

Mysore, miscellaneous pests in, 186, 207, 287; biological control of sugar-cane borers in, 207.

Mytilococcus (see Lepidosaphes).

Mytilococcus conchiformis, auct. (see Lepidosaphes ficus).

Myzocallis costalis, on pecan in Arizona, 27.

Myzus ajugae, on Ajuga reptans in Holland, 367.

Myzus caryophyllacearum, sp. n., on Dianthus in Holland, 367.

Myzus certus, on Viola arvensis in Holland, 367.

Myzus festucae (see Macrosiphum).

Myzus ornatus, in Britain, 70; in Hawaii, 274; relation of, to virus diseases of lettuce and crucifers, 70, 71; on other plants, 274.

Myzus persicae, in Argentina, 299; in Australia, 231, 385, 386; in Britain, 70, 113, 114; in Holland, 99, 367; in Nyasaland, 62, 205; in U.S.A., 72, 182, 224, 226, 239, 240, 243; transmitting virus yellows of beet and mangel, 113, 114; on crucifers, 113, 226, 231, 240, 367, 385, 386; relation of, to virus diseases of crucifers, 71, 182; on potato, 72, 224, 239, 240, 243; transmitting potato leaf-roll, 226; on Prunus spp., 113, 226, 367, 385; relation of, to virus diseases of tobacco, 62, 70, 206; on other plants, 62, 206, 240, 367; relation of, to virus diseases of other plants, 70, 99; fungus infesting, 299; bionomics and forms of, 113, 206, 367; measures and experiments against, 72, 113, 114, 206, 224, 226, 231, 239, 240, 243, 385, 386; zinc-arsenite increasing infestation by, 224.

Myzus pseudosolani (see Macrosiphum solani).

Myzus ribis (see Capitophorus).

N.

Nabis, killed by DDT in Nebraska, 224. Nabis lativentris (see Himacerus).

naevana, Rhopobota.

Nanophyes croceus, sp. n., on banana in

Senegal, 14.

Naphtha, as solvent for DDT, 45, 382. Naphthalene, ineffective in repellent spray against Anthonomus pomorum, 273; against Bruchids in stored beans, 126; greenhouse fumigation with, against mites, 161; soil treatments with, against pests, 17, 74, 192, 319; effect of, on germination of seeds, 74, 192; as solvent for DDT, 157.

β-Naphthol, spraying with, ineffective against Anthonomus pomorum, 273.

Napomyza (see Phytomyza).

narangae, Zacharops.

nararia, Natada.

Narcissus, Aphids and virus disease of, in Holland, 99.

Narosa conspersa, natural enemies of, in Ceylon, 47, 325.

Nasturtium (see Tropaeolum).

Natada nararia, parasite of, on tea in Ceylon, 324.

natadae, Platyplectrus.

natalensis, Crypticerya; Elasmus masii. nazarena, Anomala (Phyllopertha).

Nebraska, cereal pests in, 28, 192, 312, 322; potato pests in, 28, 29, 223, 224; Anasa tristis on squash in, 312, 313; Cydia molesta in, 373.

nebulosa, Cassida.

Necremnus duplicatus, parasite of Ceuthorrhynchus assimilis in Washington, 72, 76.

Necrobia ruficollis, in carcasses in Fiji and New Zealand, 101; resistance of, to cold, 101.

Necrobia rufipes, fumigation against, in dried meat in Morocco, 66; in carcasses in Fiji and New Zealand, 101; resistance of, to cold, 101.

Nectarine, Aonidiella aurantii on, in W. Australia, 364; Anuraphis schwartzi on, in Britain, 272; Cydia molesta on, in California, 110; Rhynchites bacchus on, in France, 370; as winter food-plant of Myzus persicae, 367.

Nectria cinnabarina, 41.

Necydalis major, parasites of, on Populus tremula in Finland, 41.

Nematodes, 135, 299; infesting insects, 160, 164, 177, 210, 343, 344.

Nematus erichsoni (see Pristiphora).

Nematus ribesii (on gooseberry), in Britain, 273; in Denmark, 116; insecticides against, in Holland, 59.

Nemeritis, parasite of Rhyacionia buoliana in Argentina, 372.

Nemeritis moderator, L. (Phaedroctonus cremastoides), parasite of Cydia strobilella in Finland, 40; systematic position and synonymy of, 252.

Nemeritis palmaris, establishment of, against Coleoneura trichogramma in

Fiji, 366.

nemoralis, Neurotoma. nemorata, Phyllotoma.

Nemorilla floralis, parasite of Lamprosema indicata in Porto Rico, 219.

nemorum, Phyllotreta.

nenuphar, Conotrachelus.

Neoaplectana, Diplogaster displacing, in cultures and insects, 344.

Neoaplectana glaseri, experiments with insects and, imported into Zealand, 210.

Neocerata rhodophaga, bionomics and control of, on greenhouse roses in Ohio, 78.

Neodiprion, characters and systematic position of, 14.

Neodiprion lecontei, on pine in Quebec, 173.

eodiprion sertifer (on pines), in Finland, 9, 38, 39; in Sweden, 130, Neodiprion 131; on spruce, 131; bionomics of, 38, 130; natural enemies of, 9, 39, 130.

Neodiprion swainei, on Pinus banksiana in Quebec, 173.

Neodiprion tsugae, bionomics and control of, on Tsuga spp. in Br. Columbia, 377, 378, 379.

Neopales pavida, parasite of Taeniocampa gracilis in Finland, 42.

Neoplectrus maculatus, bionomics of, in Ceylon and Java, 324, 325.

Neopristomerus appalachianus, parasite of Laphygma exigua on maize in Texas, 3.

Nephantis serinopa, use of Trichospilus pupivora against, on coconut Ceylon, 188.

Nephotettix apicalis, intercepted in transpacific aircraft, 2.

nerii, Deilephila.

Nerium oleander, new Coccid on, in Iraq, 16.

Nesodiprion, systematic position of, 14. Netherlands Indies, parasite of Artona catoxantha in, 325.

Neurotoma nemoralis, on stone fruits in France, 160.

neustria, Malacosoma.

Nevada, Anabrus simplex in, 177.

New Brunswick, miscellaneous insects in, 263, 319.

New Caledonia, thrips on sugar-cane and onion in, 275.

New Hampshire, Tineola biselliella in-

451

New Jersey, cereal pests in, 137, 147, 177, 203; Cydia spp. on fruit trees in, 139, 203, 204; new Agromyzid on lima beans in, 57; miscellaneous pests in, 142, 194, 204; beneficial insects and biological control in, 139, 142, 177, 194, 204.

New Mexico, Cydia pomonella in, 308;

pests of Pinus edulis in, 57.

New South Wales, outbreak areas of Austroicetes cruciata in, 123, 124; fruit pests in, 71, 72, 207, 208, 289, 365, 382, 383, 384, 385; pests of vegetables in, 208, 382, 383, 384, 385, 386; pests of stored products in, 21, 383, 385; miscellaneous pests in, 208, 382, 385; beneficial insects in, 82, 208.

New York, orchard pests in, 195, 246, 248, 249; potato pests in, 238, 239; miscellaneous pests in, 105, 137, 176, 194; beneficial insects in, 194, 263.

New Zealand, Coccids on Citrus in, 21; Coptotermes spp. in, 169; Odontria zealandica and Oxycanus spp. in, 20, 58, 208, 210; miscellaneous pests in, 58, 101, 209, 210, 323; parasites, diseases and biological control of insects in, 210, 211, 323; insects introduced against noxious plants in, 58, 323.

Newfoundland, forest pests in, 377.

newsteadi, Lepidosaphes.

Nezara viridula (in Australia), biological control of, on beans, 126; on sorghum, measures and experiments against, 126, 383, 386.

Nicandra, Nysius vinitor on, in New South Wales, 382.

Nicotiana spp., Agalliana ensigera not breeding on, in Argentina, 380; Gnorimoschema heliopa on, in Nyasaland, 62; experiments with turnip viruses and, 182. (See Tobacco.)

Nicotine, in repellent mixture against Anastrepha ludens, 376; Aphids, 53, 72, 78, 125, 145, 169, 177, 178, 184, 236, 241, 242, 244, 309, 356, 385; effect of, on natural enemies of Chromaphis juglandicola, 145; against Capsids, 54, 243, 276; against Coccids, 79, 184; against other Hemiptera, 89, 243, 267; against Coleoptera, 72, 85, 243; against Cydia pomonella, 8, 52, 175, 195, 196, 197, 198, 202, 203; against other Lepidoptera, 52, 53, 70, 92, 126, 137, 185, 199, 216, 236, 249, 339, 385; against Hoplocampa spp., 6, 37; against mites, 77, 175; against thrips, 67, 78, 118, 125; toxicity of, to bees, 55; site of action

of, in Periplaneta americana, 5; effect of detergents on penetration of insect 145, 169, 177, 178, 184, 185, 195, 197, 236, 267, 276, 309, 356, 385; effects of carriers on toxicity of, 236; green-78, house fumigation with, apparatus for field fumigation with, 370; spraying with, 6, 37, 52, 53, 54, 70, 78, 89, 92, 100, 137, 175, 184, 196, 197, 198, 199, 202, 203, 216, 249, 267, 376: fixed compounds of, 52, 53, 54, 77, 137, 175, 178, 195, 197, 198, 199, 202, 203, 236, 249, 356; formulae containing, 6, 37, 53, 70, 77, 92, 100, 137, 169, 184, 195, 197, 198, 199, 202, 203, 216, 236, 242, 243, 249, 276, 356, 376, 385; and Bordeaux mixture, 53, 54, 249, 267; and calcium arsenate. 72, 178, 356; and cryolite, 52; decomposition of DDT by, 157; and dinitro compound, 100; and lead dinitro compound, 100; arsenate, 195, 197, 202, 203, 385; and oil emulsions, 175, 184, 197, 198, 199, 202, 203, 249, 376; and pyrethrum, 54; and rotenone insecticides, 137, 177; wetting agent for, 6; report on methods of analysis of, 388.

Nicotine Sulphate; against Aphids, 125, 177, 178, 231, 311, 319, 356; against Coccids, 200: against other Hemiptera, 54, 126, 247, 248, 384; against Coleoptera, 85, 177, 237; against Cydia pomonella, 4, 106, 175, 197, 198; against other Lepidoptera, 52, 112, 126, 199, 209, 231, 249, 372; against Diptera, 18, 125, 384, 385; against mites, 54, 77; against product of the collection of the collec sawflies, 229, 365; against thrips, 53, 118, 125, 218, 240; in bait-spray, 385; dusting with, 177, 178, 231, 311, 356, 384; grain bins treated with, 177; formulae containing, 4, 18, 52, 54, 77, 106, 112, 118, 125, 197, 199, 200, 209, 218, 231, 247, 248, 249, 365, 372, 385; and bentonite, 197, 199; and Black Leaf 155, 197; Bordeaux mixture, 52, 247, and calcium arsenate, 311, 356; and Dreft, 199; and lead arsenate, 4, 106, 197, 198, 200, 247; and lime, 198, 199; and magnesite, 231; and oil emulsions, 18, 51, 52, 106, 125, 197, 199, 200, 247, 248, 249, 372, 384; and rotenone, 356; and soap, 18, 53, 54, 118, 125, 199, 237, 365.

niger, Thysanus.

Nigeria, cacao pests in, 46, 296; new Curculionid on Cinchona in, 206; pests and disease of cotton in, 96, 169;

nigricana, Cydia. nigricans, Euxoa (Agrotis). nigricarpus, Euphorus. nigricornis, Taeniothrips. nigripes, Phyllotreta; Pleurotropis. nigritula, Baris. nigritulus, Elachertus (Elachistus), nigriventris, Proctacanthus. nigroareolata, Icerya. nigrofasciatum, Eulecanium (Lecanium). nigrofasciatus, Dysdercus. nigrolimbatus, Podisus (Apateticus). nigronervosa, Pentalonia. nigropilosus, Thrips, nigrorepletus, Hieroglyphus. nigrovariegatus, Megastigmus. Nilotaspis halli, campaign against, in California, 108, 373; in Iraq, 16; on peach, 16, 373. nitens, Anaphoidea. nitida, Erynnia. nitidalis, Diaphania. nitidella, Argyresthia. nitidipennis, Xyloperthodes. nitidula, Cladis. nitidulus, Anastoechus. nitidus, Hadrobregmus (Anobium): Phymatodes. Nitocris princeps (see Dirphya). Nitriles, 256, 339. Nitrobenzene, as solvent for DDT, 157. Nitromethane, apparatus for removing, from pyrethrum concentrates, 360. ω-Nitrostyrene Dibromide, effect of solvents on penetration of insect cuticle by, 87. nivella, Scirpophaga. njalensis, Pseudococcus. noctilio, Sirex. Nomadacris septemfasciata, in Nyasaland, **63.** Nomia melanderi, pollinating lucerne in Utah, 260. Noorda blitealis, on Moringa oleifera in Travancore, 288. Norway, Aphids on fruit trees, etc., in. 7. notabilis, Aesiotes. notatus, Pissodes. Notolophus antiquus (see Orgyia). Nova Scotia, forest pests and their biological control in, 129, 130: Popillia japonica in, 130; potato aphids in, 319. nubilalis, Pyrausta. nubilella, Monoptilota (see M. pergratialis). numitor, Telenomus. nursei, Goryphus (Melcha).

Nyasaland, cotton pests in, 62, 168, 332;

miscellaneous insects in, 62, 63; pests

fruit-piercing moths in,

beneficial insects in, 296.

14,

15:

of tea in, 58, 62; pests and diseases of tobacco in, 62, 206, 207; restrictions on importation of dahlia into,

Nygmia phaeorrhoea, on fruit trees, etc., in Denmark, 116; Batrachedra

ledereriella in nests of, 361.

Nysius ericae, bionomics and control of, on Taraxacum spp. in Russia, 35, 36. Nysius vinitor, food-plants Australia, 382, 384, 388; of, in dusts against, 382, 384.

Oak, Alsophila pometaria on, in Canada, 129; Lachnosterna on, in Connecticut,

Oats, pests of, in Britain, 330, 381; Sesamia vuteria on, in Morocco, 257; Nysius ericae on, in Russia, 36; pests of, in U.S.A., 250, 375; varieties and time of sowing of, in relation to pests, 376, 381; resistance of, to Hylemyia coarctata, 216, 368.

obesus, Itoplectis. oblineatus, Lygus.

obliqua, Diacrisia.

obliquus, Listroderes.

oblongus, Phyllobius. obovatus, Brevipalpus.

obscura, Cytherea; Melanaspis.

obscurata, Brachymeria.

obscurus, Agriotes; Anaphothri, Athous; Rhabdoscelus; Tenebrio. Anaphothrips; obsoleta, Heliothis (Chloridea) (see H. armigera).

obsoletana, Tortrix (Archips). obsoletus, Bruchus (Acanthoscelides);

Carpophilus.

obtectus, Bruchus. ocellana, Spilonota.

ochrifascia, Thamnonoma. Ochropyga lepesmei, sp. n., on Coffea robusta in Middle Congo, 362.

Odontoglossum (see Orchids).

Odontria, disease of, in New Zealand, 210; experiments with Neoaplectana glaseri and, 210.

Odontria zealandica (in New Zealand), bionomics and control of, 20; native and imported bacilli infecting, 210; experiments with Neoaplectana glaseri and, 210; apparatus for collecting emerging adults of, 58.

oeceticola, Tetrastichus.

Oeceticus (see Oeketicus).

Oedaleus decorus, in Spain, 268.

kirbyi, parasites of, in Oeketicus Argentina, 300.

Oeme rigida (linearis), on Juniperus bermudiana in Bermuda, 191.

Ohio, Pyrausta nubilalis on maize in, 136, 222; miscellaneous pests in, 43, 78, 178; orchard pests in, 128, 195, 201, 202.

Oil Emulsions, against Aphids, 125, 191; against Coccids, 21, 33, 51, 73, 79, 80, 108, 127, 184, 191, 200, 268, 289, 290, 291, 294, 364, 373, 384; against other Hemiptera, 89, 191; against Coleoptera, 93, 101, 204, 234, 273, 321; against Cydia pomonella, 50, 73, 106, 175, 195, 197, 198, 199, 202, 203, 309; against other Lepidoptera, 52, 75, 125, 177, 199, 200, 372; treatments of maize ears with, against Heliothis armigera, 147, 177; against Diptera, 18, 75, 125, 191, 221, 376, 384; ineffective against Hoplocampa flava, 37; against mites, 19, 50, 73, 74, 91, 107, 127, 161, 175, 319, 371, 384; against Orthoptera, 191, 208; against thrips, 125; types of oils for, 18, 50, 51, 73, 74, 75, 79, 80, 89, 91, 101, 106, 141, 175, 197, 198, 199, 202, 204, 229, 238, 249, 294, 309, 319, 321, 373, 385; emulsifiers for, other than soap, 50, 175, 238, 294; formulae for, 18, 21, 50, 51, 52, 73, 74, 75, 78, 79, 80, 89, 101, 106, 125, 141, 143, 175, 184, 195, 197, 198, 199, 200, 202, 208, 221, 229, 234, 238, 247, 249, 273, 275, 289, 290, 291, 294, 309, 319, 321, 372, 373, 383, 384, **385**; formulae containing, **18**, **52**, **73**, 74, 75, 79, 80, 141, 198, 199, 204, 221, 294, 385; and barium fluosilicate, 321; and bentonite, 51, 199, 234; and Bordeaux mixture, 21, 52, 247, 294, 319; and calcium arsenate, 50. 51; and cryolite, 50, 74, 175, 200, 208, 321; and DDT, 75, 79, 106, 143, 185, 195, 238, 247, 275, 383, 384, 385; and dinitro compounds, 73, 74, 93, 141; and lead arsenate, 50, 51, 73, 74, 75, 101, 106, 191, 199, 200, 202, 203, 204, 220, 221, 234, 247, 249, 309; and Lethane B-72, 247; and limesulphur, 73; and nicotine, 18, 52, 106, 125, 175, 184, 197, 198, 199, 200, 202, 203, 247, 249, 372, 384; and pentachlorphenol, 141; and phenothiazine, and pyrethrum, 177; rotenone compounds, 80, 229; zinc sulphate, 50, 199, 202, 203, 309; "inverted" mixtures of insecticides and, 50, 51; droplet size of, 268; and injury to plants, 50, 51, 75, 89, 91, 191, 197, 289, 291.

Oil Palm (see Elaeis guineënsis).
Oils, as repellent for Chlorops, 286; soil treatment with, against Epitrix hirtipennis, 316; spraying with, against locusts, 16, 227; spraying with, against Neocerata rhodophaga, 78;

orthodichlorbenzene in, against Coleoptera in elm logs, 54; atomised or vaporised sprays of insecticides in, 62, 302, 339, 346, 374, 377; in DDT fog mixture, 262; and supplementary materials, treatment of maize ears with, against Heliothis armigera, 147; testing of, on larvae of Gnorimoschema operculella, 134; use and functions of, in aerosols, 239, 240, 241, 242, 338; in baits, 25, 339, 341; dusts impregnated with, 50, 195, 197, 248, 338; analysis of DDT and pyrethrins in, 256.

Oils, Essential (see Sassafras Oil).

Oils, Vegetable, emulsions of, against Aphid and mite on cotton, 293; and insecticides, treatments of maize ears with, against Heliothis armigera, 147; as adhesive for sprays, 72, 196, (See Cashew-nut Shell, Sesame and Tea-seed Oils.)

Oklahoma, Listroderes obliquus in, 310; Melanoplus spp. in, 339, 340; miscellaneous pests in, 180, 373, 375.

Okra (see Hibiscus esculentus).

oleae, Dacus; Eriophyes; Filippia (Philippia); Parlatoria; Prociphilus; Saissetia.

oleellus, Prays.

Oleic Acid, and acrylonitrile, aerosol of,

oleivorus, Phyllocoptruta (Phyllocoptes). oleracea, Polia (Mamestra).

oleraceum, Eurydema.

olivacea, Caedicia. Olive, Flatid on, in Bermuda, 190; Coccids on, in California, 26, 375; Saissetia oleae on, in Chile, 206; pests of, in Greece, 326; Coccids on, in Iraq and Persia, 16; pests of, in Spain, 267, 268.

Olive Scale (see Parlatoria oleae).

olivina, Euphyllura.

Olla abdominalis, effects on, of nicotine dust against Chromaphis juglandicola in California, 145.

Omophlus proteus, survival of, in crop rotations in U.S.S.R., 157.

Omotrachelus bigranulatus, sp. n., on Cinchona in Nigeria, 206.

Oncidium (see Orchids).

Oncocoris, measures against, on beans in Queensland, 126.

Oncopeltus fasciatus, tests of insecticides on, 301, 302.

Oncopera fasciculata, bionomics of, on grasses in New Zealand, 324.

Onion, Acrolepia assectella on, in Britain, 69, 70; Euxoa spp. on, in Finland,38, 100; pests of, in Hawaii, 274, 277; Thrips tabaci on, in India, 118; T. tabaci on, in New Caledonia, 275: T. tabaci on, in New South Wales, 384; Acrolepia assectella on, in Switzerland, 214, 216; pests of, in U.S.A., 53, 57, 240, 243, 247; dinitro dust injuring, 53.

Onion Thrips (see Thrips tabaci).

onobrychis, Macrosiphum.

Ontario, forest pests in, 263, 320, 377; miscellaneous pests in, 319, 320; beneficial insects and biological control in, 263, 320.

Ovencyrtus kuvanae, parasite of Lyman-

tria dispar in U.S.A., 193.

Ovencyrtus prenidis, sp. n., parasite of Panoquina in Porto Rico, 64.

opaca, Silpha (Blitophaga). opalescens, Aegeria (Sanninoidea).

operculella, Gnorimoschema (Phthorimaea).

Operophtera brumata (on fruit trees), in Denmark, 8; in Holland, 90, 94; sprays against, 8, 90.

Ophiomyia (see Agromyza).

Ophion moderator, F. (see Porizon).

Opius, parasite of Pegomyia hyoscyami betae in Britain, 331, 332.

Opsimus quadrilineatus, on Sequoia sempervirens in California, 31.

Opuntia, Diaspis boisduvali on, in California, 145; biological control of, in Mauritius, 165; Ceratitis capitata not ovipositing on, in Palestine, 132.

Orange, Tenuipalpus pseudocuneatus causing "lepra explosiva" in, Argentina, 298; pests of, in Australia, 126, 289, 384; Flatid on, in Bermuda, 190; Saissetia oleae on, in Chile, 206; Icerya on, in Belgian Congo, 82; Lamellicorns on, in Costa Rica, 128; Anastrepha ludens on, in Mexico, 376; fruit-piercing moths attacking, in Nigeria, 14, 15; pests of, in U.S.A., 24, 25, 79, 80; effects of sprays on, 79, 80; (fruit), vapour-heat treatment of, against Anastrepha, 112; as food for adults of A. fraterculus, 138; Myelois venipars in mummified, in California, 375.

Orange, Mandarin or Tangerine, moths attacking fruit of, in Nigeria, 14; resistance of, to Aceria sheldoni in Queensland, 126.

Orange, Satsuma, Philaenus leucophthalmus on, in Hawaii, 274.

orbona, Agromyza.

Orchestes (see Rhynchaenus).

orchidii, Anaphothrips.

Orchidophilus peregrinator, dust against,

on orchid in Hawaii, 275.

Orchids, Orchidophilus peregrinator on, in Hawaii, 275; pests of, in U.S.A., 109, 314, 315, 374; new Coccid intercepted in California on, from

455

Colombia and Guatemala, 112; list of insects associated with, 277.

Oregon, Camponotus spp. in, 27; mites on filbert in, 111; orchard pests in, 74, 143; pests of small fruits in, 79, 160, 292; miscellaneous pests in, 185, 247, 250, 318.

Oreodoxa regia, new Coccid on, in Belgian Congo, 82; Rhabdoscelus obscurus on, in Queensland, 1.

Oreopsyche angustella, damaging tures in France, 349.

Organothrips bianchii, on Colocasia in

Hawaii. 276.

Orgyia antiqua, Pentatomid predacious on, in Chile, 352; on elm and maple in Nova Scotia, 130.

Orgyia antiqua badia, bionomics and control of, in Br. Columbia, 377, 378,

Oria musculosa, bionomics of, in Russia and Britain, 157, 252.

Oriental Fruit Moth (see Cydia molesta). orientalis, Anomala; Aonidiella; Blatta. Orissa, Apion corchori on jute in, 117; use of Trichogramma against sugarcane borers in, 207.

Orius, killed by DDT in Nebraska, 224. Orius australis, predacious on sorghum insects in Queensland, 388.

Orius majusculus, injury to maize by, in Holland, 95.

Ormenis infuscata, bionomics of, on Juniperus bermudiana in Bermuda, 190.

ornatulopsis, Phanerotoma.

ornatus, Myzus.

orophila, Helopeltis.

Orthezia insignis, bionomics and control of, in S. Rhodesia, 33.

Ortheziopa, Acropyga associated with, on cacao in Trinidad, 372.

Orthobracon (see Bracon).

Orthodichlorbenzene, against ponotus spp., 27; in spray against Coleoptera in elm logs, 54, 55; ineffective against Kalotermes piceatus, tests with, against molecrickets, 191; in DDT aerosols, 242; causing injury to plants, 191, 242; review of data on, as an insecticide, 32. orthogonia, Agrotis.

Orthol K Ready Mix Oil, 199.

useless against Ortho-nitrodiphenyl, potato pests, 224.

Chilo; orvzae, Calandra (Sitophilus); Latheticus; Lissorhoptrus; Pachydiplosis.

Oryzaephilus surinamensis, in dried bananas in France, 363; (in stored cereals), in Punjab, 296; in Victoria, 230: effects of competition with

other insects on, 97, 98; experiments against, 296, 342, 343,

Oscinella (Oscinosoma) frit, parasite of, in N. America, 295; effect of cultural measures on, on cereals in U.S.S.R. and Britain, 157, 381.

ostreaeformis, Quadraspidiotus.

Othreis, parasites and control of, attacking fruits in Ceylon, 188, 281, 282.

Othreis ancilla, in Ceylon, 281. Othreis aurantia, in Ceylon, 281.

Othreis fullonia (fullonica), attacking fruits in Nigeria and Ceylon, 14, 281, 282; bionomics and control of, 282.

Othreis hypermnestra, in Ceylon, 281. Othreis materna, in W. Australia, 365; in Ceylon, **281.**

Othreis salaminia (see Eumaenas).

Otiorrhynchus singularis, spray against, on black currant in Britain, 272.

Otiorrhynchus sulcatus (in California), on Begonia, 109; in lawns, 374; measures against, 109, 374.

ovatus, Platycampus.

ovijentatus, Anaphes (Anagrus).

ovinus, Melophagus.

Oxycanus cervinatus (in New Zealand), bionomics of, on grasses, 208, 209; experiments with Neoaplectana glaseri and, **210.**

Oxycarenus, on cotton in Sudan, 166. Oxycarenus luctuosus, in W. Australia,

oxycoccus, Aspidaspis.

P.

pabulinus, Lygus.

Pachydiplosis oryzae, bionomics of, on rice in China, 119.

Pachymerus cassiae, methyl bromide against, in stored ground-nuts in Morocco, 65.

Pachyneuron, host relations of, Belgian Congo, 82; parasite Dacus passiflorae in Fiji, 101.

Pachyrhizus, tests for rotenone in seeds of, 218.

Pachyrhizus erosus, insecticidal properties of, 218, 303.

Pachyrhizus tuberosus, dust of, as an insecticide, 303.

Pachyzancla (see Psara).

pacificus, Tetranychus.

packardi, Cydia (Grapholitha); Melanoplus.

Palarus, bionomics and control of, destroying bees in Punjab, 297.

Pale Western Cutworm (see Agrotis orthogonia).

Paleacrita vernata, on maple, etc., in Quebec, 173.

E2

Pales pavida (see Neopales).

Palestine, Anomala nazarena on wheat in, 17; Aphid on cabbage in, 17; Capnodis spp. in, 369; Ceratitis capitata in, 132, 133.

palliatus, Tanymecus. pallidum, Callidium.

pallidus, Tarsonemus.

pallipes, Priophorus; Pristiphora.

palmaris, Nemeritis. palmatus, Tenuipalpus.

Palmitic Acid, solid mixture of DDT and,

in traps for *Popillia japonica*, **245.**Palms, new Coccid on, in Belgian Congo, **82**; insects on, in California, 145, 375; termites attacking, in Danger Is., 211; Rhabdoscelus obscurus on, in Queensland, 1. (See Coconut, Date Palm and Elaeis.)

palpata, Eupithecia. paludosa, Tipula.

Panama Canal Zone, Eretmocerus serius imported into Mexico from, against Aleurocanthus woglumi, 179.

panicea, Sitodrepa.

Panicum purpurascens, thrips on, in Hawaii, 277.

Panicum repens, Pachydiplosis oryzae on, in China, 119.

panicus, Plesiothrips.

Paniscus, introduced into U.S.A. against Heliothis armigera, 179.

Panoquina, new parasite of, on sugarcane in Porto Rico, 64.

Pantilioforma modesta and thoracica (see Ealincola).

Pantomorus (Graphognathus) (Whitefringed Beetles), Nematodes infesting, in U.S.A., 177, 343.

Pantomorus godmani, on Citrus in New South Wales, 72, 383; sprays against,

Pantomorus peregrinus, Diplogaster aerivora infesting, in U.S.A., 343.

Papaver somniferum, new Cecidomyiid on, in Sweden, 59.

papaveris, Carpodiplosis.

Papaya, fumigation of fruit of, against introduction of fruit-flies into U.S.A. from Hawaii, 190.

Paper, bags of, to protect fruits from 282; penetration of, by moths, Calandra oryzae, 121; testing attractiveness of, to Lepismids and cockroaches, 127, 128.

papillosa, Parlaspis (Parlatoria).

2,2-bis (Parachlorphenyl)-1,1,1-trichlorethane (see DDT).

αα - bis (Parachlorphenyl - ββdichlorethylene, and analogues, experiments with, 254, 255.

Paradichlorbenzene, fumigation of beehives with, against Galleria mellonella, 297; against Bruchids in stored beans, 126; against borers in trees, 35, 202, 347, 348; soil treatment with, against pests, 17, 315; doubtful value of mixture containing, against mole-crickets, 191; unsatisfactory against Tyrophagus lintneri in mushroom houses, 77; ineffective in repellent spray against Anthonomus pomorum, 274; preparation of, in miscible oil, 202.

Paraffin Wax, uses of, as protection

against termites, 291. parallela, Tortrix (Cacoecia).

parancilla, Frontiniella.

Parasa lepida, Neoplectrus maculatus feeding on, in Ceylon, 325.

Parascalecide, 202.

Parasites, of insect pests, catalogue of,

Paratetranychus citri, on Citrus U.S.A., 25, 80; sulphur dust against, 25; DDT increasing infestation by,

Paratetranychus pilosus (on fruit trees), in Br. Columbia, 175, 319; in Denmark, 8, 116; in Holland, 91; in U.S.A., 50, 52, 106, 107, 197, 203, 221, 248; bionomics of, 91; measures and experiments against, 8, 50, 52, 91, 175, 203, 319; treatments sometimes increasing infestation by, 106, 107, 175,197, 221, 248.

Paratheresia (see Theresia).

Paratrioza cockerelli, on potato in U.S.A., 29, 181, 223, 224, 358; causing Psyllid yellows, 358; insecticides against, 29, 224, 358, 359.

pardalina, Locustana.

Parexorista caridei, fungus infesting, in Argentina, 300.

Paria canella (in U.S.A.), on rose, 78; bionomics of, on strawberry, 200; measures against, 78, 200.

pariana, Anthophila.

Paris Green, in baits, 2, 8, 23, 191, 340; in dusts, 84, 135, 164, 180, 207, 340, 354; carriers for, 164, 354; in sprays, 84, 100, 187, 334; and calcium arsenate, 340, 355; DDT compatible with, 157; and lead arsenate, 334; and sulphur, 135, 180; tomatos scorched by, 354; effect on plants of soil containing, 355; risks residues of, on wheat, etc., 340, 341. parkeri, Porizon.

Parlagena inops, gen. et sp. n., in China,

Parlaspis, gen. n., for Parlatoria papillosa, 205.

Parlatoreopsis chinensis (longispina), type of genus, 205.

Parlatoreopsis pyri, 205.

Parlatoria, revision of, 205.

Parlatoria blanchardi, on date palms in

Iraq, **16.**

Parlatoria camelliae, methyl bromide against, on Camellia in California, 109. Parlatoria chinensis (see Parlatoreopsis). Parlatoria mangiferae (see Genaparlatoria pseudaspidiotus).

Parlatoria morrisoni, sp. n., food-plants

of, in Iraq and Persia, 16.

Parlatoria oleae, measures and experiments against, in California, 26, 108, 373; in Greece, 326; in Persia, 16; on olive, 16, 26, 326; on other plants, 16, 26, 373; bionomics of, 26.

Parlatoria papillosa (see Parlaspis). Parlatoria pergandei, on Citrus in Spain, 269; P. phyllanthi distinct from, 205. Parlatoria phyllanthi, status of, 205.

Parlatoria proteus, spray against, on

orchids in Maryland, 315.

Parlatoria pyri (see Parlatoreopsis). Parlatoria ziziphus, on Citrus in Persia, 16.

parlatoriae, Syngenaspis.

parlatorioides, Pseudoparlatoria.

Parnara guttata, bionomics and control of, on rice in China, 151.

Parsley, Vasates eurynotus on, in California, 374: Philaenus thalmus on, in Hawaii, 274.

Parsnip, Chilosia illustrata on, in Finland, 38; Aphid on, in Queensland,

169.

Parthenium argentatum, pests of, in U.S.A., 26, 135, 149, 318, 346.

Partridges, question of poisoning of, by arsenicals, 83.

parvula, auct., Epitrix (see E. hirtipennis).

parvulus, Euplectrus.

parvus, Anicetus.

orbiculare, Pachydiplosis Paspalum

oryzae on, in China, 119.

Passiflora edulis (Passion Fruit), thrips on, in Hawaii, 277; Ceratitis capitata not ovipositing on, in Palestine, 132. passiflorae, Dacus (Chaetodacus).

Pavetta, Antestia spp. on, in Uganda,

pavida, Neopales (Pales).

pavonacella, Brenthia. Pea, little attacked by Listroderes obliquus in W. Australia, 23; resistance of, to wireworms in Britain, 330; Cydia nigricana on, in Canada, 73; experiments with Bruchus obtectus and, in Chile, 205; Polia oleracea on, in Finland, 100; Thrips tabaci on, in Hawaii, 276; Cydia on, in Holland, 90; Bruchus pisorum on, in U.S.S.R., 157; pests of, in U.S.A., 57, 73, 76, 133, 177, 181, 242, 243;

aerosols injurious to, 242: Bruchus pisorum in stored, in Morocco, 65.

Pea (see Macrosiphum ono-Aphis brychis).

Pea Moth (see Cydia nigricana).

Peach, Aphids on, in Britain, 113, 272; Aegeriid in, in Br. Columbia, 23; pests of, in France, 160, 229, 370; Coccid on, in Iraq, 16; pests of, in Kashmir, 293; Anastrepha frater-culus on, in Mexico, 137, 138; pests of, in New South Wales, 208, 385; Capnodis spp. on, in Tadzhikistan, 253, 254; pests of, in U.S.A., 108, 109, 110, 139, 140, 141, 142, 201, 202, 234, 237, 245, 321, 335, 336, 347, 354, 373, 374, 375; as winter food-plant of Myzus persicae, 113, 367; insecticide treatments and injury to, 52, 108, 141, 230, 321, 322; fumigation of packed fruit of, against Cydia molesta, 110.

Peach Aphis, Green (see Myzus per-

sicae).

Peach Tree Borer (see Aegeria exitiosa). Pear, Quadraspidiotus perniciosus on, in Algeria, 364; pests of, in W. Australia, 364; pests of, in Britain, 35, 272, 273; pests of, in France, 160, 229, 349, 362, 370; pests of, in Holland, 67, 91; Anastrepha fraterculus experimentally infesting, in Mexico, 138; Eulecanium bituberculatum on, in Persia, 16; pests of, in U.S.A., 24, 73, 74, 109, 176, 374.

Pear Psylla (see Psylla pyricola). Pear Thrips (see Taeniothrips inconse-

Pecan, pests of, in U.S.A., 27, 108, 176, 373, 374.

pectinicornis, Corymbites.

Pectinophora gossypiella (see Platyedra). pectoralis, Platycampus.

Pediculoides graminum, on barley in

Britain, 273.

Pediculoides ventricosus, associated with Anobium punctatum in New Zealand, 211.

Pediculopsis graminum (see Pediculoides).

Pegohylemyia (see Hylemyia).

Pegomyia hyoscyami (var. betae on beet), in Britain, 330-332; in Denmark, 8, 116; bionomics of, 8, 331, 332; bait-sprays against, 331, 332.

pellionella, Tinea.

pellucens, Fundella.

pellucida, Camnula. peltigera, Heliothis (Chloridea).

Pemphigus, on Populus suaveolens in Central Asia, 156.

Pemphigus cornicularius, on Pistacia terebinthus in Jugoslavia, 328.

penicillatus, Dolycoris. pennipes, Trichopoda.

Pennisetum, Zonocerus variegatus on, in Belgian Congo, 280.

Pennisetum typhoideum, grasshoppers on,

in India, 47, 207.

Pennsylvania, orchard pests in, 140, 141, 195, 196, 197, 317; miscellaneous pests in, 105, 176, 247; Reduviid destroying Popillia japonica, etc., in, 145.

pennsylvanicus, hercu-Camponotus leanus; Chauliognathus.

Pentachlordiphenvlethane (see DDT). Pentachlorphenol, in spray against

insects and weeds, 141: treatments of timber with, against pests, 12, 172.

pentagona, Pseudaulacaspis (Diaspis). Pentalonia nigronervosa, on greenhouse plants in France, 362.

Pentatoma rufipes, bionomics and control of, on fruit trees in Britain, 115,

116. Pentodon idiota, survival of, in crop rotations in U.S.S.R., 157.

pepon, Phorma.

Peppers (see Capsicum).

peregrina, Schistocerca (see S. gregaria).

peregrinator, Orchidophilus.

peregrinus, Pantomorus. perezi, Steropleurus.

pergandei, Parlatoria.

pergratialis, Monoptilota.

Pericerya (see Icerya).

Perilampus fulvicornis, possibly hyper-parasitic on Tetralopha scortealis in Virginia, 305.

Perilitus coccinellae, bionomics of, in Britain, 114.

Perilla Oil, increasing adhesiveness of lead-arsenate sprays, 220.

Periplaneta americana, killed by DDT aerosol residue in greenhouses

Maryland, 240; sites of action DDT and nicotine in, 5.

parasite of Noctuids Perisierola, Texas, 3.

Perissopterus buscki, parasite of Coccids in Nyasaland, 63.

perniciosus, Pseudococcus; Quadraspidiotus (Aspidiotus).

Peronea variana (see Acleris).

Perrisia (see Dasyneura).

Persia, Coccids in, 16: Schistocerca gregaria in, 352.

persicae, Myzus.

persicae-niger, Anuraphis.

Persimmon (see Diospyros kaki).

persuasoria, Rhyssa.

Peru, new Capsid on cotton and beans in, 128.

Petrol, against Lamiids in coffee, 103; and creosote, treatment of timber with, against Camponotus spp., 27; soil treatment with, against Epitrix hirtipennis, 316.

Petroleum Ether, as a solvent, 359.

Petunia, experimentally infected with turnip viruses, 182; doubtful relation of, to Argentine curly-top virus, 379,

peucedani, Aceria.

Phaedon brassicae, bionomics and control of, in China, 119.

Phaedroctonus (see Nemeritis).

Phaeogenes arcticus, parasite of Acleris variana in Br. Columbia, 379.

phąeorrhoea, Nygmia.

Phalaenopsis (see Orchids).

Phanerotoma ornatulopsis, sp. n., parasite of Earias biplaga in Belgian Congo. 81.

Phanerotoma phycitinoma, sp. n., parasite of Cryptoblabes gnidiella in Belgian Congo, 81.

Phanurus (see Telenomus).

pharaonis, Monomorium

phaseli, Lampides (Jamides).

phaseoli, Agromyza; Conotrachelus.

phaseolunata, Agromyza.

Phaseolus vulgaris (see Beans).

Pheasants, question of poisoning of, by arsenicals, 83.

Pheidole, associated with Pseudococcus njalensis in W. Africa, 46.

Phenacaspis prunorum, on fruit trees in Persia, 16.

Phenacoccus colemani, method of feeding

Phenacoccus solani, on Echium in California, **109.**

Phenol, as a bee repellent in sprays, 55. Phenothiazine, against Coleoptera, 178, 316; against Lepidoptera, 175, 197, against Rhagoletis pomonella,

50; favouring mites and Aphids on apple, 175, 197; in conditioned dusts, 197, 316; spraying with, 50, 175, 178, 197, 316; adhesives and spreaders for, 50, 175, 316; and lead arsenate, 197; and oil emulsions, 175; micronised form of, 175, 178, 197, 316;

formulae containing, 50, 175, 197, 316. Phenothioxin (Phenoxathiin), unsatisfactory against cotton pests, 178, 179. Phenyl Acetic Acid, in solid baits for

Cydia pomonella, 309. phidilealis, Hellula.

philadelphicus, Proctacanthus.

Philaenus leucophthalmus, food-plants of, in Hawaii, 274; DDT aerosols against, on peas in U.S.A., 243.

Philippia (see Filippia).

Philippines, risk and prevention of introduction of pests into Hawaii by aircraft from. 2.

Phillyrea media, new Aphid on, in Greece, 326.

Phleum pratense, experiments with Hylemyia coarctata and, in Britain, 368.

Phloeophagosoma cucurbitae, sp. n., on cucurbits in Senegal, 206.

Phloeophthorus spinulosus, parasite of, on spruce in Finland, 39.

Phloeothrips ficorum (see Gynaikothrips). Phloeotribus scarabaeoides, on olive in Greece, 326.

Phlyctaenia rubigalis, measures against, on greenhouse rose in Ohio, 78.

phoenicis, Asterolecanium.

Phoenicococcus marlatti, on date palm in Persia, 16.

Phoenix canariensis, Coccotrypes dactyliperda on, in California, 375.

Phoenix dactylifera (see Date Palm).

Phorma pepon, new parasite of, on coffee in Belgian Congo, 80.

Phorocera, introduced into U.S.A. against armyworms, 179.

Phorostoma, parasite of Bixadus sierricola in Belgian Congo, 96.

Photinia spp., Cydia molesta on, in California, 109, 373.

Phthorimaea (see Gnorimoschema).

phthorimaeae, Chelonus.

Phthorophloeus spinulosus (see Phloeophthorus).

phycitinoma, Phanerotoma.

phyllanthi, Parlatoria.

Phyllobius oblongus, bionomics and control of, on pear in France, 349.

Phyllocoptes destructor (see Vasates).

Phyllocoptes eurynotus (see Vasates).

Phyllocoptes gracilis, on raspberry in Washington, 75, 160.

Phyllocoptes lycopersici, DDT retarding infestation by, on tomato in New South Wales, 385.

Phyllocoptes oleivorus (see Phyllocoptruta).

Phyllocoptes schlechtendali, sprays against, on pear in Washington, 73.

Phyllocoptruta oleivorus (on Citrus), in Australia, 127, 384; in Cyprus, 371; sprays and dusts against, 127, 371, 384; DDT possibly increasing infestation by, 384.

Phyllopertha (see Anomala).

Phyllophaga (see Lachnosterna).

Phyllotoma nemorata, spread of, on birch in Quebec, 173.

Phyllotreta (on crucifers), in Britain, 114, 256; in Denmark, 7; flight habits of, 114; dusts against, 7, 256.

Phyllotreta atra, on crucifers in Britain, 114.

Phyllotreta columbiana, measures against, on cabbage in Washington, 75.

Phyllotreta consobrina, on crucifers in Britain, 114.

Phyllotreta cruciferae, on crucifers in Britain, 114.

Phyllotreta diademata, on crucifers in Britain, 114.

Phyllotreta nemorum, on crucifers in Britain, 114.

Phyllotreta nigripes, on crucifers in Britain, 114.

Phyllotreta undulata, on crucifers in Britain, 114.

Phyllotreta vittata, measures against, on crucifers in China, 118, 344.

Phyllotreta vittula, on crucifers in Britain,

Phylloxera vitifoliae (on vines), bionomics and control of, in Argentina, 299; study of fungus infesting, in France, 33.

Phymatodes nitidus, in Sequoia sempervirens in California, 31.

Physalis, Agalliana ensigera transmitting curly-top virus to, in Argentina, 379.

Physalis alkekengi, Leptinotarsa decemlineata not feeding on, 264.

Phytometra (see Plusia).

Phytomonas rubrilineans, favouring infestation of sugar-cane by Rhab-doscelus obscurus in Queensland, 1.

Phytomyza lateralis, bionomics and control of, in chicory in Belgium, 212, 213. Phytonomus (see Hypera).

Phytophaga (see Mayetiola).

Phytoptus avellanae, Cecidophyes vermiformis in galls of, on filbert in Oregon, 111.

Picea (see Spruce).

Picea abies, Hylemyia anthracina on, in Finland, 38; Neodiprion sertifer on, in Sweden, 131.

Picea engelmanni, Dendroctonus engelmanni on, in Colorado, 176.

Picea excelsa (see P. abies).

Picea rubra, pests of, in Nova Scotia, 129. piceae, Chermes (Adelges).

piceaperda, Dendroctonus.

piceatus, Kalotermes (Cryptotermes).

Picric Acid, use of, in determining HCN in fumigated insects, 150.

Picromerus bidens (predacious on noxious insects), bionomics of, in Belgium, 213; in Sweden, 130.

pictipennis, Platyspathius.

pictipes, Aegeria; Ephialtes (Epiurus). Pieris brassicae (on crucifers), in Denmark, 7; in Finland, 271; in U.S.S.R., 131; experiments with Pentatomids and, 213; diseases of, and their utilisation, 131, 213, 271; tests of DDT and analogues against, 7, 255.

Pieris rapae (on crucifers), in Denmark, 7; in New South Wales, 383, 384, 386; parasites established against, in New Zealand, 323; use of fungus against, in U.S.S.R., 131; in U.S.A., 75, 313, 318, 337; insecticides against, 7, 313, 318, 337, 383, 384, 386; tests of DDT analogues on, 255.

Piezodorus hybneri (rubrofasciatus), measures against, on beans in Queens-

land, 126.

Pigweed, Black (see Portulaca oleracea). pilosus, Atomus; Paratetranychus.

pimpinellae, Anthrenus.

Pimpla brevicornis (see Ephialtes).

Pimpla curvipes (see Thalessa). Pimpla kangasi (see Ephialtes).

Pimpla sanguinipes, parasite of Autographa egena in California, 30.

Pine, Hylotrupes bajulus in, in S. Africa, 12; Rhyacionia buoliana on, in Argentina, 371; pests of, in Canada, 129, 173, 320, 329; pests of, in Finland, 9, 38, 39, 42, 99; pests of, in France, 257, 329, 371; Dendrolimus pini on, in Germany, 99; pests of, in Holland, 87, 88, 93; Neodiprion sertifer on, in Sweden, 130, 131; Bupalus piniarius on, in U.S.S.R., 131; pests of, in U.S.A., 57, 158, 176, 236, 237, 311; injection of chemicals into, for timber preservation, 56.

Pine, Jack (see Pinus banksiana).

Pine, Lodgepole (see *Pinus contorta latifolia*).

Pine, Ponderosa (see Pinus ponderosa).

Pine, Red (see Pinus resinosa).

Pine, White (see Pinus strobus).

Pine Bark Weevil (see Aesiotes notabilis).

Pine Beetle, Mountain (see *Dendroctonus monticolae*).

Pine Budworm, Jack (see Harmologa fumiferana).

Pine Geometrid (see *Bupalus piniarius*). Pine Lasiocampid (see *Dendrolimus pini*). Pine Sawfly (see *Diprion pini*).

Pine-tar Oil, in baits for Cydia pomonella, 308.

Pineapple, Pseudococcus njalensis on, in Gold Coast, 46; pests of, in Hawaii, 134, 307.

Pineapple Mealybug (see *Pseudococcus brevipes*).

Pineus (see Chermes).

pinguioides, Promasipoda.

pinguis, Agromyza (Ophiomyia).

pini, Cinara; Dendrolimus; Diprion; Eupithecia.

piniarius, Bupalus; Hylurdrectonus. pinifoliae, Chermes (Pineus); Chionaspis.

piniperda, Myelophilus. piniphilus, Pissodes.

Pinus (see Pine).

Pinus banksiana, pests of, in Canada, 173, 320.

Pinus caribaea, Cinara carolina on, in S. Carolina, 311.

Pinus contorta var. latifolia, Dendroctonus monticolae on, in Canada, 360.

Pinus echinata, pests of, in U.S.A., 158,

Pinus edulis, pests of, in U.S.A., 57.

Pinus insignis (see P. radiata). Pinus maritima (see P. pinaster).

Pinus mugo (montana), Neodriprion sertifer on, in Sweden, 131.

Pinus nigra, experimentally attacked by Gilpinia frutetorum, 236.

Pinus pinaster, pests of, in France, 329, 371.

Pinus ponderosa, Harmologa fumiferana on, in U.S.A., 176.

Pinus radiata, Rhyacionia buoliana on, in Argentina, 371.

Pinus resinosa, Gilpinia frutetorum on, in Connecticut, 236, 237; Pissodes strobi on, in Quebec, 173.

Pinus rigida, pests of, in U.S.A., 158,

Pinus strobus, pests of, in Canada, 129, 173; Cinara strobi on, in S. Carolina, 311.

Pinus sylvestris, sawflies on, in Sweden and U.S.A., 130, 236.

Pinus virginiana, Cinara spp. on, in U.S.A., 311.

Pinyon (see Pinus edulis).

Piophila casei, natural sources of, 150.

piri, Anthonomus.

pisi, Bruchus (see B. pisorum); Macro-siphum (see M. onobrychis).

pisorum, Bruchus.

Pissodes approximatus, on Pinus resinosa in Connecticut, 237.

Pissodes notatus, parasites of, in Finland, 39, 40; measures against, on Pinus pinaster in France, 371.

Pissodes piniphilus, new parasite of, in Finland, 39.

Pissodes strobi, on Pinus spp. and spruce in Quebec, 173.

Pissodes validirostris, new parasite of, in Finland, 39, 40.

Pistacia, Capnodis cariosa on, in Palestine, 369.

Pistacia terebinthus, pests of, in Jugo-slavia, 327, 328.

Pistacia vera, Coccids on, in Persia, 16.

pistaciae, Eriophyes; Lepidosaphes (Mytilococcus); Pulvinaria. Pithecellobium trapezifolium, dust from,

as an insecticide, 304.

Pittosporum, Coccid on, in California, 109.

pityocampa, Thaumetopoea.

Pityogenes (see Ibs).

Plant Pest Legislation, in U.S.A., 112, 291, 292; summaries of, in other countries, 112, 292.

Plantago, new Agromyzid on, in U.S.A.,

Plants, studies of, as sources of insecticides, 137, 217, 218, 301, 303, 304, 359, 381; bibliography of, as sources of insecticides, 32, 192.

Plaster of Paris, in solid baits for Cydia pomonella, 308.

platensis, Tetrastichus.

Plathypena scabra (in U.S.A.), parasite of, 3; tests of dusts against, on beans,

144, 317.

Platycampus ovatus, on larch in Holland,

Platycampus pectoralis, on larch in Holland, 69.

platycephalus, Egius.

Platyedra gossypiella (Pink Cotton Bollworm), 159; in W. Africa, 96; in W. Australia, 366; Bracon greeni reared on, in Bihar, 165; in Morocco, 66; in Sudan, 166; in Tanganyika, 167; in U.S.A., 178, 315; legislation against importation of, into U.S.A., 291; varieties of cotton in relation to, 166, 178; bionomics of, 96, 178; field measures against, 178, 315; fumigation of cottonseed against, 66.

Platyedra scutigera, measures against, on cotton in Queensland, 166.

platyhypenae, Euplectrus.

Platymoideus acutus, transmitting virus disease of lucerne in Washington, 74. Platyphytoptus jonesi (see Setoptus).

Platyplectrus natadae, propodeum of, 47. Platyspathius spp., associated Coleoptera in Acacia log in Nyasaland, 63.

Plecoptera reflexa, parasites of, on Dalbergia sissoo in India, 285, 286.

plecopterae, Euplectrus (see E. parvulus);

Microgaster; Rogas.

Plemeliella abietina, in spruce cones in Finland, 38, 64; parasite of, 38. Plesiocoris rugicollis, spray against, on

apple in Britain, 273.

Plesiothrips panicus, on maize in Hawaii,

Pleurotropis, hyperparasites of Antestia in Uganda, 278.

Pleurotropis nigripes, parasite of Apanteles transvaalensis in Nyasaland, 63. Plodia interpunctella, measures against, in dried fruit, etc., in Morocco and

W. Australia, **65, 365.**

Plum, 253; Quadraspidiotus perniciosus on, in Algeria, 363, 364; pests of, in Australia, 208, 364, 365; pests of, in Britain, 35, 115, 272; pests of, in France, 229, 370; pests of, in Holland, 67, 91; Eulecanium bituberculatum on, in Iraq, 16; Anastrepha fraterculus experimentally infesting, in Mexico, 138; Aphids on, in Norway, 7; pests of, in Palestine, 132, 369; Coccids on, in Persia, 16; Hoplocampa spp. on, in Sweden, 37; pests of, in U.S.A., 24, 108, 245, 321, 374, 375; methyl-bromide fumigation of fruits of, **363, 365.**

Plum Curculio (see Conotrachelus

nenuphar).

Plumbago capensis, Icerya purchasi on, in India, 287.

Plusia argentifera, dust against, on beans in Queensland, 126.

Plusia brassicae, on cabbage in U.S.A., 75, 313, 318; insecticides against, 313, 318.

Plusia californica, on lucerne in California, **111.**

Plusia chalcites, dust against, on beans in Queensland, 126.

lucerne Plusia circumflexa, on Tadzhikistan, 253.

Plutella maculipennis (on crucifers), in Australia, **231, 383, 384, 385, 386**; in Fiji, 101; in New Zealand, 209, 323; in Nyasaland, 63; in U.S.A., 75, 319; parasites and biological control of, 63, 323; insecticides against, 101, 209, 231, 319, 383, 384, 385, 386.

Poa annua, Hylemyia coarctata on, in Britain, 368.

Poa pratensis, Conoderus bellus damaging, in Kentucky, 183.

Poa trivialis, Hylemyia coarctata on, in Britain, 368.

podana, Tortrix (Cacoecia).

Podisus nigrolimbatus, predacious Orgyia antiqua in Chile, 352.

Podisus serieventris, predacious Acleris variana in Br. Columbia, 379.

Podocarpus dacrydioides, susceptibility of timber of, to Anobium punctatum in New Zealand, 211.

Podocarpus spicatus, resistance of timber of, to Anobium punctatum in New Zealand, 211.

Podocarpus totara, resistance of timber of, to Anobium punctatum in New Zealand, 211.

Podonta daghestanica, survival of, in crop rotations in U.S.S.R., 157.

Poecilonota variolosa, parasite of, on Populus tremula in Finland. 41.

Hibiscus Poeciloscytus cognatus, on cannabinus in Uzbekistan, 155.

Poeciloscytus vulneratus, on Hibiscus cannabinus in Uzbekistan, 155.

Poinciana regia, Lyncestis acontioides on, in Florida, 356.

Poinsettia, Pseudococcus citri on, in U.S.A., 79.

Polia oleracea, bionomics of, in Finland,

Pollination, wild bees causing, in lucerne, 260.

Pollinia pollini (on olive), in California, 375; in Greece, 326.

Polyethylene Glycol, cetyl ethers of, increasing permeability of cuticle, 60.

Polygonum pennsylvanicum, 304.

Polygraphus rufipennis, in spruce and

Abres in Succession Polyhedral Disease, in mortality Lepidoptera infected with, 213.

polytoma, Gilpinia (Lophyrus).

pomaria, Typhlocyba.

Pomegranate, Pseudococcus citri on, in Persia, 16; Pteroma on, in Travancore, 283; Cydia molesta developing on, 109.

pometaria, Alsophila.

pomi, Aphis.

pomonella, Cydia (Carpocapsa); Rhago-

pomorum, Anthonomus. Pongamia glabra, 165.

Popillia japonica, in Nova Scotia, 130; in U.S.A., 51, 107, 141, 145, 176, 194, 210, 233, 234, 244, 245, 304, 341; insects predacious on, 145, 194; parasites and disease of, and their utilisation, 51, 234; bacillus of Odontria not infecting, 210; treatments of plants against adults of, 141, 234, 244, 245; method evaluating sprays against, 304; other measures and experiments against, 107, 176, 245, 341.

popilliavora, Tiphia.

Poplar (Populus), Aphids on, in Central Asia, 156; pests of, in Canada, 129, 173. (See Aspen.)

Poppy (see Papaver somniferum).

Popular Names (of insects), list of, in U.S.A., 352.

populnea, Saperda. Populus (see Poplar).

Populus tremula (see Aspen).

Porizon, doubtful status of, 252; species of,* introduced into U.S.A. against Listroderes obliquus, 104, 179.

Porizon argentinensis, sp. n.* (parasite of Listroderes), in Argentina, 104; imported into U.S.A., 104.

Porizon moderator, F., doubtful identity of, 252.

Porizon parkeri, sp. n.* (parasite of Listroderes), in S. America, 104; imported into U.S.A., 104.

Porthesia chrysorrhoea, auct. (see Nygmia phaeorrhoea).

Porthetria (see Lymantria).

Porto Rico, Kalotermes brevis in, 172, 344; pests of leguminous plants in, 64, 104, 218, 219; miscellaneous pests in, 31, 218, 219; Lepidoptera on sugar-cane in, 31, 64, 179, 320; beneficial insects and biological control in, 31, 32, 64, 179, 219, 320; summary of plant quarantines in, 292; pests intercepted in quarantine in, 183; Hambletonia pseudococcina imported into U.S.A. from, 179, 219.

Portulaca, Agalliana ensigera on, in Argentina, 380.

Portulaca oleracea, Hypurus bertrandi on, in France, 349; Hymenia recurvalis on, in Queensland, 169.

postica, Hypera (see H. variabilis).

Potash, manuring with, reducing infestation of jute by Apion corchori,

Potassium Ammonium Silicate, Australian pyrophyllite containing, 231.

Potassium Cyanide, treatment of ants' nests with solution of, 297.

Potassium Fluosilicate, dusting with, against Anthonomus grandis, 356.

Potassium Metabisulphite, 15. Potassium Palmitate, 239.

Potato, pests of, in Australia, 23, 382, 387; Gnorimoschema operculella on, in Bermuda, 103, 300; pests of, in Britain, 161, 370; pests of, in Canada, 264, 319; Gryllotalpa on, in Egypt, 19; Gnorimoschema operculella on, in Eritrea, 154; Leptinotarsa decemlineata on, in France, 9, 10, 32, 83, 84, 327; Heliothis armigera on, in Hawaii, 274; L. decemlineata on, in Spain, 100; L. decemlineata on, in Switzerland, 327; pests of, in U.S.A., 28, 29, 43, 53, 72, 74, 100, 149, 180, 223, 224, 226, 235, 238, 239, 240, 243, 264, 337, 355, 358; Aphids transmitting leaf-roll of, 74, 226; effect of, on infestation of cereals by *Hylemyia*

^{*}In view of divergent opinions regarding the status of Porizon, the generic position of the species cited in these references is uncertain. Porizon argentinensis and P.parkeri are more closely allied to Thersilochus than to Porizonidea but differ in some characters from Thersilochus as understood by European authors.-Ed.

463 INDEX.

coarctata, 216, 367; (tubers), Gnorimoschema operculella in, 65, 154, 301; influence of moisture of, on rearing of Macrocentrus on G. operculella, 305; DDT mixed with, against Pseudococcus spp., 334.

Potato Beetle, Colorado (see Leptino-

tarsa decemlineata).

Potato Flea-beetles (see Epitrix cucumeris and E. tuberis).

Potato Leafhopper (see *Empoasca fabae*). Potato Psyllid (see Paratrioza cocker-

Potato Tuber Moth (see Gnorimoschema

operculella).

Potentilla canadensis, Tyloderma fragariae on, in Kentucky, 292.

P.P.666, 256.

praefica, Prodenia.

praemorsus, Leptostylus.

praetor, Euphorus.

pratensis, L., Lygus.

pratensis, auct., Lygus (see L. obline-

Prays oleellus (on olive), parasites of, in Greece, 326; in Spain, 267.

Prenes (see Panoquina).

prenidis, Ooencyrtus.

pretiosum, Trichogramma.

Prickly Pear (see Opuntia).

Prince Edward Island, Aphids potato in, 319.

princeps, Dirphya (Nitocris).

Prionomeion, systematic position of, 14. Priophorus pallipes, bionomics of, on plum in Britain, 272.

Pristiphora erichsoni (on larch), Holland, 69; parasites and disease of, in Quebec, 173.

Pristiphora laricis, on larch in Holland,

Pristiphora pallipes, bionomics and control of, on gooseberry in Britain,

Pristiphora wesmaeli, on larch in Holland, 69.

Pristonychus terricola, damaging stored apples in Britain, 59.

Privet, Parlatoria oleae on, in California,

Proatractonema sciarae, gen. et sp. n., infesting Sciara in Denmark, 160.

procephali, Aphidius.

Proceras sticticraspis (on sugar-cane in India), use of Trichogramma minutum against, 207; effects of earthing-up of cane on, 187, 188.

Proceras venosatus, use of Trichogramma minutum against, on sugar-cane in

India, 207.

Prociphilus oleae, sp. n., on olive and Phillyrea media in Greece, 326.

Proctacanthus spp., predacious on Popillia japonica in New York, 194.

Prodenia, parasites of, in Texas, 3. Prodenia eridania (see Laphygma).

Prodenia litura, Euplectrus gopimohani reared on, in Bengal, 164; inter-cepted in aircraft in Pacific Is., 2; dusts against, on cabbage in Queensland, 231; new parasite of, Senegal, 96.

Prodenia praefica, on lucerne in Cali-

fornia, 111.

Prodioctes haematicus, bionomics and control of, on Elettaria cardamomum in Travancore, 283.

prolixus, Rhodnius

prolongata, Xiphydria.

Promasipoda pinguioides, bionomics of, in Brazil, 5, 6.

Prontaspis citri (see Unaspis).

Propane, use of, for producing aerosols,

Propylea quatuordecimpunctata, experiment with Perilitus coccinellae and, 114.

Propylene Dichloride (see Dichlorpropane).

prosectus, Alaus.

Prosena, parasite of Dermolepida albohirtum in Queensland, 1.

proserpina, Megamelus.

Prosopis chilensis, new Bruchid in seeds of, in W. Indies, 64.

Prospaltella berlesei, introduced parasite Pseudaulacaspis pentagona France, 362.

Prostephanus punctatus, in barrel staves in Georgia, 194.

proteus, Omophlus; Parlatoria.

Protodiaspis cinchonae, sp. n., on Cinchona calisaya in Bolivia, 112.

Protoleucania albilinea, on cereals in New Jersey, 203.

Protoparce (on tobacco in U.S.A.), Engytatus geniculatus predacious on, 347: measures against, 178.

Protoparce sexta, bionomics and control of, on tobacco in U.S.A., 335, 356.

Protrigonia zizanialis, bionomics of, on Moringa oleifera in Travancore, 288.

Prune, Paratetranychus pilosus on, in Br. Columbia, 319; pests of, in U.S.A., 24, 143, 149.

prunifoliae, Rhopalosiphum. prunorum, Phenacaspis.

Prunus, Cydia molesta on, in California, 109; relation of Myzus persicae to species of, 367. (See Cherry, Peach, Plum, etc.)

Prunus divaricata, Rhynchites bacchus on, in France, 370.

Prunus emarginata, Rhagoletis spp. on, in U.S.A., 108.

Prunus institia, Anuraphis schwartzi on, in Britain. 272.

Prunus nigra, winter food-plant of Myzus persicae, 226, 367; methods of destroying, 226.

Prunus padus, Anuraphis schwartzi on,

in Britain, 272.

Prunus spinosa, insects on, in Britain, 185, 272.

Psallus ancorifer, dust against, on onion in Oregon, 247.

Psallus seriatus, dusts against, on cotton in U.S.A., 180, 345.

Psara bipunctalis, on soy bean in Porto Rico, 219.

Pseudabutilon lozani, Platyedra gossypiella overwintering on, in Texas, 178. Pseudaonidia trilobitiformis, on fig and

lemon in Nyasaland, 63.

Pseudaphycus, use of, against Pseudococcus comstocki in U.S.A., 234, 336; effect of DDT spray deposits on, 336.

pseudaspiaiotus, Genaparlatoria.

Pseudaspidoproctus vayssieriellus, sp. n., parasites of, on palms in Belgian Congo, 82.

Pseudaulacaspis pentagona, food-plants and introduced parasite of, in France,

362

Pseudoarchytopsis, introduced into U.S.A. against armyworms, 179.

pseudococcina, Hambletonia.

Pseudococcus, DDT against, in stored potatoes in Brazil, 334.

Pseudococcus adonidum, on Aleurites in Nyasaland, 63.

Pseudococcus brevipes (Pineapple Mealybug), in Florida, 179, 219; in Hawaii, 135, 274, 307; in Porto Rico, 219; on Gliricidia, 274; effects of, on pineapple, 135, 307, 308; artificial feeding and oral secretions of, 307; natural enemies and biological control of, 179, 219, 274; soil treatments against, 135.

Pseudococcus citri, experimentally transmitting strains of swollen-shoot of cacao in Gold Coast, 102; on mulberry in Iraq, 16; parasites of, on beans in Nyasaland, 63; on pomegranate and vine in Persia, 16; on Citrus in Spain, 269; measures and experiments against, on greenhouse plants in U.S.A., 79, 240.

Pseudococcus comstocki, in stored potatoes in Brazil, 334; on apple in U.S.A., 31, 106, 179, 234, 336; method of feeding of, on various plants, 31; parasites and biological control of, 179, 234, 336; DDT against, 106,

334.

Pseudococcus exitiabilis (see P. njalensis).

Pseudococcus filamentosus, establishment of Scymnus uncinatus against, on Ceratonia in Hawaii, 274.

Pseudococcus juniperi, spray against, on Juniperus in Kansas, 200.

Pseudococcus maritimus, DDT against, in stored potatoes in Brazil, 334.

Pseudococcus njalensis, transmitting swollen-shoot of cacao in Gold Coast, 46, 102; food-plants and distribution of, in W. Africa, 46; ants associated with, 46; characters and synonymy of, 46.

Pseudococcus perniciosus, on Aleurites in

Nyasaland, 63.

pseudocuneatus, Tenuipalpus. pseudoeceticola, Tetrastichus.

pseudohesperidum, Coccus. pseudomagnoliarum, Coccus.

Pseudoparlatoria parlatorioides, on orchids in U.S.A., 109, 315; DDT spray against, 315.

pseudosolani, Myzus (see Macrosiphum

solani).

Pseudotsuga taxifolia, pests of, in Br. Columbia, 320, 378; pests of, in U.S.A., 31, 76, 160, 176; recorded as Abies, 76.

psidii, Conotrachelus.

Psila rosae (on carrots), in Britain, 44, 45; in Br. Columbia, 319; bionomics of, in Germany, 68; in Oregon, 250; in Switzerland, 43; DDT against, 43-45, 250; naphthalene against, 319.

Psyche atra (see Oreopsyche angustella). Psylla mali, sprays against, on apple in Denmark, 8.

Psylla pyricola, measures against, on pear in U.S.A., 74, 176; colour response in, 176.

Psylla simulans, bionomics of, on pear in Britain, 272.

Psyllia (see Psylla).

Psyllid Yellows, caused by Paratrioza cockerelli in potato, 358.

Pteroma, bionomics of, on pomegranate in Travancore, 283.

Pteromalus puparum, utilisation of, against Pieris rapae in New Zealand, 323.

Pteronus (see Nematus).

pterophori, Ephialtes (Epiurus).

Ptilinus fuscus, on Populus tremula in Finland, 41.

Ptinus tectus, in stored ants' pupae in Czechoslovakia, 212; mass rearing of, 61.

Ptychomyia selecta, parasite of Pristiphora erichsoni in Quebec, 173.

pubescens, Graphops.

Puccinia spp., varieties of wheat resistant to Mayetiola destructor and, 5. pulicaria, Chaetocnema.

INDEX. 465

pulverea, Holcocera.

Pulvinaria betulae, on apricot in Persia,

Pulvinaria flavescens, fungus infesting, in Argentina, 299.

Pulvinaria floccifera, fumigation against, on Camellia in California, 109; on Citrus in Persia, 16.

Pulvinaria jacksoni, on Aleurites fordi in

Nyasaland, 63.

Pulvinaria mesembryanthemi, bionomics of, on Mesembryanthemum in France, 349.

Pulvinaria pistaciae, on Pistacia vera in Persia, 16.

pumila, Agathis (Bassus).

Pumpkin, little attacked by Listroderes obliquus in W. Australia, 23; Diabrotica melanocephala transmitting Bacillus tracheiphilus to, in U.S.A., 148.

punctata, Hypera; Xanthopimpla.

punctatum, Anobium.

punctatus, Eysarcoris (Asopus); Prostephanus.

punctiferalis, Dichocrocis.

punctiventris, Bithotrotis.
punctiventris, Cleonus (Bothynoderes).
punctulatus, Deraeccoris (Camptobrochis).
Punjab, Earias spp. on cotton in, 282,
287; miscellaneous pests in, 47, 117,
118, 285, 296; natural enemies of

honey bees in, 297.
puparum, Pteromalus.
pupivora, Trichospilus.
purchasi, Icerya (Pericerya).

tantanas Clausonia

purpurea, Clausenia.

Purslane (see Portulaca oleracea).

pusilla, Agromyza. pusillus, Lophocateres.

putrescentiae, Tyrophagus (Tyroglyphus). Pyemotes ventricosus (see Pediculoides).

pygmaeus, Cephus. pyramicus, Dorymyrmex.

Pyrausta nubilalis (European Corn Borer), in Canada, 320; experiments with Bacterium prodigiosum and, in U.S.S.R., 131; in U.S.A., 53, 136, 137, 177, 221, 235, 236; resistance of inbred lines of maize to, 221; on potato, 53, 136, 137; parasites and biological control of, 177, 320; not parasitised by Agrothereutes stokesi, 139; insecticides against, 53, 136, 137, 235, 236.

Pyrax, 311, 312, 319.

Pyrax ABB, 301, 302, 312.

Pyrene, protecting wood from termites, 172.

Pyrethrins, content of, in pyrethrum dusts and sprays, 49, 77, 88, 89, 91, 147, 200, 243, 244, 301, 311, 312, 313, 317, 334, 338, 339; analysis of DDT and, in kerosene, 256.

Pyrethrum (including extracts), against ants, 238; against Aphilds, 73; against Jassids, 244, 267, 312, 313, 374, 376, 377; against Lygus, 54, 261, 281, 312; against other Hemiptera, 89, 281, 301, 311; against Coleoptera, 23, 49, 62, 72, 76, 105, 149, 270, 334, 338, 339; against Lepidoptera, 49, 78, 91, 147, 152, 200, 243, 317, 338, 339; and oils or emulsions, treatment of maize ears with, against Heliothis, 147, 177; testing of, on Bombyx mori, 91; against Melanoplus femur-rubrum, 312; against mites, 77; against sawflies, 37, 88, 89; against thrips, 24, 338; toxicity of, to honey bees, 55; penetration of insect cuticle by, factors affecting insecticidal action of, 376; aerosols of, 241, 338; dusting with, 23, 54, 55, 72, 73, 76, 78, 88, 89, 91, 149, 152, 200, 243, 244, 261, 267, 270, 281, 301, 311, 312, 313, 317, 338, 377; carriers for, 23, 89, 311, 312, 317; oil-impregnated dust of, 338; decreased toxicity of, stored in calcium cyanamide, 89; and DDT, 157, 256; and derris, 89, 317; and nicotine, 54; atomised sprays of oil containing, 62, 339, 377; vapour spray of diesel oil containing, 374; and rotenone insecticides, 24, 338; sesame oil as activator for, 338; and soap, 49, 54; and sulphur, 244, 261, 267, 312, 317; and thiocyanates, 339, 374; and tobacco, 152; apparatus for removing solvents from concentrates of, 360; supplement to book on, 352; Lasioderma serricorne damaging anti-mosquito smoke spirals containing, 101.

Pyrethrum Marc, in spray against Lepidoptera, 75.

Pyrethrum Plant (see Chrysanthemum cinerariaefolium).

pyri, Eriophyes; Parlatoreopsis (Parlatoria).

pyricola, Psylla.

Pyridine, fumigation with, against *Tyrophagus lintneri* in mushroom houses, 77.

Pyrilla, on sugar-cane in India, 187.

pyrina, Zeuzera.

Pyrocide, 200, 261.

Pyroderces dendrophaga, on sorghum in Queensland, 388.

Pyroderces risbeci, sp. n., on sorghum in Senegal, **96.**

Pyronota festiva (see Calonota).

Pyrophorus luminosus, establishment of, against Clemora smithi in Barbados, 159.

Pyrophyllite, as a carrier for dusts, 50, 105, 137, 144, 149, 180, 197, 224, 235, 236, 238, 239, 244, 245, 247, 249, 250, 301, 302, 311, 312, 315, 338, 345, 382, 383, 386; composition of Australian, as a dust carrier, 231; spray and paint suspension of DDT in, 238.

Pyruvic Acid, in solid baits for Cydia

pomonella, 309.

Pythium, D-D mixture against, 135.

Q.

Quadraspidiotus ostreaeformis, on apple

in Persia, 16.

Quadraspidiotus perniciosus (on fruit trees), in Algeria, 363, 364; in Kashmir, 293; in U.S.A., 73, 75, 321; on cranberry, 75; sprays against, 73, 75, 293; fumigation of harvested fruits against, 363, 364.

quadridentata, Ascogaster. quadrilineatus, Ceroplastes; Opsimus.

quadripes, Xylotrechus.

quadripunctella, Incurvaria (Lampronia). quadriseriata, Ctenolepisma.

Quarantine, pests intercepted in: in Pacific Is., 2, 183; in Porto Rico, 183; in U.S.A., 112, 183, 277.

Quassia Sprays, ineffective against Eurydema oleraceum, 89; against Hoplocampa spp., 37, 229; preparation of, 37, 229.

Quassia amara, 229.

quatuordecimpunctata, Propylea.

Quebec, forest pests in, 173, 263, 377; Pyrausta nubilalis on maize in, 320; beneficial insects and biological con-

trol in, 173, 263, 320.

Queensland, pests of Araucaria cunninghami in, 258-260; pests of Citrus in, 126, 127; cotton pests in, 166; pests of sorghum in, 387, 388; sugar-cane pests in, 1; pests of vegetables in, 125, 126, 169, 231, 288, 386, 387; pests of stored products in, 126, 388; natural enemies of insect pests in, 1, 2, 126, 260, 388; biological control of Lantana in, 33.

Quick-lime, in washes against Melanauster chinensis on Citrus, 270.

Quince, Caliroa limacina on, in W. Australia, 365; Zeuzera pyrina on, in England, 35; pests of, in New York, 246, 249; Phenacaspis prunorum on, in Persia, 16; DDT retarding ripening of, 249.

Quinoline, protecting wood from ter-

mites, 172.

R.

Rabbit Skins, pests imported into U.S.A. in, 275.

Radish, Ceuthorrhynchus assimilis on, in Washington, 76. (See Raphanus spp.)

Ragweed Borer (see Eucosma strenuana).

Railway Waggons, use of compressed air against pests of stored grain in, 320.

rainbowi, Kalotermes (Calotermes).

Rainfall, relation of locusts and grass-hoppers to, 124, 268; relation of other insects to, 24, 158, 174, 175, 187, 283, 300, 306, 307, 374, 375.

Raisins (see Fruit, Dried).

Ramburiella hispanica, in Spain, 268.

rapae, Diaeretus; Pieris. rapax, Hemiberlesia.

Rape, Meligethes aeneus on, in Germany, 97.

Rape Beetle (see Meligethes aeneus).
Raphanus, Ceuthorrhynchus assimilis

on, in Washington, 76.

Raphanus raphanistrum, methods of destroying, against Aphids in Maine, 226.

Raphanus sativus (see Radish).

Raphanus sativus hortensis (Chinese Radish), Crocidolomia binotalis on, in Fiji, 101, 366; erroneously cited as Brassica sativus hortensis, 101, 366.

Raphanus sativus longipinnatus (Daikon Radish), Philaenus leucophthalmus on,

in Hawaii, 274.

Raphia, new weevil on, in Ivory Coast, 14.

raphiae, Leurostethus.

Raspberry, Byturus urbanus on, in Denmark, 7; Taeniocampa gracilis on, in Finland, 42; pests of, in U.S.A., 75, 78, 142, 160, 292.

Raspberry Beetle (see Byturus urbanus).

ratzeburgi, Scolytus.

ratzeburgiana, Zeiraphera.

Ravenala madagascariensis, Rhabdoscelus obscurus on, in Queensland, 1.

recessus, Bracon (Microbracon).

recurvalis, Hymenia.

Redwood, Coast (see Sequoia semper-virens).

Red-legged Earth Mite (see *Halotydeus destructor*).

Red-legged Grasshopper (see *Melanoplus femur-rubrum*).

reflexa, Plecoptera.

religiosa, Mantis.

Resin, in adhesive mixture, 118.

Resin Residue, emulsion of, as adhesive for DDT, 246.

Reticulitermes flavipes, in buildings in Missouri, 49.

Reviews:—Barnes (H. F.), Gall Midges of Economic Importance. Vols. I & II, 251; del Cañizo Gomez (J.), The Potato Beetle, 100; Chamberlin Chamberlin (W. J.), Entomological Nomenclature and Literature, 113; Escherich (K.), The Forest Insects of Central Europe. Vol. V., 69; Ferrière (C.), Défago (G.) & Roos (K.), Lutte contre le doryphore de 1923 à 1943, 327; Gnadinger (C. B.), Pyrethrum Flowers. Supplement, 352; Herce (P.), Análises de insecticidas, 20; Lizer y Trelles (C. A.), Insects and other Enemies of Market Gardens, 133; López Cristóbal (U.), Insectos útiles a la agricultura, McClintock (J. A.) & Fisher (W. B.), Spray Chemicals and Application Equipment, 151; Massee (A. M.), The Pests of Fruits and Hops. 2nd edn., 101; Moritz-Romanova (Z. E.), Berezhkov (R. P.) & Davuidov (P. N.), Pests and Diseases of the cultivated Plants of Western Siberia, 326; Raucourt (M.) & Bégué (H.), Formulaire phyto-pharmaceutique, 66; Silvestri (G.), Compendio di entomologia applicata. Parte speciale, Vol. II, 10; Thompson (W. R.) Ed., A Catalogue of the Parasites and Predators of Insect Trujillo Peluffo (A.), Pests, 96: Insects and other Pests of Agriculture and Agricultural Products in Uruguay, 104: West (T. F.) & Campbell (G. A.), DDT, the Synthetic Insecticide, 327; Some Properties and applications of D.D.T., 348.

Rhabdoscelus obscurus, causes of decreased importance of, on sugar-cane in Queensland and Hawaii, 1, 170; alternative food-plants of, 1.

Rhagoletis cingulata (in U.S.A.), insecticides against, on cherry, **75**, **143**; distribution and association of, with *Prunus emarginata*, **108**.

Rhagoletis fauta (in U.S.A.), sprays against, on cherry, **75**; distribution and association of, with *Prunus emarginata*, **108**.

Rhagoletis pomonella (on apple in U.S.A.), seasonal activities of, 235, 306; insecticides against, 49, 196, 220, 221, 235, 306.

Rhagoletis suavis var. completa, experiment with HCN and puparia of, 151. rhamni, Aphis.

Rhamnus alnifolia, destruction of, against Aphis rhamni in Maine, 226.

Rhamphus senegalensis, sp. n., on salads and cowpeas in Senegal, 206.

Rhaphidopalpa (see Aulacophora).

Rheedia brasiliensis, Conotrachelus on, in Brazil. 6.

Rhizoecus moruliferus, Acropyga associated with, on cacao in Trinidad, 372.

Rhizoglyphus echinopus, on imported lily and hyacinth bulbs in France, 350.

Rhizomyrma (see Acropyga).

Rhizopertha dominica (in stored cereals), in Africa, 63, 328; in Australia, 21, 122, 123, 188, 222, 223, 230; in India, 296, 325; in U.S.A., 177; bionomics and physiology of, 21, 97, 122, 123, 188, 222, 223, 230; effects of competition with other insects on, 97, 98; boring in wood, 328; measures and experiments against, 177, 230, 296, 325.

Rhode Island, Calomycterus setarius in, 105.

Rhodesia, Southern, cotton pests in, 166; Orthezia insignis in, 33; Rhizopertha dominica not recorded in, 63; modification of restrictions on importation of dahlias into Nyasaland from, 62.

Rhodnius prolixus, permeability of cuticle of, by insecticides, etc., 60.

Rhododendron, Heliothrips haemorrhoidalis on, in Hawaii, 277. (See Azalea.) rhodophaga, Neocerata (Dasyneura).

Rhoecocoris sulciventris, DDT sprays against, on Citrus in New South Wales, 383.

Rhopalosiphum prunifoliae (on wheat), experiments with Aphidius granarius against, in Britain, 46; DDT dust against, in New South Wales, 385.

Rhopalosiphum rufomaculatum, DDT aerosol against, on greenhouse chrysanthemums in Maryland, 240.

Rhopalosiphum subterraneum (see Cerosipha).

Rhopobota naevana, sprays against, on cranberry in Washington, 75.

Rhubarb, pests of, in New South Wales, 382, 384.

Rhyacionia buoliana, bionomics and control of, on Pinus radiata in Argentina, 371, 372.

Rhynchaenus rufipes, bionomics and control of, on willow in Connecticut, 237.

Rhynchites aequatus (on plum), bionomics of, in Britain, 272; measures against, in France, 370; on Crataegus and apple, 272.

Rhynchites auratus ferganensis, measures against, on apricot in Central Asia, 49.

Rhynchites bacchus, bionomics and control of, on fruit trees in France, 370. Rhynchites bicolor, probably destroying Megastigmus nigrovariegatus on rose in Illinois, 293.

Rhyssa persuasoria, utilisation of, against Sirex noctilio in New Zealand, 323.

ribesii, Nematus (Pteronus). ribis, Capitophorus (Myzus).

Rice, new weevil on, in Argentina. 171; Chilo loftini on, in California, 375; pests of, in China, 119, 151, 293; Hieroglyphus nigrorepletus on, India, 47; Chilo on, in Spain, 269.

Rice (Stored), 66; Coleoptera in, 101, 120; and products, in baits, 191, 294. Rice Stem Midge (see Pachydiplosis

oryzae).

Rice Weevil (see Calandra oryzae).

Ricine, 53.

Ricinus communis (see Castor).

rigida, Oeme.

Rimu (see Dacrydium cupressinum). Rio de Oro, Schistocerca gregaria in, 265. Rioxa araucariae, rendering Araucaria unsuitable for Aesiotes in Queensland,

Riptortus serripes, measures against, on beans in Queensland, 126.

visbeci, Apanteles; Eurystylus; derces.

riveti, Cyrtorhinus.

Robinia pseudacacia, factors affecting infestation of, by Cyllene robiniae in N. Dakota, 233.

robiniae, Cyllene.

laphygmae, introduced Rogas Hawaii from Texas, 3; bionomics of,

Rogas plecopterae, parasite of Plecoptera veflexa in India, 286.

Rollinia longifolia, Arsenura armida on, in Brazil, 6.

Rosa (see Rose).

vosaceana, Tortrix (Archips).

rosae, Athalia; Crypticerya; Eurytoma;

Macrosiphum; Psila.

Rose, Aonidiella aurantii on, in W. Australia, 364: Crypticerya natalensis on, in Belgian Congo, 82; Incurvaria quadripunctella on, in France, 348; Heliothrips haemorrhoidalis on, Hawaii, 277; new Coccid on, in Iraq, 16; Popillia japonica on, in Nova Scotia, 130; pests of, in U.S.A., 26, 54, 78, 79, 237, 292, 293, 373; importance and destruction of, as winter food-plant of injurious Aphids, 226, 367; susceptibility of species of, to Megastigmus nigrovariegatus, 292; insecticide treatments causing injury to, 54, 374.

Rose Apple (see Eugenia jambos). rossi, Lindingaspis (Chrysomphalus).

Rotenone, against Aphids, 73, 143, 144, stimulating 177, 239, 319, 356; reproduction of Aphids, 143, 144; against Coleoptera, 29, 72, 75, 76, 100, 178, 239, 339, 356, 359, 370; against Empoasca fabae, 239; against Lepidoptera, 78, 216, 305, 313, 337, 339; against Rhagoletis pomonella, 49, 221; against thrips, 24; toxicity of, to honey bees, 55; agents affecting penetration of insect cuticle 60, 87; aerosol of, 339; in dusts, 24, 29, 49, 55, 72, 73, 75, 76, 78, 100, 143, 144, 177, 178, 216, 239, 305, 313, 337, 356, 359; in sprays, 24, 78, 221, 239, 319, 356; formulae containing, 24, 29, 239, 313, 337, 359; and arsenates, 356; and Bordeaux mixture, 239; and cryolite, 29, 356; DDT comand cryolite, 29, 356; DDT compatible with, 157; and nicotine, 177, 356; and pyrethrum, 24, 339; and sulphur, 29, 239, 359; and thiocyanate, 177; materials reducing effect of light on, 50; modification and results of test for, in plants, 218. Derris, Cubé and Timbo.)

rowani, Telenomus (Phanurus).

Royal Poinciana (see Poinciana regia). Rubber, new mite on, in Brazil, 104.

rubens, Epineura. rubi, Anthonomus. rubigalis, Phlyctaenia.

rubriceps, Hamitermes. rubrinervis, Helopeltis bergrothi.

rubrocinctus, Selenothrips.

rubrofasciatus. Piezodorus (see hybneri).

Rudbeckia, thrips on, in greenhouses in Maryland, 240.

rufa, Formica.

ruficollis, Necrobia.

ruficorne, Sinoxylon.

ruficornis, Cerotoma; Magdalis.

ruficrus, Apanteles.

rufimanus, Bruchus.

rufipennis, Polygraphus.

rufipes, Hylastes (Hylurgopinus); Necrobia; Pentatoma; Rhynchaenus (Orchestes).

rufiscutellaris, Glypta.

rufomaculatum, Rhopalosiphum.

ruforum, Achrysocharella.

rufoscapus, Chelonus (Chelonella).

rufovillosum, Xestobium. rugicollis, Plesiocoris.

rugulosus, Scolvtus.

Rumex, Anuraphis tulipae on, in Queensland, 169.

rumicis, Aphis. ruralis, Voria. rusci, Ceroplastes. Russian Thistle (see Salsola).

Russian Union (see Union of Socialist Soviet Republics).

russica, Monolepta.

Rust-red Grain Beetle (see Laemophloeus ferrugineus).

rusticus, Xylotrechus. rutila, Masicera (Erycia).

Ryanex, tests of, against Lepidoptera, 137, 249.

Ryania speciosa, Ryanex (q.v.) a pre-

paration of, 137, 249.

Rye, 158; Hylemyia coarctata on, in Britain, 367; Anomala horticola on, in Denmark, 116; experiments with wireworms and, in Germany, 92; Nysius ericae on, in Russia, 36; Sitodiplosis mosellana on, in Washington, **251.**

Ryegrass (see *Lolium*).

Sabadilla, against Coleoptera, 72, 144; badilla, against Hemiptera, 72, against Hemiptera, 72, 313; 72, 144, 261, against Lepidoptera, 144, 313; against Melanoplus femur-rubrum, 301, 312; action of, on insects, 144, 313; toxicity to insects of alkaloids of, 301, 302; treatments increasing toxicity of, 144, 301, 312; in dusts, 72, 144, 261, 301, 311, 312, 313; carriers for, 144, 301, 311, 312, 313; in sprays, 144, 312, 313; formulae containing, **72, 144, 261, 301, 311, 312, 313**; and sulphur, 144, 312, 313.

sabinusalis, Sylepta.

sabulifera, Anomis (Cosmophila).

saccharalis, Diatraea.

saccharicolus, Diarthrothrips.

saccharina, Lepisma. saccharoni, Thrips.

Safflower (see Carthamus tinctorius).

Safrole, in solid baits for Cydia pomonella, 308, 309.

Sahlbergella singularis, parasite of, on cacao in W. Africa, 296.

sahlbergellae, Euphorus.

St. John's Wort (see Hypericum perforatum).

Saissetia coffeae (hemisphaerica), on orchids and Adiantum in Maryland, 315; on Citrus in New Zealand, 21; sprays against, 21, 315.

Saissetia oleae, in Chile, 206; in Greece, 326; in New Zealand, 21; in U.S.A., 25, 206; on Citrus, 21, 25, 206; on olive, 206, 326; parasites and biological control of, 206, 326; insecticides against, 21, 25; experiments with HCN and races of, 150.

salaminia, Eumaenas (Othreis).

salicis, Stilpnotia.

Salix caprea, pests of, in Finland, 41. Salix pentandra, Rhynchaenus rufipes on, in Connecticut, 237.

sallei, Lygus.

Sallow (see Salix caprea).

Salsola, destruction of, against Eutettix tenellus in California, 374.

Salsola kali var. tenuifolia, summer foodplant of Eutettix tenellus in Idaho, 353.

Salsola pestifer, S. kali tenuifolia recorded as, 353.

samoanus, Kalotermes (Calotermes).

San José Scale (see Quadraspidiotus perniciosus).

sanctae-helenae, Heteronychus.

Sand, for protecting stored cereals from Calandra oryzae, 121.

sanguinipes, Pimpla (Ephialtes).

Sanninoidea (see Aegeria).

Saperda, in elm logs in Connecticut, 236. Saperda carcharias, on Populus tremula in Finland, 41.

Saperda populnea, trees attacked by, in Finland, 41.

Saperda scalaris, trees attacked by, in Finland, 9, 41.

Saperda tridentata (see Eutetrapha). Sapota, White (see Casimiroa edulis).

Sarcophaga kellyi, parasite of Melanoplus mexicanus in Br. Columbia, 227.

Saskatchewan, pests of field crops in, 159, 174, 319.

Sassafras Oil, 58; in bait for Cydia pomonella, 198.

satyriniformis, Melittia.

Sawdust, in baits for locusts, 294, 339; in solid baits for Cydia pomonella, 308.

Sawfly, Gooseberry (see Nematus ribesii).

Sawfly, Pine (see Diprion pini).

Sawfly, Wheat Stem (see Cephus cinctus).

scabra, Plathypena.

scalaris, Saperda.

Scale, California or Citrus Red (see Aonidiella aurantii).

Scale, Fluted (see *Icerya purchasi*). Scale, Olive (see Parlatoria oleae).

Scale, San José (see Quadraspidiotus perniciosus).

Scale Insects, natural enemies of, 63, 81, 82, 145, 187, 190, 206, 234, 238, 274, 361, 362; ants associated with, 33, 46, 238, 372; and plant virus disease, 46, 102; classification and disease, 46, 102; classification and new species of, 16, 82, 112, 205, 256, 329; of Fiji, 388.

Scambus (see Ephialtes):

Scapteriscus acletus, measures against, in seed beds in Florida, 192.

Scapteriscus vicinus, in seed beds in Florida, 192; bionomics and biological control of, in W. Indies, 31, 32, 191; other measures against, 31, 191, 192.

scapularis, Chalepus.

scarabaeoides, Phloeotribus.

Scelio bipartitus, parasite of Gastrimargus musicus in Queensland, 2.

Schinus molle, thrips on, in U.S.A., 24. Schistocerca gregaria (peregrina), in N. and W. Africa, 10, 11, 165, 265, 351; breeding places and migrations of, in Arabia and Iraq, 15, 16; in Kenya and Persia, 352: in Somalia, 294: study on physiology and phases of, 350-351; mass movements of hoppers of, 352; intestinal bacteria of, 10; baits for, 16, 165, 295; other measures and experiments against, 11, 16.

Schizoneura lanigera (see Eriosoma).

schlechtendali, Phyllocoptes.

schoenei, Tetranychus.

schoenobii, Shirakia; Tetrastichus.

Schoenobius bipunctifer (incertellus), bionomics and control of, on rice in China, 152.

Schoenocaulon (see Sabadilla).

Schoenocaulon officinale, preparation of sabadilla insecticides from, 144, 312. schwartzi, Anuraphis.

Sciara, new Nematode infesting, on Convallaria in Denmark, 160.

Sciara inconstans, killed by DDT aerosol in greenhouses in Maryland, 240.

sciarae, Proatractonema.

Sciopus exul, predacious on other insects in hothouses in Holland, 99.

Scirpophaga nivella (in India), method of estimating infestation of sugar-cane by, 47; Bracon greeni reared on, 165. Scirtothrips citri, bionomics and control of, on Citrus in U.S.A., 24, 263.

Scirtothrips longipennis, spray against, on Cinchona in Porto Rico, 218.

Sclerotinia spp., relation of insects to, in fruits, **321, 370.**

Sclerotium rolfsi, 338.

scoliaeformis, Aegeria (Synanthedon). Scolytus mali, bionomics of, on cherry in Holland, 67.

Scolytus multistriatus (in U.S.A.), relation of, to Dutch elm disease, 236; sprays against, in elm logs, 55.

Scolytus ratzeburgi, on birch in Finland,

Scolytus rugulosus, on apple in England, 35; on Sorbus aucuparia in Finland, 41; on peach in Tennessee, 321.

Scolytus scolytus, in dead apple tree in Holland, 67.

Scopula emissaria, Euplectrus gopimohani reared on, in Bengal, 164.

Scorodite (see Ferric Arsenate).

scortealis, Tetralopa.

scrophulariae, Anthrenus.

scrupulosa, Teleonemia.

scutellata, Sturmia

scutellatus, Gonipterus; Monochamus.

Scutellista cyanea, parasite of Saissetia oleae in Greece, 326.

scutigera, Platyedra.

Scutigerella immaculata, measures

against, in Oregon, 318.

Scymnus uncinatus, establishment of, against mealybugs in Hawaii, 274.

Scythropia crataegella, bionomics and control of, on Cotoneaster spp. in Britain, 185.

Seed-corn Maggot (see Hylemyia cilicrura).

segetum, Agrotis.

Selatosomus (see Corymbites).

selecta, Ptychomyia (Bessa).

selenaria, Boarmia (Ascotis)

Selenothrips rubrocinctus, on azalea and Passiflora in Hawaii, 277.

Selinon, 93, 94.

Semanotus ligneus sequoiae, bionomics of, in Sequoia sempervirens in California, 30, 31.

senegalensis, Rhamphus.

senilis, auct., Ceromasia (see Lydella stabulans grisescens).

Sepiomus casuarinae, sp. n., on Casuarina in Annam, 362.

septemfasciata, Nomadacris.

septempunctata, Coccinella. septentrionalis, Croesus.

Septobasidium, infesting Howardia biclavis in Nyasaland, 63.

Sequoia sempervirens, Coleoptera in, in California, 30.

sequoiae, Semanotus ligneus.

seriatus, Psallus.

Serica tristis, on poplar and spruce in Quebec, 173.

sericaria, Ephestia (Hyphantidium) (see

E. elutella). sericeus, Metamasius; Spermophagus

(Euspermophagus). serieventris, Podisus.

serinopa, Nephantis.

serius, Eretmocerus.

serricorne, Lasioderma. serripes, Riptortus.

sertifer, Neodiprion (Diprion, Lophyrus). Sesame Oil, as an activator for pyrethrum in aerosols, 241, 338.

Sesamia vuteria, bionomics Morocco, 257.

Sesamum, Halticid on, in Nyasaland, 63. Sesamum orientale, as trap-crop for Laphygma exigua in Bengal, 164.

INDEX. 471

Sesia (see Aegeria).

Setaria italica, grasshoppers feeding on, in India, 47, 207.

setarius, Calomycterus.

setifacies, Lypha (Didyma).

Setoptus, gen. n., 110.

Setoptus jonèsi, type of genus, 110.

setosa, Chaetophleps.

sexannulatus, Cryptus.

sexdentatus, Ips.

sexta, Protoparce.

seychellensis, Microphanurus.

Shallot, Acrolepia assectella on, in Britain, 70.

shastae, Cryptaspidiotus.

sheldoni, Aceria (Eriophyes).

Shirakia schoenobii (dorsalis), parasite of Schoenobius bipunctifer in China, 152.

shirakii, Cremastus.

Shrews, destroying Gilpinia frutetorum in Connecticut, 237.

Siberia, book on pests of plants in, 326. sidae, Dysdercus.

Sierra Leone, Pseudococcus njalensis on coffee and cacao in, 46.

sierricola, Bixadus.

sigillatus, Gryllodes. signatum, Trypodendron.

signatus, Anthonomus.

Silica, ineffective against Anthonomus pomorum, 274.

Silkworm (see Bombyx mori).

Silpha opaca, bionomics and control of, on beet in France, 10.

Silpha turkestanica, on lucerne Tadzhikistan, **253.**

Silvanus surinamensis (see Oryzaephilus).

similalis, Loxostege.

simonyi, Lygus. simplex, Anabrus; Caedicia; Chilo; Dasus; Lissorhoptrus; Taeniothrips.

Simsim (see Sesamum)

simulans, Psylla (Psyllia).

Sind, Bruchus analis in stored pulses in, 229.

Sinea diadema, predacious on Megastigmus nigrovariegatus in Illinois, 293.

sinensis, Ceroplastes. singaporensis, Apanteles.

singularis, Otiorrhynchus; Sahlbergella.

sinicus, Adoretus.

Sinox, 226. Sinoxylon ruficorne, measures against, in timber in S. Africa, 12; possible parasites of, in Nyasaland, 63.

Sinoxylon transvaalense, measures against, in timber in S. Africa, 12.

Sirex cyaneus, 115.

Sirex juvenous, forms of, in Canadian spruce in Britain, 115.

Sirex noctilio, utilisation of Rhyssa persuasoria against, in New Zealand, 323.

Situation Situation Situation Situation of Situation Situation of Situation on wheat in Sweden and N. America, 36, 251; measures against, 251.

Sitodrepa panicea, in stored ants' pupae in Czechoslovakia, 211.

Sitona, crop rotations ineffective against, in U.S.S.R., 157.

Sitona callosus, bionomics of, on lucerne in Uzbekistan, 156.

Sitona crinitus, bionomics of, on lucerne in Uzbekistan, 156.

Sitona cylindricollis, on Melilotus in N. Dakota, 29, 30; on lucerne in Uzbekistan, 156; bionomics of, 29, **30, 156;** measures against, **30.**

Sitona fronto, on lucerne in Central Asia, **156, 253**; bionomics of, **156.**

Sitona hispidulus, on lucerne California, 111.

bionomics of, Sitona humeralis; on lucerne in Uzbekistan, 156.

Sitona lineellus, bionomics of, on lucerne in Uzbekistan, 156.

Sitona longulus, bionomics of, on lucerne in Uzbekistan, **156.**

Sitophilus (see Calandra).

Sitotroga cerealella (in stored cereals), in India, 296, 325; in Queensland, 388; on field sorghum, 388; effects of competition with other insects on 97, 98; natural reservoirs of, 150; measures and experiments against, 296, 325, 388; Trichogramma reared on eggs of, **314.**

Sloe (see Prunus spinosa).

smithi, Clemora (Lachnosterna).

Smoke Candles, for distributing insecticides, 379.

Smudge Fires, as repellents against insects, 20, 282.

Soap, as an emulsifier, 89, 106, 294; other uses of, in mixed sprays, 18, 37, 49, 53, 54, 118, 125, 199, 229, 237, **274**, **365**, **385**; reducing adhesion of sprays, 220; strawberry injured by sprays containing, 199; types of, 89, 106, 118, 125, 199, 274; formulae containing, 18, 37, 49, 54, 89, 118, 125, 199, 229, 274, 294, 365.

Sodium Antimony Lactophenolate, in sprays against rose pests, 78.

Sodium Arsenate, in preparation of nascent lead and aluminium arsenates,

Sodium Arsenite, in baits, 15, 16, 23, 268, 282, 295; Aphids favoured by, in bait-spray against cotton stainers, 167; sap-stream treatment of trees with, as timber preservative, 56; uses of, to destroy Prunus nigra, 226. Sodium Bicarbonate, effect of,

Tribolium in biscuits, 140.

Sodium Bisulphate, in spray against weeds, 226.

Sodium Carbonate, treatment of sabadilla with, 301.

Sodium Dichromate, sap-stream treatment of trees with mixture containing, as timber preservative, 56.

Sodium Dinitro-o-cresylate (see under

Dinitro-o-cresol).

Sodium Fluoaluminate (see Cryolite). Sodium Fluoride, in baits and bait-spray, 8, 23, 191, 331; uses of, against Camponotus, 27, 28; DDT compatible

with, 157.

Sodium Fluosilicate, in baits, 8, 23, 25, 35, 165, 191, 192, 200, 282, 292; dusting with, 72, 76, 315, 319, 321, 355; spraying with, 315, 356, 383; causing injury to plants, 321, 355, and calcium cyanamide for defoliating cotton, 315; DDT compatible with, 157.

Sodium Hypochlorite, uses of, 142. Sodium Lauryl Sulphate, in dusts, 316; in sprays, **106, 242, 246, 316.**

Sodium Nitrate, decreasing injury by

cryolite to peach, 322.

Sodium Oleyl Sulphate, as a spreader for

sprays, 244.

Soil, effect of types of, on Austroicetes cruciata, 124; apparatus and method of sampling Arthropod fauna in, 372. solana, Empoasca.

solani, Gargaphia; Macrosiphum (Aulacorthum); Phenacoccus. solanifolii, Macrosiphum.

Solanum dulcamara, experiment with Leptinotarsa decemlineata and, 264. Solanum melongena (see Egg-plant).

Solanum nigrum, experiment with Leptinotarsa decemlineata and, 264.

Solanum tuberosum (see Potato).

Solar Heat, use of: against Scolytids in logs, 55; against pests of stored products, 121, 325, 365.

Solidago, Calomycterus setarius on, in Connecticut, 105.

Solidago leavenworthii, pests of, U.S.A., 309.

solitarius, Apanteles; Metaplectrus. solstitiale, Amphimallon.

Somalia, Schistocerca gregaria in, 294.

Soot, dusting with, against Caliroa limacina, 365. Sooty Moulds, associated with Homo-

ptera, 33, 171, 246, 388. Sophocarpin, solution of, against

Tetranychus turkestani, 49. Sophora japonica, Hispid on, in Arizona, 27.

sorbiella, Argyresthia.

Sorbus aucuparia, pests of, in Finland, 41.

sordidus. Cosmopolites.

sorghicola, Contarinia.

Sorghum, thrips on, in Hawaii, 277; pests of, in India, 47, 117, 118, 207; Sesamia vuteria on, in Morocco, 257; pests of, in Queensland, 387; new pests of, in Senegal, 96; Lygus simonyi on, in Uganda, 13; pests of, in USA 180 275 in U.S.A., 180, 375; (stored), pests in and their control, 388; (stalks), as carrier for locust bait, 294, 295.

Sorghum sudanense, thrips on, in Hawaii,

soror, Diabrotica (see D. undecimpunctata).

Soy Bean, pests of, in Porto Rico, 218, 219; pests of, in U.S.A., 27, 240, 244, 317; cyclohexanone causing injury to, 240.

Soybean Flour, as an adhesive for lead arsenate, 199, 220, 248.

Soybean Meal, in baits for Paria canella,

200, 201. Soybean Oil, effect of, in lead-arsenate sprays, 196, 220.

Spaghetti, Calandra oryzae in, in W. Australia, 365.

Spain, Leptinotarsa decemlineata in, 263. 264, 265; locusts and their parasites in, 265, 266, 268; miscellaneous pests in, 266-269; use of Trichogramma against Cydia pomonella in, 288.

Spalangia drosophilae, factor affecting diapause of, parasitising Oscinella frit

in N. America, 295.

Spanioza (see Trioza).

Spatulifimbria castaneiceps, new parasite of, in Ceylon, 47.

speciosus, Anaphothrips.

Speedsprayer, 195, 196. Spermatolonchaea inquilina, possibly predacious on Hylemyia anthracina in Finland, 64.

Spermophagus dorsopictus, sp. n., in imported dried beans in France, 329; considered a synonym of S. subfasciatus, 329.

Spermophagus sericeus, on safflower and Convolvulus in Uzbekistan, 155.

Spermophagus subfasciatus, S. dorsopictus considered a synonym of, 329.

sphenarioides, Colemania.

Sphingomorpha chlorea, attacking Citrus fruits in Nigeria, 14; measures against, 15.

Spicaria prasina, infesting Lepidoptera in Porto Rico and Argentina, 219, 300. Spiders, destroying noxious insects, 24,

Spilochalcis bergi, bionomics of, in Argentina, 300.

152, 379.

Spilonota ocellana, bionomics and control of, on Cotoneaster in Britain, 185; on INDEX.

plum in California, 375; on cherry in Holland, 90.

Spinach, mite on, in Belgium, 161; pests of, in U.S.A., 243, 375; experimentally infected with turnip viruses, 182; DDT causing injury to, 176.

Spinach Beet (see Beet).

spinarum, Athalia (see A. rosae).

Spindle Tree (see Euonymus).

spiniceps, Chirothrips.

spirulosus, Phloeophthorus (Phthorophloeus).

splendidella, Dioryctria.

Spodoptera abyssinia, biochemical differences between phases of, 47.

Spondias, Anastrepha fraterculus experimentally infesting, in Mexico, 138. Sprays, equipment for applying, 141,

176, 195, 346, 356, 379.

Spraysoy A; 199.

Spruce (Picea), Sirex juvencus in imported timber of, in Britain, 115; pests of, in Canada, 129, 173, 263, 378; pests of, in Finland, 38, 39, 64; Neodiprion sertifer on, in Sweden, 131; pests of, in U.S.A., 76, 176, 237; effect of methyl bromide on, as Christmas trees, 316.

Spruce, Engelmann (see Picea engel-

manni).

Spruce, Red (see Picea rubra).

Spruce Beetle, Engelmann (see Dendroctonus engelmanni).

Spruce Budworm (see Harmologa fumiferana).

squamulata, Glyptoscelis.

Squash, Thrips hawaiiensis on, in Hawaii, 277; pests of, in U.S.A., 52, 72, 140, 148, 311, 313, 337, 355; Diabrotica melanocephala transmitting Bacillus tracheiphilus to, 148; DDT retarding development of, 338.

Squash Bug (see Anasa tristis). Squash Vine Borer (see Melittia satyrini-

formis).

Squill, pests of, in greenhouses in Maryland, 240.

stabulans, Lydella.

Stantonia lamprosemae, parasite of Pyralids in Porto Rico, 219.

Starch, as adhesive for insecticides, 319, 334.

Steam, use of, against Tyrophagus lintneri, 78.

steinwedeni, Acaphylla.

Stellaria media, Agalliana ensigera experimentally transmitting Argentine curly-top virus to, 379.

Stenoma decora, on cacao in Brazil, 5. Stephanotis, Pseudococcus citri on, in U.S.A., 79.

Stephegyne diversifolia, Pyralid on, in Burma, 285.

Sterictophora, 317.

Steropleurus perezi, in Spain, 268.

Sthenarus, new parasite of, on cotton and Albizzia in Uganda, 296.

473

sticticalis, Loxostege.

sticticraspis, Proceras (Argyria).

Stictocephala, 317.

Stictococcus, Cryptoblabes gnidiella predacious on, in Belgian Congo, 81.

stigma, Chilocorus.

stigmaticus, Coeloides.

Stigmatium ventrale, predacious on Aesiotes notabilis in Queensland, 260. Stilpnotia salicis, introduced parasite of, on poplar in Canada, 129.

Stinoplus militaris, parasite of Scolytids in Finland, 39, 41.

Stock (see Matthiola).

stokesi, Agrothereutes (Gambrus).

Strawberry, pests of, in Britain, 101;
Anthonomus rubi on, in Denmark, 7;
Philaenus leucophthalmus on, in
Hawaii, 274; virus diseases of, in
Holland, 90; Anthonomus signatus
on, in New Brunswick, 319; Heteronychus sanctae-helenae on, in New South
Wales, 382; Odontria zealandica on,
in New Zealand, 20; pests of, in
U.S.A., 78, 142, 160, 181, 199, 200,
292; Monolepta russica on, in Uzbekistan, 155; Capitophorus fragariae
transmitting virus diseases of, 90.

Strawberry Leaf-roller (see Ancylis:

comptana).

Strawberry Weevil (see Anthonomus signatus).

strenua, Caedicia.

strenuana, Eucosma (Epiblema).

strigatus, Eumerus.

strobi, Cinara; Dasyneura (Perrisia); Pissodes.

strobicola, Eutelus.

strobilella, Cydia (Laspeyresia).

strobilellae, Ephialtes.

Strontium Arsenate, against Leptinotarsa decemlineata, 83.

Strumeta (see Dacus).

Sturmia, utilisation of, against Gilpinia hercyniae in Canada, 130, 173.

Sturmia cubaecola, parasite of Lamprosema indicata in Porto Rico, 219.

Sturmia inconspicua, parasite of Neodiprion sertifer in Finland, 39.

Sturmia scutellata, parasite of Lymantria dispar in U.S.A., 193.

Styphelia, thrips on, in Hawaii, 277.

suavella, Eurhodope.

suavis, Macrodactylus; Rhagoletis.

subcrinita, Epitrix.

subericinella, Aristotelia.

subfasciatus, Spermophagus. subpusilla, Agromyza.

subspinosus, Macrodactylus.

subterranea, Cerosipha (Rhopalosiphum); Feltia.

Subterranean Clover (see Trifolium subterraneum).

subtilis, Cosmia.

Sucrose (see Sugar).

Sudan, Anglo-Egyptian, Anthrenus coloratus in, **59**; cotton pests in, **166**. Sudan Bollworm (see Diparopsis castanea).

Sudan Grass (see Sorghum sudanense).
Sugar, in baits, 198, 200, 201, 283, 376;
in bait-sprays for fruit-flies, 133, 385;
unsatisfactory in bait-spray for Pegomyia hyoscyami, 331; in sprays against thrips, 24, 25, 78, 240, 243, 338, 383, 384, 385; tests and uses of, as food for insects, 138, 143, 215, 308, 325, 328, 368.

Sugar-cane, pests of, in Australia, 1, 382; Argyria in, in China, 121; pests of, in Hawaii, 170, 275; pests of, in India, 47, 187, 188, 207, 297; Chilo loftini on, in Mexico, 375; thrips on, in New Caledonia, 275; pests of, in U.S.A., 177, 375; pests of, in W. Indies, 31, 64, 159, 320; Aphids and mosaic of, 297; Phytomonas rubrilineans favouring Rhabdoscelus obscurus in, 1; variety of, resistant to R. obscurus, 170; stalks of, as bait for Heteronychus sanctae-helenae, 382.

Sugar-cane Borer (see Diatraea sacchar-alis).

Sugar-cane Mite, West Indian (see Tarsonemus bancrofti).

sulcatula, Anomala.

sulcatus, Otiorrhynchus (Brachyrrhinus). sulci, Syngenaspis (see S. parlatoriae). sulciventris, Rhoecocoris.

Sulphite Lye, in DDT sprays, 34.

Sulphur, against Capsids, 135, 180; against Coccids, 25; against Empo-asca fabae, 244, 337, 338; against Epitrix tuberis, 358, 359; against mites, 25, 95, 100, 126, 127, 164, 384; increasing infestation of apple Paratetranychus pilosus, against Paratrioza cockerelli, 29, 358; against thrips, 25, 338; toxicity of, to honey bees, 55; method of burning, against Odontria zealandica, 20; dusting with, 25, 95, 100, 126, 135, 164, 180, 244, 317, 337, 338, 358; in mixed dusts, 29, 78, 126, 127, 135, 137, 138, 143, 144, 179, 180, 195, 197, 224, 239, 244, 247, 261, 267, 303, 312, 313, 317, 318, 345, 358, 359; in mixed sprays, 29, 52, 106, 138, 198, 202, 203, 220, 221, 224, 248, 318, 322, 384; fused on bentonite, 200; effect of, on DDT, 106, 157, 245; forms of, 100, 143, 197, 248, 384; formulae containing, 29, 52, 78, 126, 127, 135, 138, 143, 144, 164, 179, 195, 197, 198, 221, 224, 239, 247, 267, 270, 312, 313, 318, 322, 345, 358, 359, 384.

Sulphur Dioxide, fumigation with, 77, 101.

sulphureus, Systoechus.

Sulphuric Acid, spraying with, against weeds, 226.

Superphosphates (manuring with), reducing infestation of jute by Apion, 117; ineffective against Hylemyia coarctata, 217; increasing cryolite injury to peach, 322.

superstitiosus, Dysdercus. suranus, Microphanurus.

Surinam Cherry (see Eugenia uniflora). surinamensis, Oryzaephilus (Silvanus).

swainei, Neodiprion.

Sweden, forest pests in, 40, 130, 131; orchard pests in, 7, 37; Cecidomyiids on wheat and poppy in, 36, 59; natural enemies of Neodiprion sertifer in, 130.

Swedes, Phyllotreta spp. on, in Denmark and Britain, 7, 114; Polia oleracea on, in Finland, 100; Eurydema oleraceum on, in Germany, 89; time of sowing of, in relation to Heteronychus sanctae-helenae in New South Wales, 382; Ceuthorthynchus assimilis on, in Washington, 76.

Sweet Clover (see Melilotus).

Sweet Clover Weevil (see Sitona cylindricollis).

Sweet Gum (see Liquidambar styraciflua). Sweet Pea (see Lathyrus odoratus).

Sweet Potato, pests of, in Barbados, 159; Herse cingulata on, in Florida, 146; Dendrothripoides ipomeae on, in Hawaii, 274; Heteronychus sanctaehelenae on, in New South Wales, 382; (tubers), as bait for H. sanctaehelenae, 382.

swezeyi, Anaphothrips.

Swietenia mahagoni (see Mahogany). Switzerland, pests of vegetables in, 43, 214, 215, 216, 327.

Sword Beans (see Canavalia gladiata).

sycophanta, Calosoma.

Sylepta derogata (Cotton Leaf-roller), in China, 120, 293; in Belgian Congo, 81, 82; in Tanganyika, 82; experiments with varieties of cotton and, 120; on Hibiscus esculentus, 82; parasites of, 81, 82; sprays against, 293.

Sylepta sabinusalis, new parasite of, on Boehmeria nivea in Belgian Congo, 81. syleptae, Eurytoma; Microgaster.

Syllitus araucariae, on Araucaria cunninghami in Queensland, 260.

Synanthedon (see Aegeria).

Syngenaspis parlatoriae (sulci), 235. Syntone, 239.

Syrphus luniger, DDT toxic to, 45.

Syrup, in baits for Cydia pomonella, 308, 309.

Systates, on coffee in Belgian Congo, 153. Systates variabilis, on Cinchona in Belgian Congo, 323.

Systena blanda, on potato in Nebraska, 28.

Systoechus spp., parasites of Dociostaurus maroccanus in Spain, 265.

T.

tabaci, Thrips. taedae, Cinara.

Taeniocampa gracilis, bionomics of, on raspberry, etc., in Finland, 42.

Taeniothrips alliorum, on onion in Hawaii, 277.

Taeniothrips atratus, on beans in Finland, 37.

Taeniothrips inconsequens, bionomics and control of, on fruit trees in U.S.A., 24, 143.

Taeniothrips nigricornis, measures against, on beans in Australia, 125, 383; bionomics of, 125.

Taeniothrips simplex (on gladiolus), in Hawaii, 277; insecticides against, in New South Wales, 385.

taiganus, Urocerus gigas.

Talc, as a carrier for dusts, 50, 89, 137, 138, 143, 149, 169, 238, 258, 304, 312, 313, 316, 317, 333, 386.

Talinum, Halticus citri on, in Bermuda,

tamarindus, Drosichiella (Monophlebus). Tamarix, Chionaspis etrusca on, in California, 109.

Tanganyika Territory, cotton pests in, 82, 167; locusts in, 280; pests of timber in, 103; identity of parasite of Sylepta derogata in, 82.

Tanymecus palliatus, survival of, in crop rotations in U.S.S.R., 157.

tapetzella, Trichophaga.

Taphrina deformans, combined spray against Coccid and, on peach in Kashmir, 293, 294.

taprobanes, Autoplectrus.

Tar, in burning mixture against fruit-

piercing moths, 282.

Tar Distillates, as dormant sprays against pests of fruit trees and Cotoneaster, 90, 93, 116, 184, 349; as summer sprays against Aphids, 7; experiments with, against Xylotrechus quadripes, 187; causing injury to trees, 7; formulae for, 7, 93, 184, 187.

Taraxacum kok-saghyz, Nysius ericae on, in Russia, 35, 36.

Taraxacum megalorrhizon, Nysius ericae on, in Russia, 36.

Taraxacum officinale, Aphids transmitting mosaic of, to lettuce in Britain, 70.

Tarnished Plant Bug (see Lygus oblineatus).

Taro (see Colocasia).

INDEX.

Tarsonemus bancrofti, on sugar-cane in Florida, 177.

Tarsonemus latus (see Hemitarsonemus). Tarsonemus pallidus, on Begonia in Holland, 95; characters of, 95.

Tarsonemus translucens, Green (see

Hemitarsonemus latus).

Tartar Emetic, in bait-sprays against fruit-flies, 133, 276, 385; preventing ovarian development in Dacus cucurbitae, 308; in sprays against thrips, 24, 25, 78, 240, 243, 338, 383, 384, 385; resistance of Scirtothrips citri to, 25; against Lepidopterous larvae, 78; ineffective against Pantomorus godmani, 383; and sugar, 24, 25, 78, 133, 240, 243, 338, 383, 384, 385; formulae containing, 24, 25, 78, 243, 276, 383, 385.

Tasmania, Caliroa limacina in, 365.

taylori, Aridelus.

Tea, pests of, in Ceylon, 71, 211, 324, 325, 366; pests of, in Nyasaland, 58, 62.

Tea Mite, Yellow (see Hemitarsonemus latus).

Tea-seed Oil, in adhesive mixture for trapping *Phyllotreta*, **118**.

Teak, experiments with Xylotrechus quadripes and, in Mysore, 186; risk to, of liberating Teleonemia scrupulosa in India, 287.

Tecoma, Orthezia insignis on, in S. Rhodesia, 33.

Tectona grandis (see Teak).

tectus, Ptinus.

telarius, Tetranychus.

Telegraph Poles, Camponotus spp. boring in, in Oregon, 27.

Telenomus beneficiens, parasite of Schoenobius bipunctifer in China, 152. Telenomus dignus, parasite of Schoeno-

bius bipunctifer in China, 152.
Telenomus numitor, possible parasite of

Pentatomid on Aleurites in Nyasaland, 63.

Telbrowus rowani parasite of Schoene-

Telenomus rowani, parasite of Schoenobius bipunctifer in China, 152.

Teleonemia scrupulosa (attacking Lantana), risk to teak of liberating, in India, 287; utilisation of, in Queensland, 33; suggested introduction of, into S. Rhodesia, 33.

Temnochila virescens, associated with Cerambycids on Sequoia in California,

Temnosternus imbilensis, on Araucaria cunninghami in Queensland, 260.

Temperature, relation of locusts and grasshoppers to, 124, 180, 340, 350, 351; relation of other insects to, 2, 21, 22, 42, 45, 48, 54, 121, 122, 123, 133, 168, 174, 175, 188, 189, 212, 213, 214, 215, 222, 223, 237, 258, 259, 265, 283, 284, 306, 321, 335, 353, 369; effect of, on transpiration through insect cuticle, 60, 61; effect of, on insecticidal action of pyrethrum, 376, 377.

tenax, Melanaspis.

tenebricosa, Capnodis.

Tenebrio spp., experiments on foods for, reared for feeding fowls, 328.

Tenebrionids, characters of species of, infesting stored products, 329.

tenebrionis, Capnodis.

Tenebroides mauritanicus, probably predacious on Lasioderma serricorne in Nyasaland, 62; measures and experiments against, in stored cereals in U.S.A. and Punjab, 177, 296, 342.

tenellus, Eutettix.

Tennessee, peach pests in, 321, 322; miscellaneous pests in, 144, 178, 340; hosts of Macrocentrus ancylivorus in, 322.

tenthredinis, Mesoleius. tenuicornis, Habrocytus.

Tenuipalpus, scope of, 104.

Tenuipalpus heveae, sp. n., on rubber in Brazil, 104.

Tenuipalpus palmatus, type of genus, 104.

Tenuipalpus pseudocuneatus, causing "lepra explosiva" of orange in Argentina, 298.

tenuis, Cyrtopeltis.

Tephrina disputaria, parasite of, on Acacia in India, 285.

Tephrosia, Icerya on, in Belgian Congo, 82.

Tephrosia candida, flowers of, for feeding Neoplectrus maculatus, 324.

Tephrosia toxicaria, toxicity of, to Epilachna, 303.

Tephrosia virginiana, toxicity of, to Epilachna, 303.

Tephrosia vogeli, Etiella zinckenella on, in Belgian Congo, 81; insecticidal action and constituents of, 304.

Terastia meticulosalis, new parasite of, on Erythrina in Belgian Congo, 81. terastiae, Bracon (Microbracon).

Terminalia (Myrobalans), Anthrenus psp. imported into Britain in, 59.

Terminalia catappa, Trypetid on, in Mexico, 137.

terminifera, Chortoicetes.

Termites, attacking living plants, 167, 191, 211, 366; attacking timber, 32, 49, 103, 116, 169, 170, 172, 345, 366, 388; timbers resistant to, 366; damaging textiles and books, 291; measures and experiments against, 32, 116, 170, 172, 191, 291, 345, 380; influence of reproductive forms on control of, 169, 170.

Termites, Harvester (see Hodotermes). terricola, Pristonychus (Laemostenus).

tessmanni. Mesohomotoma.

testudinea, Hoplocampa.

testulalis, Maruca.

Tetrachlorphenol, protecting wood from termites, 172.

Tetraleurodes abutilonea, on egg-plant in California, 109.

Tetralopa scortealis, bionomics and control of, in U.S.A., 304.

Tetramethylthiuram Disulphide, sprays against insects, 236, 273.

Tetranychus, on cotton in China, 293; life-cycle of, on greenhouse plants in Massachusetts, 54; sprays against, 54, 293.

Tetranychus atlanticus, food-plants of,

in California, **374.**

Tetranychus bimaculatus, on Cinchona in Belgian Congò, 323; on greenhouse plants in U.S.A., 78, 240; measures against, 78; not affected by DDT aerosols, 240.

Tetranychus pacificus (on apple), in Br. Columbia, 175; in Washington, 73, 75; sprays against, 73, 175; insecticides favouring infestation by, 75, 175.

Tetranychus schoenei, DDT sprays increasing infestation of apple by, in

Virginia, 106.

Tetranychus telarius, on leguminous plants in Australia, 126, 384; on hops and tomato in Britain, 100, 273; dusts and sprays against, 100, 126; DDT ineffective against, 384.

Tetranychus turkestani, sprays against, on cotton in Central Asia, 49.

Tetranychus urticae (see T. telarius).

Tetrastichus carpatus, parasite of Tineola biselliella in U.S.A., 194.

Tetrastichus inunctus, parasite of Dasyneura strobi in Finland, 39.

Tetrastichus isis, parasite of Oeketicus kirbyi in Argentina, 300.

Tetrastichus oeceticola, parasite of Oeketicus kirbyi in Argentina, 300.

Tetrastichus platensis, parasite of Oeketicus kirbyi in Argentina, 300. Tetrastichus pseudoeceticola, bionomics of, in Argentina, 300.

INDEX.

Tetrastichus schoenobii, parasite of Schoenobius bipunctifer in China, 152. texanus, Chelonus.

Texas, cereal pests in, 3, 375; cotton pests in, 178, 247, 345; Fundella argentina in, 104; parasites of Noctuids in, and their introduction into

Hawaii, 2, 3.

Textiles, insects damaging, 95, 127, 291. textor, Lamia.

Thalessa curvipes, parasite of Necydalis major in Finland, 41.

Thannonoma ochrifascia, on Juniperus bermudiana in Bermuda, 190, 191; spray against, 191.

Thaumetopoea pityocampa, on Pinus pinaster in France, 371.

Thea vigintiduopunctata, experiment with Perilitus coccinellae and, 114.

Theocolax formiciformis, associated with Anobium punctatum in New Zealand, 211.

Theresia claripalpis (parasite of Diatraea saccharalis), imported into Porto Rico from Brazil, 179, 219; failure to establish, in Barbados, 159.

Thermobia domestica, testing of attractiveness of fabrics and paper to, in

U.S.A., 127.

Thersilochus, status of, 252; cf. also

p. 462 (note).

Thiocyanates, unsatisfactory against Aphids, 177, 178, 356; doubtful effectiveness of, against Lepidoptera and Epilachna varivestis, 199, 316, 317; against Jassids, 247, 248, 313, 374; against Pseudococcus citri, 79; against Tetranychus, 54; in dusts, 177, 313, 316, 317, 356; in sprays, 54, 79, 199, 247, 248; vaporised spray of oil containing pyrethrum extract and, 374; and other insecticides, 177, 247, 248, 316, 317, 356; causing injury to plants, 54, 79, 199; uses of, to destroy Prunus nigra, 226.

thoracica, Ealincola (Lycidocoris, Pan-

tilioforma).

Thosea cervina, new parasites of, in Ceylon, 47.

thoseae, Metaplectrus.

Thrips angusticeps, bionomics of, on lettuce in Finland, 37.

Thrips flavus, on Chrysanthemum cinerariaefolium in Nyasaland, 63.

Thrips hawaiiensis, on squash and lily in Hawaii, 277.

Thrips lini, insecticides against, on flax in Holland, 67.

Thrips nigropilosus, food-plants of, in Hawaii, 277; DDT aerosol against,

on greenhouse chrysanthemums in Maryland, 240.

Thrips saccharoni, on maize in Hawaii, 276.

Thrips tabaci, in Hawaii, 276, 277; bionomics of, in India, 118; in New Caledonia, 275; in New South Wales, 384; in Nyasaland, 63; in U.S.A., 53, 239, 240, 243; on onion, 53, 118, 240, 243, 275, 277, 384; on other plants, 63, 118, 240, 276; measures and experiments against, 53, 118, 239, 240, 243, 384.

Thrips, Onion (see Thrips tabaci).

Thrips, Pear (see Taeniothrips inconsequens).

Thrips, Tobacco (see Frankliniella fusca).

Thuja, Coccid on, in U.S.A., 200.

Thuja plicata, Geometrid on, in Br. Columbia, 378.

Thyridanthrax fenestratus, parasite of Dociostaurus maroccanus in Spain, 265.

Thysanoptera, relation of, to plant diseases, 206; natural enemies of, 19, 37, 218; predacious species of, 37, 218; new species of, 275; of Finland, 64.

Thysanus niger, 219.

Til (see Sesamum orientale).

tiliaria, Erannis.

Timber, Camponotus spp. in, 27; Coleoptera in, 11, 12, 13, 103, 116, 211, 333; species of, resistant to Anobium punctatum, 211; termites in, 32, 49, 103, 116, 169, 170, 172, 345, 366; Urocerus spp. in, 115; treatment of, with preservatives against pests, 32, 55, 56, 116, 172.

Timbo, dusting with, against insects, 84, 231; mixed with maize against Calandra oryzae, 334; rotenone contents of, 231, 334. (See Lonchocarpus.)

Timothy Grass (see *Phleum pratense*). Tinea ditella, genitalia of, 128. Tinea insectella, genitalia of, 128.

Tinea pellionella, parasite of, 315.

Tineola biselliella, parasite of, in stored wool in U.S.A., 194; natural sources

of, **150.**Tiphia popilliavora, establishment of, against Popillia japonica in Connec-

ticut, 234. Tiphia vernalis, establishment of, against Popillia japonica in Connecticut, 234.

Tipula paludosa, plants attacked by, in Denmark, 116; survival of, in crop rotations in U.S.S.R., 157.

tipuliformis, Aegeria (Sesia).

Tirathaba trichogramma (see Coleoneura). Tlascala finitella, bionomics and control of, on cranberry in Massachusetts, 51.

Toads, destroying Gonipterus scutellatus,

102. (See Bufo marinus.)

Tobacco, 380; pests and virus diseases of, in Africa, 62, 70, 206; insects associated with, in Argentina, 299; Gnorimoschema operculella in, in Bermuda, 301; Tipula paludosa on, in Denmark, 116; pests of, in U.S.A., 129, 178, 183, 235, 335, 347, 356; Leptinotarsa decembineata not feeding on, 264.

Tobacco (Stored), pests in, and their control, 62, 338; insects damaging

drying leaves of, 116, 207.

Tobacco (as an insecticide), dusting with, 152; greenhouse fumigation with, 78; spraying with, 293; other uses of, against pests, 77, 78; as a carrier for dusts, 137, 236, 356, 383; doubtful value of stems of, against Schoenobius bipunctifer, 153; Lasioderma serricorne damaging smoke spirals containing, 101.

Tobacco Aphis (see Myzus persicae).

Flea-beetle Tobacco (see Epitrix hirtipennis).

Tobacco Hornworms (see Protoparce). Tobacco Thrips (see Frankliniella

fusca).

Togoland, British, Pseudococcus njalensis and associated ants in, 46.

Togoland, French (see Africa, French West).

tokionis, Agrotis.

Tomato, 380; pests of, in Australia, 23, 288, 382; Gnorimoschema operculella on, in Bermuda, 301; pests of, in Brazil, 171, 172; Tetranychus telarius on, in Britain, 100; mite causing leaf-curl of, in Ceylon, 188; pests of, in Finland, 99, 100; Leptinotarsa decemblineata on, in France, 10; pests of, in Hawaii, 149, 276; Achaea attacking fruit of, in Nigeria, 15: as reservoir of Bemisia and crinkle virus of tobacco in Nyasaland, 206; L. decembineata on, in Spain, 264; mole-crickets damaging, in Trinidad, 192; pests of, in U.S.A., 109, 177, 181, 204, 240, 310, 354, 374; experiments with Ceratitis capitata and, 132; insecticides causing injury to, 100, 176, 240, 354; (fruits), fumigation of, against fruit-flies, 190; as baits for Othreis, 188.

Tomato Fruitworm (see Heliothis armi-

gera),

tomentosus, auct., Byturus (see B.

Tonga Beans (see Dolichos lablab).

Torocampus eques, parasite of Neodiprion sertifer in Finland, 9, 39.

Tortrix obsoletana, 304.

Tortrix parallela, flooding against, on cranberry in Massachusetts, 51.

Tortrix podana, on cherry in Holland, 90. Tortrix rosaceana, measures against, on greenhouse rose in Ohio, 78; parasite of, in Quebec, 263.

Torymus spp., characters of, parasitising insects in spruce cones in Finland, 38.

Totara (see Podocarpus totara).

Toxoptera aurantii, fungus infesting, in Argentina, 299; not transmitting swollen-shoot of cacao in Gold Coast, 102.

Toxoptera graminum, fungi infesting, in Argentina and Uruguay, 299, 300; outbreak of, on cereals in U.S.A., 375; varieties of cereals resistant to, 375, 376.

translucens, Green, Tarsonemus (Acarus)

(see Hemitarsonemus latus).

transvaalense, Sinoxylon. transvaalensis, Apanteles.

Traps, for Pegomyia, 331; uses of DDT in, for Popillia japonica, 245; mobile types of, for Eurygaster and Phyllotreta, 48, 118. (See Light-traps.)

Travancore, miscellaneous pests in, 283,

287, 288,

Treehopper, Buffalo (see Ceresa bubalus). Trees, methods of injecting chemicals into, for timber preservation, 55, 56. tremae, Icerya.

Trenches, to trap pests of stored grain,

Trephognathus apii, gen. et sp. n., on

celery in Senegal, 189.
Trialeurodes, DDT spray against, in greenhouses in Brazil, 334; Sciopus exul predacious on, in Holland, 99.

Trialeurodes vaporariorum, on beans in Hawaii, 275; on greenhouse vegetables in Maryland, 240; uses of DDT against, 240, 275.

Triaspis, introduced into U.S.A. against Listroderes obliquus, 179.

Tribolium, methyl bromide against, in banana flour in Morocco, 65.

Tribolium castaneum, in dried bananas in France, 363; (in stored cereals), in Fiji, 101; in India, 287, 296, 325; in Victoria, 230; measures and experiments against, 101, 296, 325, 342, 343.

Tribolium confusum, factors affecting retarded development of, on biscuits, 139: method of determining amounts of HCN in fumigated, 150.

1,1,2-Trichlorethane, tests with, as fumigant against Tenebroides mauri-

tanicus, 342.

Trichlorethylene, fumigation of bulk with mixture containing, wheat against pests, 230; in DDT aerosol formula, 242.

1 - Trichlor - 2 - o - chlorphenyl - 2 - p chlorphenylethane (o,p'-DDT), in technical DDT, 255, 360; toxicity of, to insects, 255, 360.

1-Trichlor-2,2-bis (parachlorphenyl)-

ethane (see DDT).

Trichogramma, technique of rearing, on eggs of Gnorimoschema operculella, 314.

Trichogramma japonicum, parasite of Schoenobius bipunctifer in China, 152.

Trichogramma minutum, parasite Acleris variana in Br. Columbia, 379: utilisation of: against sugar-cane borers in India, 207; against Cydia pomonella in Spain, 288; against Diatraea saccharalis in W. Indies, 31, 159, 320.

Trichogramma pretiosum, experiments with, against Cydia pomonella in

Spain, 288.

trichogramma, Coleoneura (Tirathaba).

Trichomalus fasciatus, parasite Ceuthorrhynchus assimilis in Washington, 72, 76.

Trichophaga tapetzella, in rabbit skin imported into U.S.A. from Hawaii, 275.

Trichopoda pennipes, establishment of, against Anasa tristis in Washington,

Trichospilus pupivora, use of, against Nephantis serinopa in Ceylon, 188.

Trichosteresis, possibly parasitic on Anthonomus pomorum in Holland, 94. Triclistus fulvipes, parasite of Tetralopha

scortealis in Virginia, 305.

tridentata, Eutetrapha (Saperda). Trifolium (see Clover).

Trifolium procumbens, effect of Halotydeus destructor on yield of, in W. Australia, 163.

Trifolium subterraneum, pests of, in Australia, 162, 163, 324.

trigonum, Eutrombidium.

trilobitiformis, Pseudaonidia. trimaculata, Hartigia (Adirus).

Trinidad, miscellaneous insects in, 57, 191, 219, 372.

triodiae, Eutermes.

Trionymus, on Amaryllis in California,

Trionymus lounsburyi, 109.

Trioza apicalis, on carrot in Denmark, 116.

Trioza erytreae, on Citrus in Nyasaland, 63.

Triphleps (see Orius).

Tripterygium forrestii, dust prepared from, against Coleoptera, 344.

Tripterygium wilfordi, insecticides prepared from, against Phaedon brassicae, 120.

tristis, Anasa; Lachnosterna; Serica. tritici, Contarinia; Euxoa (Agrotis); Harmolita; Margarodes.

Tritolestes, new subgenus of Cryptochetum, 82.

Triton, 314.

Triton N.E. 100, 245.

Triton X-100, **244.**

Triumfetta rhomboidea, jute Buprestid on, in Bengal, 164.

Trochilium apiforme, on Populus tremula in Finland, 41.

Trogoderma granarium, measures and experiments against, in stored cereals in India, 296, 297, 325.

Troilus luridus, bionomics of, in Belgium,

213.

Tropaeolum, new Agromyzid on, in U.S.A., 57.

truncatella, Litomastix (Copidosoma). tryoni, Dacus (Strumeta) ferrugineus.

Trypodendron signatum, on birch in Finland, 41.

Trypophloeus (see Cryphalus).

Tsuga, Acleris variana on, Br. in Columbia, 320.

Tsuga heterophylla, pests of, in Br. Columbia, 377, 378.

Tsuga mertensiana, pests of, in Columbia, 378.

tsugae, Neodiprion.

tuberculatus, Eumerus; Lepidosaphes (see L. machili).

tuberis, Epitrix.

Tulip, 367; tolerant to methyl bromide fumigation, 110.

tulipae, Aceria (Eriophyes); Anuraphis.

Tung (see Aleurites).

Turf, Odontria zealandica in, in New Zealand, 20; injured by oil emulsions against mole-crickets in Trinidad, 191; Coleoptera in, in U.S.A., 105, 237, 374.

turkestani, Tetranychus (Eotetranychus).

turkestanica, Silpha (Aclypea). Turkey, pests of wheat in, 48, 256.

turneri, Platyspathius.

Turnip, Athalia rosae on, in Denmark, Gryllotalpa on, in Egypt, 19; Cidaria fluctuata on, in Finland, 38; pests of, in U.S.A., 76, 182, 375; Aphids and virus diseases of, 182; (chopped), in bait for Listroderes obliquus, 23.

Tychius, on lucerne in Tadzhikistan, 253.

Tychius femoralis, 253.

Tyloderma fragariae, measures against, on strawberry in Kentucky, 292.

Typhlocyba pomaria, insecticides against, on apple in U.S.A., 50, 52.

Tyroglyphids, methods of estimating populations of, in stored products, 98. Tyroglyphus, species of Tyrophagus recorded as, 161.

Tyroglyphus farinae, products infested by, 98.

Tyrophagus, measures against, damaging greenhouse spinach in Belgium, 161.

Tyrophagus dimidiatus, 161.

Tyrophagus lintneri, bionomics and control of, on mushrooms in N. America, 76-78.

Tyrophagus longior, 161. Tyrophagus putrescentiae, 161.

U.

ubacahy, Conotrachelus.

Uganda, Antestia spp. on coffee, etc., in, 277-279; cotton pests in, 13, 166, 296: Capsids on other plants in, 13, 296; Lepidosaphes gloveri in, 82; pests of timber in, 103; beneficial insects and biological control in, 13, 278, 279, 296.

uhleri, Aceratagallia.

Ulex europaeus, liberation of Apion ulicis against, in New Zealand, 323. ulicis, Apion.

ulmi, Lepidosaphes (Mytilococcus).

ulmi, auct., Metatetranychus (see Paratetranychus pilosus).

Ultra-violet Rays, effect of, on insecticides, 235.

Unaspis citri, spray against, on Citrus in Queensland, 127.

uncinatus, Scymnus.

undalis, Hellula.

Diotimana undulata, (Diotima);

Phyllotreta.

Union of Socialist Soviet Republics, pests of field crops in, 35, 48, 131, 157, 252; pests of fruit trees in, 131, 374; use of bacteria and fungi against insects in, 131. (See Siberia and Asia, Central.)

unionalis, Margaronia (Glyphodes).

unipuncta, Cirphis.

United Provinces (India), Eriosoma lanigerum on apple in, 228; Chlorops causing short circuits on high tension cables in, 286; Hieroglyphus nigro-repletus in, 47; sugar-cane pests in, 187, 188; pests of stored cereals and pulses in, 325; beneficial insects in, 228, 285, 286.

United States of America, cereal pests in, 136, 177, 192, 357, 375; cotton pests in, 315; pests of forest and shade trees in, 115, 176, 193, 200, 236; orchard pests in, 107, 110, 141, 142, 176; pests of vegetable and root crops in, 57, 100, 133, 177, 178, 264, 267; pests of stored cereals in, 177; Lepismids and cockroaches in, 127; Popillia japonica in, 51, 107, 141, 145, 176, 194, 210, 233, 234, 244, 245, 304, 341; natural enemies and diseases of insects and their utilisation in, 104, 141, 142, 177, 179, 193, 194, 210, 343; pests intercepted in quarantine in, 183, 277; pests in imported rabbit skins in, 274; plant pest legislation in, 112, 291, 292; list of popular names of insects in, 352. (See also under the various States.)

urbana, Ctenolepisma. urbanus, Byturus.

Urocerus californicus, in imported timber in Britain, 115; in N. America, 115. Urocerus gigas taiganus, in Britain, 115. uroxys, Apanteles.

urozonus, Eupelmus.

urticae, Tetranychus (see T. telarius).

Uruguay, book on pests in, 104; natural enemies of Listroderes in, 104, 299; entomogenous fungi in, 299, 300; beneficial insects imported into U.S.A. from, 179.

Utah, Cydia molesta in, 373; lucerne pests in, 260, 261.

utilis, Allotropa. uzeli, Gynaikothrips.

V.

vaccinii, Mineola.

Vacuum Fumigation, 65, 66, 180, 363. Valanga irregularis, bait for, on sorghum in Australia, 387.

Valeric Acid, in baits for Cydia pomo-

nella, 308.

validirostris, Pissodes. Vanda (see Orchids).

vanduzeei, Xerophloea.

vaporariorum, Trialeurodes.

varia, Codophila.

variabilis, Erythroneura; Hypera (Phytonomus); Systates.

variana, Acleris (Peronea).

varians, Cyrtopeltis (see Engytatus geniculatus).

variegata, Gilpinia.

variegatus, Zonocerus.

varinervis, Antholcus.

variolosa, Poecilonota.

varivestis, Epilachna.

Vasates destructor, on tomato in Colorado,

Vasates eurynotus, on carrot and parsley in California, 374.

vayssieriellus, Pseudaspidoproctus. Vegetable Weevil (see Listroderes obliquus).

Velsicol AR 60, effect of dusts containing, on Aphids, 143, 144; as solvent for DDT, 319.

Venezuela, Fundella argentina in, 104.

venipars, Myelois.

venosatus, Proceras (Diatraea).

ventrale, Stigmatium.

ventralis, Chrysopa.

ventricosus, Pediculoides (Pyemotes).

vepretella, Gelechia.

Veratridine, toxicity of, to insects, 302. Veratrine, toxicity of, to insects, 301, 302.

verbasci, Anthrenus.

Verbascum, Engytatus geniculatus on, in N. Carolina, 347.

vermiformis, Cecidophyes (Eriophyes).

vernalis, Tiphia.

vernata, Paleacrita.

Vernonia, as reservoir of Bemisia and virus of tobacco in Nyasaland, 206.

Veronica, Asterolecanium arabidis on, in California, 109.

verrillata, Alcis (see Thamnonoma ochrifascia).

vesicalis, Mordwilkoja.

Vesiculaphis caricis, Cerosipha subterranea (californica) recorded as, in Hawaii, 149.

vesiculata, Yangambia.

Vespa, control of species of, damaging fruits in France, 229.

Vespa crabro, 229.

Vespa germanica, 229; in beehive in New Zealand, 323.

Vespa vulgaris, 229.

vespiformis, Franklinothrips.

Vetches, Xerophloea vanduzeei on, in California, 227; eradication of, as source of Cydia nigricana in Washington, 76.

Viburnum, Diaspidiotus ancylus on, in California, 374.

Vicia faba (see Broad Beans).

vicinus, Camponotus maculatus; Scapteriscus.

Victoria, pests of stored cereals in, 21, 121, 230.

vigintiduopunctata, Thea.

vigintioctopunctata, Epilachna.

Vigna unguiculata (sinensis) (see Cowpeas).

Vigna unguiculata var. sesquipedalis (see Long Beans).

Vine, Grape, 145; Phylloxera vitifoliae on, in Argentina, 299; pests of, in Australia, 23, 364, 382; pests of, in France, 33, 328, 362; Agrilus derasofasciatus on, in Germany, 328; Anastrepha fraterculus not developing on, in Mexico, 138; Ceratitis capitata on, in Palestine, 132, 133; Pseudococcus citri on, in Persia, 16; Empoasca lybica on, in Spain, 267; pests of, in U.S.A., 52, 78, 109, 112, 227, 234, 245, 246, 247, 262; Euzophera bigella on, in Uzbekistan, 156; Cydia molesta experimentally developing on, 109.

Vine Phylloxera (see Phylloxera viti-

foliae).

vinitor, Nysius.

vinula, Dicranura.

Viola arvensis, Aphids on, in Holland, 367.

Viola odorata (Violet), Dasyneura affinis on, in Egypt, 18.

virens, Agromyza; Brachynema (see B. germari).

virescens, Heliothis (Chloridea); Temnochila.

virgata, Ferrisia (Ferrisiana).

Virginia, apple pests in, 4, 31, 105, 106, 195, 336; pests of leguminous plants in, 138, 244, 304, 316, 317; Cinara spp. on Pinus virginiana in, 311; pests of field and stored tobacco in, 128, 338; beneficial insects in, 305, 336, 337.

viridis, Agrilus; Coccus.

viridula, Nezara.

viridulus, Eulophus.

Virus Diseases, in Neodiprion sertifer, 130; (in plants*), relation of Aphids to, 70, 71, 74, 90, 99, 102, 113, 114, 182, 206, 226, 297, 367, 370; Aleurodids transmitting, 169, 206; Coccids transmitting, 46, 102; relation of Jassids to, 74, 111, 353, 354, 374, 379, 380; negative experiment with Psyllid and, 102; thrips transmitting, 206; mechanical transmission of, 70, 182, 380.

Viteus (see Phylloxera).

vitifoliae, Phylloxera (Viteus).

Vitis californica, Harrisina brillians on, in California, 112.

Vitis vinifera (see Vine, Grape).

vitis, Eriophyes.

vittata, Diabrotica (see D. melanocephala); Phyllotreta.

vittula, Phyllotreta.

Volck Oil, 52.

vorax, Anthrenus.

Voria ruralis, parasite of Anticarsia gemmatalis in Porto Rico, 219. vosseleri, Lygus (see L. simonyi).

vulgaris, Vespa.

vulneratus, Poeciloscytus.

vulpinus, Dermestes (see D. maculatus). vuteria, Sesamia.

^{*}These references are also indexed under the individual plants infected.

W.

Walnut, Caliroa limacina on, in W. Australia, 365; pests of, in California, 104, 145, 309, 374.

Walnut, Black (see Juglans nigra).
Walnut Aphis (see Chromaphis juglandicola).

Washington, Hylemyia cilicrura on seedling conifers in, 76, 160; orchard pests in, 50, 73, 74, 75, 175; pests of small fruits in, 75, 160; pests of vegetables in, 72, 73, 74, 75, 76, 140, 182; Sitodiplosis mosellana on wheat in, 251; wireworms in, 74; insects and virus diseases of plants in, 74, 182; beneficial insects and biological control in, 72, 73, 76, 251.

Wasmannia auropunctata, habits and control of, in Citrus groves in Florida,

238.

Wasps (see Vespa germanica and Vivulgaris).

Water, Hot, uses of, against pests, 78, 212.

Watermelon, Aulacophora chinensis on, in China, 344; Diabrotica on, in Indiana, 148; pulp of, in bait for Schistocerca gregaria, 165.

Wattle (see Acacia).

Wax Moth (see Galleria mellonella).

wesmaeli, Pristiphora (Lygaeonematus). West Indian Sugar-cane Mite (see

Tarsonemus bancrofti).

Western Red Cedar (see Thuja plicata). Wheat, 316; Cephus pygmaeus on, in Belgium, 185; pests of, in Britain, 46, 216, 330, 367, 368, 381; pests of, in Canada, 146, 174, 251, 319; Tipula paludosa on, in Denmark, 116; Sesamia vuteria on, in Morocco, 257; Rhopalosiphum prunifoliae on, in New South Wales, 385; Laphygma exempta on, in Nyasaland, 63; Anomala nazarena on, in Palestine, 17; Margarodes on, in Persia, 16; Eurygaster on, in Russia, 48, 131; Cecidomyiids on, in Sweden, 36; pests of, in Turkey, 48, 256; pests of, in U.S.A., 4, 5, 28, 74, 129, 136, 137, 146, 157, 177, 180, 181, 203, 251, 340, 375; varieties and time of sowing of, in relation to Mayetiola destructor, 4, 5, 28, 158, 177, 375; varieties of, resistant to Toxoptera graminum, 375. Wheat (Stored), pests in, and their control, 21, 22, 45, 65, 97, 120, 121, 122,

177, 188, 222, 223, 230, 296, 297, 325, 342, 385.
Wheat Bulb Fly (see *Hylemyia coarc*-

tata).

Wheat Gall-midges (see Contarinia tritici and Sitodiplosis mosellana).

Wheat Stem Sawfly (see Cephus cinctus). Whiteflies, transmitting plant virus diseases, 169, 206; natural enemies of, 99, 171; new species of, 64.

White-fringed Beetles (see *Pantomorus* (*Graphognathus*)).

wilkinsoni, Apanteles.

Willow (see Salix).

Wilt Disease, in Lepidoptera, 193, 379. Wimmera Rye-grass (see *Lolium rigidum*).

Wind, spread of insects by, **33, 36, 133, 184.**

Wireworms, influence of crops and soil factors on, 74, 92, 157, 235, 330; factors affecting passage of water through cuticle of, 60; naphthalene against, 74; method of removing, from soil samples, 329.

Wisconsin, miscellaneous pests in, 311, 312, 313; virus diseases of cabbage

in, 182.

Wistaria, Coccid on, in California, 109. wistariae, Chionaspis.

woglumi, Aleurocanthus.

Woodlice, tests of contact insecticides on, **255.** (See *Armadillidium*.)

Woodpecker, Green (see Gecinus viridus). Wool, pests in, **59**, **194**.

Woolly Apple Aphis (see Eriosoma lanigerum).

Wyethia mollis, preferred by Anabrus simplex in California, 25.

Wyoming, Anabrus simplex in, 177; potato pests in, 358.

X.

Xanthone, unsatisfactory in dusts against insects on beans, 316, 317. Xanthopimpla punctata, parasite of

Xanthopimpla punctata, parasite Cosmophila flava in China, 381.

xanthostigmus, Apanteles.

Xerophloea vanduzeei, bionomics of, causing disease-like symptoms in plants in California, 227, 228.

Xestobium rufovillosum, treatment of

wood against, 116.

Xiphydria prolongata, on Salix caprea in Finland, 41.

Xylaria, infesting Lachnosterna anxia in Montana, 181.

Xyleborus cryptographus, Cucujus cinnaberinus associated with, in aspen in Finland, 41.

Xyleborus eurygraphus, measures against, on Pinus pinaster in France, 371.

Xyleborus fornicatus fornicatior, factors affecting damage to tea by, in Ceylon, 71.

Xyleborus morstatti, on Crotalaria anagyroides in Mysore, 187.

Xylene (Xylol), as a solvent for DDT, 44, 244, 245, 314.

Xylion adustus, measures against, in timber in S. Africa, 12.

Xylocrius agassizi, bionomics and control of, on gooseberry in Oregon, 79.

Xylol (see Xylene).

Xyloperthodes nitidipennis, measures against, in timber in S. Africa, 12.

Xylotrechus quadripes, bionomics and control of, in Mysore, 186, 187.

Xylotrechus rusticus, on aspen and birch in Finland, 41.

Y.

Yam, damaged by Atta cephalotes in Br. Honduras, 323.

Yangambia vesiculata, gen. et sp. n., in Belgian Congo, 81; synonymy of, 81. Yeast, in baits for Lepidoptera, 308, 309, 372; in foods for insects, 62, 138, 308.

ypsilon, Agrotis.

Z.

Zabrus, crop rotation against, in U.S.S.R., 157.

Zacharops narangae, parasite of Cosmophila flava in China, 381.

Zadiprion, systematic position of, 14.

Zamia Palm (see Macrozamia).

Zantedeschia, experiment with Phenacoccus colemani and, 31.

Zatropis, parasite of Ceuthorrhynchus assimilis in Washington, 76.

zealandica, Odontria.

Zeiraphera fortunana, on spruce in Nova Scotia, 129.

Zeiraphera ratzeburgiana, on spruce in Nova Scotia, 129.

Zenillia milis, parasite of Taeniocampa gracilis in Finland, 42.

Zeuzera coffeae, on coffee in Mysore, 187.
Zeuzera pyrina, bionomics and control of, on fruit trees, etc., in England, 34.
Zicrona coerulea, in Belgium, 213;

Zicrona coerulea, in Belgium, 213; experimentally predacious on Leptinotarsa decemlineata, 213.

Zinc Arsenate, Basic, spraying with, against Rhagoletis pomonella, 50.

Zinc Arsenite, in sprays and dusts against insects on potato, 29, 224, 358, 359; increasing infestation by Myzus persicae and Aceratagallia uhleri, 224.

Zinc Chloride, sap-stream treatment of trees with, as timber preservative, **56**; and sodium dichromate, **56**.

Zinc Fluoride, protecting wood from termites, 172.

Zinc Metarsenite, sap-stream treatment of trees with, as timber preservative, 56.

Zinc Oxide, reducing effect on plants of calcium arsenate in soil, 355.

Zinc Phosphide, in bait for molecrickets, 19.

Zinc Sulphate, use of, with arsenates in sprays, 50, 51, 196, 197, 199, 201, 202, 203, 204, 309, 336.

zinckenella, Etiella.

Zinnia, Hispid on, in Arizona, 27; turnip viruses infecting, 182.

Zinnia elegans, Agalliana ensigera on, in Argentina, 380; A. ensigera transmitting curly-top virus to, 379.

Zizania latifolia, Pachydiplosis oryzae ovipositing on, in China, 119.

zizanialis, Protrigonia. ziziphus, Parlatoria.

Zizyphus, Parlatoria morrisoni on, in Persia, 16.

Zizyphus jujuba, Cydia molesta experimentally developing on, 109.

zonellus, Chilo.

Zonocerus variegatus, Empusa grylli infesting, in Belgian Congo, 280.

and the state of t

NOTICES.

Secretaries of Societies and Editors of Journals willing to exchange their publications with those of the Institute are requested to communicate with the Director.